



## **Ethiopian Oribatid Mites of the Family Scheloribatidae (Acari: Oribatida)**

Authors: Ermilov, Sergey G., Rybalov, Leonid B., and Franke, Kerstin

Source: African Invertebrates, 52(2) : 311-322

Published By: KwaZulu-Natal Museum

URL: <https://doi.org/10.5733/afin.052.0207>

---

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Ethiopian oribatid mites of the family Scheloribatidae (Acari: Oribatida)

Sergey G. Ermilov<sup>1\*</sup>, Leonid B. Rybalov<sup>2</sup> and Kerstin Franke<sup>3</sup>

<sup>1</sup>Laboratory of Entomology, Phytosanitary Department, Nizhniy Novgorod Referral Center of the Federal Service for Veterinary and Phytosanitary Inspection, prospekt Gagarina 97, Nizhniy Novgorod, 603107 Russia; ErmilovAcari@yandex.ru

<sup>2</sup>Laboratory for Soil Zoology and General Entomology, Institute of Ecological and Evolutionary Problems, Russian Academy of Sciences, Leninskiy prospekt 33, Moscow, 117071 Russia; lrybalov52@mail.ru

<sup>3</sup>Section Arachnida, Senckenberg Museum of Natural History Goerlitz, Am Museum 1, Goerlitz, 02826 Germany; Kerstin.Franke@senckenberg.de

\*Corresponding author

### ABSTRACT

We have registered eight species and three genera (*Scheloribates* Berlese, 1908, *Similobates* Mahunka, 1982, *Perscheloribates* Hammer, 1973) of Scheloribatidae from southern Ethiopia. Two species of *Scheloribates*—*S. discifer* and *S. latipes*—and the genus *Perscheloribates* are reported from Ethiopia for the first time. Two new species, *Scheloribates acutirostrum* sp. n. and *Perscheloribates crassisetosus* sp. n. from Cholomu and Harennna forests, are described. The first is from soil, litter and mosses on trees, while the second is from litter. Identification keys to Ethiopian species of *Scheloribates* and African species of *Perscheloribates* are presented.

KEY WORDS: Oribatida, *Scheloribates*, *Perscheloribates*, mites, new species, new records, identification keys, Afrotropical, Ethiopia.

### INTRODUCTION

Until recently, the oribatid mites of Ethiopia have been poorly studied (Berlese 1916; Aoki 1971; Mahunka 1982, 1983, 1984; Bernini 1988). However, investigations of these oribatids have increased, e.g. with the works of Niedbała (2008), Ermilov *et al.* (2010a–h, 2011, in press), and Niedbała & Ermilov (2011). The present work is part of a continuing study of the Ethiopian oribatid mite fauna, and focuses on species in the family Scheloribatidae. Worldwide, this family comprises 20 genera and 338 species that collectively have a cosmopolitan distribution (Subías 2004, online version 2011).

Only two species and two genera of Scheloribatidae have been reported previously from Ethiopia: *Scheloribates aethiopicus* Mahunka, 1982 and *Similobates demetorum* Mahunka, 1982. In the course of our taxonomic studies we have registered eight species and three genera (*Scheloribates* Berlese, 1908, *Similobates* Mahunka, 1982 and *Perscheloribates* Hammer, 1973) of Scheloribatidae from southern Ethiopia. We found representatives of two new species, one belonging to *Scheloribates* and another to *Perscheloribates*. The other identified scheloribatid taxa are presented below.

*Scheloribates* was proposed by Berlese (1908) with *Zetes latipes* Koch, 1844 as the type species. Currently, it comprises 221 species and is distributed worldwide (Subías 2004, online version 2011). The diagnostic characters of the genus *Scheloribates* are as following: lamellae well developed; prolamellae present, absent or partially developed; translamellar lines usually absent, but rarely incompletely present; sensilli with dilated head; pteromorphs immovable; notogaster with 10 pairs of setae or their alveoli; four or five pairs of sacculi on notogaster; four pairs of genital setae; legs tridactylous (e.g., Coetzer 1967–1968; Corpuz-Raros 1980; Lee & Pajak 1990).

*Perschelorbates* was proposed by Hammer (1973) with *Perschelorbates clavatus* Hammer, 1973 as the type species. Currently, it comprises 41 species and have pan-tropical and subtropical distribution (Subías 2004, online version 2011). The diagnostic characters of *Perschelorbates* are as following: rostrum rounded; lamellae well developed; prolamellae present, absent or partially developed; translamellar lines usually absent, but rarely incompletely present; interlamellar setae long; sensilli with dilated head; pteromorphs immovable; notogaster with 10 pairs of setae or their alveoli; four pairs of sacculi on notogaster; four pairs of genital setae; lyrifissures *iad* adanal; legs monodactylous (e.g., Hammer 1973; Corpuz-Raros 1980).

#### MATERIAL AND METHODS

Samples (soil, mosses, lichens *etc.*) were processed in Tullgren funnels if another method is not specified.

Specimens of new species were studied in lactic acid, mounted on temporary cavity slides for the duration of the study, then were stored in 70% alcohol. All body measurements are presented in micrometres. Body length was measured in lateral view, from the tip of the rostrum to the posterior edge of the ventral plate, to avoid discrepancies caused by different degrees of notogastral distension. Notogastral width refers to the maximum width in dorsal aspect. Length of body setae was measured in lateral aspect.

Formulae of leg setation are given according to the sequence trochanter–femur–genu–tibia–tarsus (famulus included). Formulae of leg solenidia are given (in square brackets) according to the sequence genu–tibia–tarsus.

The holotype material is deposited at the Zoological Institute of the Russian Academy of Sciences, St Petersburg, Russia (ZISP). The paratype material is deposited at the Siberian Zoological Museum, Novosibirsk, Russia (SZMN) and is in the personal collection of the first author (PC).

#### List of collecting sites

S-1: 8°53'N:38°09'E, 2900 m, 10 km south of Ginchi city, Cholomu Forest (woody species, in particular *Hagenia abyssinica* and *Podocarpus* forming the canopy; undergrowth of ferns), in mosses on trees, 15.xi.2009, L.B. Rybalov.

S-2: 6°42'N:39°43'E, 2249 m, Bale Mountains National Park, Harenna Forest (woody species, *H. abyssinica* forming the canopy), in mosses on trees, 23.xi.2009, L.B. Rybalov.

S-3: 6°49'N:39°49'E, 4367 m, Bale Mountains National Park, Sanetti plateau, Batu mountain peak, plants *Artemisia* spp. and *Oxytriops* spp., 23.xi.2009, L.B. Rybalov.

S-4: 6°42'N:39°43'E, 4050 m, Bale Mountains National Park, Sanetti plateau, Afro-Alpine tundra, mosses and soil near a bog, 23.xi.2009, L.B. Rybalov.

S-5: 6°38'N:39°43'E, 1883 m, Bale Mountains National Park, Harenna Forest (woody species, in particular *H. abyssinica* forming the canopy), in soil, 23.xi.2009, L.B. Rybalov.

S-6: 6°42'N:39°43'E, 2249 m, Bale Mountains National Park, Harenna Forest (woody species, in particular *H. abyssinica* forming the canopy), in soil, 23.xi.2009, L.B. Rybalov.

S-7: 8°53'N:38°09'E, 2920 m, 10 km south of Ginchi city, Wenchi crater, Cholomu Forest (*H. abyssinica* forming the canopy), plants on rock, 20.xi.2010, L.B. Rybalov & A.I. Bastrakov.

S-8: 8°53'N:38°09'E, 2810 m, 10 km south of Ginchi city, Cholomu Forest (*H. abyssinica* forming the canopy), in litter, 28.xi.2010, L.B. Rybalov & A.I. Bastrakov.

S-9: 8°53'N:38°09'E, 3300 m, 10 km south of Ginchi city, Wenchi crater, Cholomu Forest (*Erica* forming the canopy), plant mosses, 20.xi.2010, L.B. Rybalov & A.I. Bastrakov.

S-10: 8°53'N:38°09'E, 2810 m, 10 km south of Ginchi city, Cholomu Forest (*H. abissinica* forming the canopy), sifting filter, 04.xi.2010, L.B. Rybalov & A.I. Bastrakov.

S-11: same as S-10, but lichens on trees.

S-12: same as S-10, but leaf litter.

S-13: same as S-10, but mosses on trees.

S-14: same as S-10, but in soil.

S-15: same as S-10, but mosses and orchids on trees.

#### TAXONOMY

##### Family Scheloribatidae Grandjean, 1933

We found representatives of three genera and eight species of the family, as follow:

*Scheloribates acutirostrum* sp. n. Localities: S-1, S-5, S-12.

*S. aethiopicus* Mahunka, 1982. Localities: S-1, S-2, S-3, S-4, S-5, S-6, S-9, S-10, S-13, S-14, S-15.

*S. discifer* Balogh, 1959. Localities: S-3, S-4, S-5, S-6, S-8, S-10. First record for Ethiopia.

*S. latipes* (Koch, 1844). Localities: S-1, S-5, S-6, S-7, S-8, S-9, S-10, S-11, S-12. First record for Ethiopia.

*Similobates demetorum* Mahunka, 1982. Localities: S-5, S-8, S-10.

*Perscheloribates crassisetosus* sp. n. Localities: S-8, S-10.

*P. ? luminum* (Hammer, 1961). Locality: S-10. First record for Ethiopia. Our specimens are similar to the description of *P. luminum* in all characters. However Hammer (1961) does not give information about the prolamella, therefore we cannot make a definitive identification (our specimens lack a prolamella). We also have no identified material of *P. luminum* for comparison. Thus, we tentatively name our species *P. luminum*.

*P. minutus* (Pletzen, 1965). Localities: S-1, S-5. First record for Ethiopia.

##### Genus *Scheloribates* Berlese, 1908

##### ***Scheloribates acutirostrum* sp. n.**

##### Figs 1–3

**Etymology:** From Latin *acutus* (point) and rostrum, referring to the pointed rostrum.

**Diagnosis:** This species is distinguished by the following combination of character states: body size 614–697 × 448–481; dorsal and ventral surfaces of body smooth; rostrum pointed; translamellar line present, thin, interrupted medially; sensilli spindle-form, barbed, with short apical part; one pair of notogastral setae and nine pairs of notogastral alveoli; prolamellae absent; tarsi I each with 19 setae.

##### Description:

**Measurements.** Body length 647 (holotype ♂), 614–697 (paratypes: 3♂, 1♀); body width 481 (holotype ♂), 448–481 (paratypes: 3♂, 1♀).

**Integument.** Body brown. Dorsal and ventral surfaces of body smooth. Lateral surfaces of body weakly granulate.

**Prodorsum** (Figs 1A; 2A–C). Rostrum pointed in dorsoventral view. Lamellae more than half of prodorsum. Translamellar line present; thin, interrupted medially. Rostral (*ro*,




Fig. 1. *Schelorbates acutirostrum* sp. n.: (A) dorsal view, legs not shown; (B) ventral view, legs, palps and subcapitular setae not shown. Scale bar = 200  $\mu$ m.

94–102), lamellar (*le*, 155–172) and interlamellar (*in*, 196–229) setae setiform, barbed. Sensilli (*ss*, 102–116) spindle-form, barbed; apical part short. Lateral region (Fig. 2B) with prolamella absent but sublamellar line present. Sublamellar areae porosae oblong or oval (*Al*, 12–16  $\times$  4–8). Exobothridial setae (*ex*, 12) thin, slightly barbed.

*Notogaster* (Fig. 1A). Dorsosejugal suture weakly convex medially. Only one pair of thin, smooth notogastral setae (*p*<sub>1</sub>, 12) present; other nine pairs represented only by alveoli. Sacculi *Sa* and *S1* oblong, distinct; *S2* and *S3* very small, inconspicuous. Opisthonotal gland opening and lyrifissures inconspicuous (except *im*) but present in typical arrangement for genus.

*Anogenital region* (Figs 1B; 2D–F). Two pairs anal (*an*<sub>1</sub>, *an*<sub>2</sub>, 20–24), three pairs adanal (*ad*<sub>1</sub>–*ad*<sub>3</sub>, 24), one pair aggenital (*ag*, 12–16) and four pairs of genital (*g*<sub>1</sub>, 24–28; *g*<sub>2</sub>–*g*<sub>4</sub>, 16–20) setae setiform, smooth. Lyrifissures *iad* in typical position for genus. Ovipositor: length of lobes 86–90, length of cylindrical distal part (bDp) 98–110, width of cylindrical distal part 41–49. All setae of ovipositor setiform, smooth. Lobe setae  $\psi_1 \approx \tau_1$  (36–41) longer than  $\psi_2 \approx \tau_a \approx \tau_b \approx \tau_c$  (16–20). Coronal setae absent.

*Epimeral region* (Fig. 1B). Apodemes 1, 2, sejugal and circumpedal carina well-developed. Apodemes 3 present, but weakly visible. All epimeral setae (16–24) setiform, smooth.

*Gnathosoma* (Figs 3A–C). Subcapitulum longer than wide: 118–127  $\times$  106–123. Hypostomal setae setiform, weakly and indistinctly barbed; *h* (41) longer than *m* (32) and *a* (24). Lateral lips with two pairs of adoral setae (*or*<sub>1</sub>, *or*<sub>2</sub>, 12–16), setiform, barbed. Palps (length 82–102) with setation 0–2–1–3–9(+1 $\omega$ ). All setae (except some on tarsi) weakly and indistinctly barbed. Chelicerae (length 151–164) chelate-dentate; cheliceral setae setiform, barbed, *cha* (57–65) longer than *chb* (32–41).




Fig. 2. *Schelorbates acutirostrum* sp. n.: (A) rostrum; (B) lateral view of prodorsum, gnathosoma and legs not shown; (C) sensillus; (D) genital plate, right; (E) anal plate, right, and adanal setae; (F) ovipositor. Scale bars A, C, E, F = 50  $\mu$ m; B = 100  $\mu$ m; D = 20  $\mu$ m.

**Legs** (Figs 3D, 3E). Morphology of segments similar to other species of *Schelorbates* (e.g., Bayartogtokh 2000). Formulae of leg setation and solenidia: I (1-5-3-4-19) [1-2-2], II (1-5-2-4-15) [1-1-2], III (2-3-1-3-15) [1-1-0], IV (1-2-2-3-12) [0-1-0]; homology of setae and solenidia indicated in Table 1. Almost all setae barbed. Famulus short, straight, blunt-ended. Solenidia  $\omega_1$  on tarsi I,  $\omega_1$  and  $\omega_2$  on tarsi II, rod-shaped; other solenidia setiform.




Fig. 3. *Scheloribates acutirostrum* sp. n.: (A) subcapitulum; (B) palp; (C) chelicera; (D) leg I, right, paraxial view; (E) leg IV, right, paraxial view. Scale bars A, C = 50  $\mu$ m; B = 20  $\mu$ m; D, E = 100  $\mu$ m.

Holotype: ♂ ETHIOPIA: Locality S-1 (ZISP).

Paratypes: ETHIOPIA: Localities S-1 (1 ♀ PC), S-5 (2 ♂ SZMN), S-12 (1 ♂ SZMN).

Distribution: At present, this species is known only from Ethiopia.

Remarks: In having the combination of a pointed rostrum, spindle-form sensilli and prolamella absent *S. acutirostrum* sp. n. is similar to *S. sphaeroides* Hammer, 1973 from Polynesia and the Galapagos Islands (Hammer 1973). It differs from the latter in the relative length of interlamellar and lamellar setae (*in* longer than *le* in *S. acutirostrum*; *in* not longer than *le* in *S. sphaeroides*), the shape of the sensilli (head well developed, barbed and with short apical part in *S. acutirostrum*; head weakly developed, smooth and with long apical part in *S. sphaeroides*) and the presence of only notogastral setae  $p_1$  ( $p_1, h_1-h_3$  present in *S. sphaeroides*). In having the combination of a pointed rostrum and spindle-form sensilli, *S. acutirostrum* sp. n. is similar to *S. rostrodentatus* Hammer, 1977 from Pakistan (Hammer 1977), but differs from the latter by having smaller body (614–697 vs 980 in *S. rostrodentatus*) and in the absence of a prolamella (present in *S. rostrodentatus*).



TABLE 1

Leg setation and solenidia of *Scheloribates acutirostrum* sp. n. Roman letters refer to normal setae (*e* – *fa* – *fulmus*), Greek letters refer to solenidia. One apostrophe (') marks setae on anterior, double apostrophe (") setae on posterior, side of the given leg segment. Parentheses refer to a pair of setae.

Leg	Trochanter	Femur	Genu	Tibia	Tarsus
I	v'	<i>d</i> , ( <i>l</i> ), <i>bv</i> "', v"	( <i>l</i> ), v', σ	( <i>l</i> ), ( <i>v</i> ), φ <sub>1</sub> , φ <sub>2</sub>	( <i>ft</i> ), ( <i>tc</i> ), ( <i>it</i> ), ( <i>p</i> ), ( <i>u</i> ), ( <i>a</i> ), <i>s</i> , ( <i>pv</i> ), v', ( <i>pl</i> ), <i>e</i> , ω <sub>1</sub> , ω <sub>2</sub>
II	v'	<i>d</i> , <i>l</i> ' <sub>1</sub> , <i>l</i> ' <sub>2</sub> , <i>bv</i> "', v"	( <i>l</i> ), σ	( <i>l</i> ), ( <i>v</i> ), φ	( <i>ft</i> ), ( <i>tc</i> ), ( <i>it</i> ), ( <i>p</i> ), ( <i>u</i> ), ( <i>a</i> ), <i>s</i> , ( <i>pv</i> ), ω <sub>1</sub> , ω <sub>2</sub>
III	v', <i>l</i> '	<i>d</i> , <i>l</i> ', <i>ev</i> '	<i>l</i> ' σ	<i>l</i> ' ( <i>v</i> ), φ	( <i>ft</i> ), ( <i>tc</i> ), ( <i>it</i> ), ( <i>p</i> ), ( <i>u</i> ), ( <i>a</i> ), <i>s</i> , ( <i>pv</i> )
IV	v'	<i>d</i> , <i>ev</i> '	<i>d</i> , <i>l</i> '	<i>l</i> ' ( <i>v</i> ), φ	<i>ft</i> "', ( <i>tc</i> ), ( <i>p</i> ), ( <i>u</i> ), ( <i>a</i> ), <i>s</i> , ( <i>pv</i> )

Key to Ethiopian species of *Scheloribates*

- 1 Rostrum pointed; sensilli spindle-form.....**acutirostrum** sp. n.
- Rostrum with rounded apex; sensilli with rounded head distally .....2
- 2 Translamellar line present, interrupted medially; all notogastral setae developed ..... **aethiopicus** Mahunka, 1982
- Translamellar line absent; some notogastral setae presented by alveoli.....3
- 3 Sensilli very short (length of head slightly longer than diameter of bothridia), with oval head ..... **discifer** Balogh, 1959
- Sensilli long (length of head considerably longer than diameter of bothridia), with noticeably oblong head ..... **latipes** Koch, 1844

Genus *Perscheloribates* Hammer, 1973

***Perscheloribates crassisetosus*** sp. n.

Figs 4–6

**Etymology:** From Latin *crassus* (thick) and *seta*, referring to the thickened lamellar setae.

**Diagnosis:** This species is distinguished by the following combination of character states: body size 481–514 × 348–365; translamellar line present, interrupted medially; lamellar setae thickened; sensilli long, with asymmetrically dilated, barbed head; one pair of muscle sigilla present posterior to each saccule *Sa*, three pairs of muscle sigilla present posterior to each saccule *SI*; prolamella present; coronal setae of ovipositor present; hypostomal setae *a* thicker than others, setae *m* very short; tarsi I with 18 setae.

**Description:**

**Measurements.** Body length 481 (holotype ♂), 514 (paratype ♀); body width 348 (holotype ♂), 365 (paratype ♀).

**Integument.** Body brown. Dorsal and ventral surfaces of body smooth. Lateral surfaces of body weakly granulate.

**Prodorsum** (Figs 4A; 5A–C). Rostrum rounded in dorsal view. Lamellae more than half length of prodorsum. Translamellar line present; thin, interrupted medially. Rostral (57–65) and interlamellar (151–159) setae setiform, barbed. Lamellar setae (155–172) unusually thickened in basal half, ciliate. Sensilli long (94), with asymmetrically dilated head; head shorter than stalk, slightly barbed. Lateral region (Fig. 5A) with prolamella present and sublamellar line long. Sublamellar areae porosae oblong or oval (8–12 ×






Fig. 4. *Perscheloribates crassisetosus* sp. n.: (A) dorsal view, legs not shown; (B) ventral view, legs, palps and subcapitular setae not shown. Scale bar = 200  $\mu$ m.

4–6). Lateral carina (*kf*) as typical for many Scheloribatidae species, but part parallel to sublamellar line longer, more prominent. Exobothridial setae (*ex*, 8) thin, slightly barbed.

*Notogaster* (Fig. 4A). Dorsosejugal suture weakly convex medially. Only one pair of thin, smooth notogastral setae present ( $p_1$ , 16); nine other pairs presented by alveoli. All sacculi oblong, distinct. One pair of muscle sigilla (not sacculi) present posterior to each saccule *Sa*. Three pairs of muscle sigilla present posterior to each saccule *S1*. Circumgastric band of muscle sigilla distinct. Opisthonotal gland opening and lyrifissures indistinct, but developed in typical arrangement for genus.

*Anogenital region* (Figs 4B; 5D–F). Two pairs anal (8–12), three pairs adanal (20), one pair aggenital (8–12) and four pairs of genital ( $g_1$ , 20;  $g_2$ – $g_4$ , 8–12) setae setiform, smooth. Lyrifissures *iad* in typical position for genus. Ovipositor: length of lobes 86, length of cylindrical distal part 96, width of cylindrical distal part 32. All setae of ovipositor setiform, smooth. Lobe setae  $\psi_1 \approx \tau_1$  (24) longer than  $\psi_2 \approx \tau_a \approx \tau_b \approx \tau_c$  (14). Coronal setae present (*k*, 8).

*Epimeral region* (Fig. 4B). Apodemes 1, 2, sejugal, 3 and circumpedal carina well-developed. All epimeral setae (12–20) setiform, smooth.

*Gnathosoma* (Figs 6A–C). Subcapitulum longer than wide: 118  $\times$  102. Hypostomal setae setiform; *a* (26) slightly longer and thicker than *h* (24), both hardly barbed, setae *m* very short (8), smooth. Lateral lips with two pairs of adoral setae (12), setiform, barbed. Palps (length 73) with setation 0–2–1–3–9(+1 $\omega$ ). All setae (except some on tarsi) weakly and indistinctly barbed. Chelicerae (length 131) chelate-dentate; cheliceral setae setiform, barbed, *cha* (61) longer than *chb* (32).




Fig. 5. *Perschelorbates crassisetosus* sp. n.: (A) lateral view of prodorsum, gnathosoma and legs not shown; (B) lamellar seta; (C) sensillus; (D) genital plate, right; (E) anal plate, right, and adanal setae; (F) ovipositor. Scale bars A = 100 μm; B–F = 50 μm.

*Legs* (Figs 6D, 6E). Formulae of leg setation and solenidia: I (1–5–3–4–18) [1–2–2], II (1–5–2–4–15) [1–1–2], III (2–3–1–3–15) [1–1–0], IV (1–2–2–3–12) [0–1–0]; homology of setae and solenidia indicated in Table 2. Almost all setae barbed. Famulus short, straight, blunt-ended. Solenidia ω<sub>1</sub> on tarsi I, ω<sub>1</sub> and ω<sub>2</sub> on tarsi II, rod-shaped; other solenidia setiform.

Holotype: ♂ ETHIOPIA: Locality S-8 (ZISP).

Paratype: ETHIOPIA: ♀ Locality S-10 (PC).

Distribution: At present, this species is known only from Ethiopia.

TABLE 2

Leg setation and solenidia of *Perscheloribates crassisetosus* sp. n. For explanation see Table 1.

Leg	Trochanter	Femur	Genu	Tibia	Tarsus
I	v'	d, (l), bv'', v''	(l), v', σ	(l), (v), φ <sub>1</sub> , φ <sub>2</sub>	(fi), (tc), (it), (p), (u), (a), s, (pv), (pl), e, ω <sub>1</sub> , ω <sub>2</sub>
II	v'	d, l', l' <sub>2</sub> , bv'', v''	(l), σ	(l), (v), φ	(fi), (tc), (it), (p), (u), (a), s, (pv), ω <sub>1</sub> , ω <sub>2</sub>
III	v', l'	d, l', ev'	l', σ	l', (v), φ	(fi), (tc), (it), (p), (u), (a), s, (pv)
IV	v'	d, ev'	d, l'	l', (v), φ	fi'', (tc), (p), (u), (a), s, (pv)

Remarks: In having the combination of long sensilli, with asymmetrically dilated and slightly barbed head, and the presence of translamellar line (interrupted medially), *P. crassisetosus* sp. n. is very similar to *P. luminosus* from the tropics (Hammer 1961); however, it clearly differs from the latter by having a larger body (481–514 vs 420 in *P. luminosus*) and thicker lamellar setae (thin in *P. luminosus*).




Fig. 6. *Perscheloribates crassisetosus* sp. n.: (A) subcapitulum; (B) palp; (C) chelicera; (D) leg I, right, antiaxial view; (E) leg IV, right, antiaxial view. Scale bars A, C–E = 50 μm; B = 20 μm.

Key to African species of *Perscheloribates*

- 1 Translamellar line absent ..... 2  
 – Translamellar line always present (interrupted medially) ..... 4
- 2 Aggenital setae absent ..... **ethiopicus** (Mahunka, 1986)  
 – Aggenital setae present ..... 3
- 3 All notogastral setae present, considerably longer than diameter of bothridia .....  
 ..... **shiraensis** (Evans, 1953)  
 – Dorsal notogastral setae absent (only alveoli present), caudal setae not longer than  
 diameter of bothridia ..... **rustenburgensis** (Pletzen, 1963)
- 4 Prolamella short, not reaching insertion of rostral setae .....  
 ..... **minimus** Mahunka, 1992  
 – Prolamella long, reaching insertion of rostral setae or absent ..... 5
- 5 Sensilli spindle-form ..... 6  
 – Sensilli lanceolate ..... 7
- 6 Sensillar head barbed; notogastral setae  $p_1$ – $p_3$  present .... **minutus** (Pletzen, 1965)  
 – Sensillar head smooth; only notogastral setae  $p_1$  present .....  
 ..... **tzitzikamaensis** (Pletzen, 1963)
- 7 Lamellar setae clear thicker than rostral and interlamellar setae .....  
 ..... **crassisetosus** sp. n.  
 – Lamellar setae not thickened ..... **luminosus** (Hammer, 1961)

## ACKNOWLEDGEMENTS

We gratefully acknowledge Prof. Roy A. Norton (State University of New York, College of Environmental Science and Forestry, Syracuse, USA), Dr Umukusum Shtanchaeva (Caspian Institute of Biological Resources, Makhachkala, Russia), Prof. Luis Subías (Universidad Complutense de Madrid, Madrid, Spain), Dr Matthew Colloff (CSIRO Entomology, Canberra, Australia), for help with collecting literature. We gratefully acknowledge Dr Ekaterina A. Sidorchuk (Paleontological Institute, Russian Academy of Sciences, Moscow, Russia) and A.I. Bastrakov (Institute of Ecological and Evolutionary Problems, Russian Academy of Sciences, Moscow, Russia) for help with collecting Ethiopian oribatid mites. The work was performed within the framework of the Joint Russian-Ethiopian Biological Expedition, financially supported by the Russian Academy of Sciences. We are grateful to our Project Coordinators Dr Andrey Darkov and Ato Girma Yosef for management of the Expedition. We thank Dr Kemal Ali, director of the Ambo Plant Protection Research Centre, EIAR, for supporting field studies and organizing laboratory operations. We thank Prof. Roy A. Norton and an anonymous reviewer for valuable comments.

## REFERENCES

- AOKI, J. 1971. A new species of oribatid mite, *Thamnacarus moribei*, from the West Coast of the Red Sea, with records of two other species of the family Lohmanniidae. *The Japanese Journal of Zoology* **16** (3): 127–129.
- BALOGH, J. 1959. Oribates (Acari) nouveaux d'Angola et du Congo Belge (1<sup>ère</sup> série). *Companhia de Diamantes de Angola, Lisboa* **48**: 91–108.
- BAYARTOGTOKH, B. 2000. Oribatid mites of the genus *Scheloribates* (Acari: Oribatida: Scheloribatidae) from Mongolia. *Edaphologia* **65**: 61–88.
- BERLESE, A. 1908. Elenco di generi e specie nuovi di Acari. *Redia* **5**: 1–15.
- 1916. Centuria seconda di Acari nuovi. *Redia* **12**: 125–177.
- BERNINI, F. 1988. A new species of *Basilobelba* Balogh, 1958 (Acarida Oribatida) from Ethiopia. *Tropical Zoology* **1**: 223–232.
- COETZER, A. 1967–1968. New Orbatulidae Thor, 1929 (Oribatei, Acari) from South Africa, new combinations and a key to the genera of the family. *Memórias do Instituto de Investigação Científica de Moçambique* **9** (A): 15–126.

- CORPUZ-RAROS, L. 1980. Philippine Oribatei (Acarina) V. *Schelorbitates* Berlese and related genera (Oribatulidae). *Kalikasan* **9** (2–3): 169–245.
- ERMILOV, S.G., SIDORCHUK, E.A. & RYBALOV, L.B. 2010a. Two new species of oribatid mites (Acari, Oribatida) from Ethiopia. *Annales Zoologici* **60** (3): 407–417.
- 2010b. Two new species of *Austrocarabodes* (*Uluguroides*) from Ethiopia (Acari: Oribatida: Carabodidae). *Annales Zoologici* **60** (4): 617–626.
- 2010c. Two new species of oribatid mites of the family Carabodidae from Ethiopia (Acari: Oribatida). *Genus* **21** (4): 659–671.
- 2010d. New oribatid mites of the genera *Plasmobates* and *Arcoppia* (Acari: Oribatida) from Ethiopia. *Genus* **21** (4): 673–686.
- 2010e. A new species of *Dolicheremaeus* (Acari: Oribatida: Tetracondylidae) from Ethiopia. *Systematic and Applied Acarology* **15** (3): 235–243.
- 2010f. A new species of the genus *Pergalumna* (Acari: Oribatida: Galumnidae) collected in moss on trees from Ethiopia. *Systematic and Applied Acarology* **15** (3): 244–250.
- 2010g. New species of oribatid mites of the superfamily Galumnoidea (Acari: Oribatida) from Ethiopia. *Zootaxa* **2646**: 43–62.
- 2010h. *Zetorchella nortoni*, a new species of oribatid mite from Ethiopia (Acari: Oribatida: Caloppiidae). *Acarina* **18** (1): 61–65.
- 2011. A new species of *Separatoppia* (Acari: Oribatida: Oppiidae) from Ethiopia. *Systematic and Applied Acarology* **16** (1): 21–26.
- In press. Three new species of oribatid mites (Acari: Oribatida: Galumnidae) from Ethiopia. *International Journal of Acarology*.
- EVANS, G.O. 1953. On a collection of Acari from Kilimanjaro (Tanganyika). *Annals and Magazine of Natural History* **12** (6): 258–281.
- GRANDJEAN, F. 1933. Étude sur le développement des Oribates. *Bulletin de la Société zoologique de France* **58**: 30–61.
- HAMMER, M. 1961. Investigations on the oribatid fauna of the Andes Mountains. *Det Kongelige Danske Videnskaberne Selskab Biologiske Skrifter* **13** (1): 1–151.
- 1973. Oribatids from Tongatapu and Eua, the Tonga Islands, and from Upolu, Western Samoa. *Det Kongelige Danske Videnskaberne Selskab Biologiske Skrifter* **20** (3): 1–70.
- 1977. Investigations on the oribatid fauna of North-West Pakistan. *Det Kongelige Danske Videnskaberne Selskab Biologiske Skrifter* **21** (4): 1–71.
- KOCH, C.L. 1835–1844. *Deutschlands Crustaceen, Myriapoden und Arachniden*. Ein Beitrag zur deutschen Fauna. Regensburg: Friedrich Pustet. (original not seen)
- LEE, C.L. & PAJAK, G.A. 1990. *Schelorbitates* Berlese and *Megaschelorbitates* gen. nov. from Southeastern Australia, with comments on Schelorbitatidae (Acari: Cryptostigmata: Oripodoidea). *Invertebrate Taxonomy* **90** (4): 206–246.
- MAHUNKA, S. 1982. Oribatids from the Eastern Part of the Ethiopian Region (Acari) I. *Acta Zoologica Academiae Scientiarum Hungaricae* **28** (3–4): 293–336.
- 1983. Oribatids from the Eastern Part of the Ethiopian Region II. *Acta Zoologica Academiae Scientiarum Hungaricae* **29** (1–3): 151–180.
- 1984. Oribatids of the Eastern Part of the Ethiopian Region (Acari). V. *Acta Zoologica Hungarica* **30** (1–2): 87–136.
- 1986. Oribatids from Africa (Acari, Oribatida), IV. *Annales Historico-Naturales Musei Nationalis Hungarici* **78**: 301–317.
- 1992. New and interesting mites from the Geneva Museum LXIII. A survey of the oribatid mites fauna of Senegal (Acari: Oribatida). *Revue Suisse de Zoologie* **99** (3): 673–712.
- NIEDBALA, W. 2008. Description of a new species of ptyctimous mites (Acari, Oribatida) from Ethiopia and a checklist of ptyctimous mites of the Afrotropical Region. *Tropical Zoology* **21**: 1–9.
- NIEDBALA, W. & ERMILOV, S.G. 2011. New and little known species of ptyctimous mites (Acari, Oribatida) from Ethiopia. *Zootaxa* **2739**: 60–68.
- PLETZEN, R. 1963. Studies on South African oribatei (Acarina). *Acarologia* **5** (4): 690–703.
- 1965. Studies on the South African oribatei (Acari). III. Further new species of the genus *Schelorbitates* Berlese, 1908. *Acarologia* **17** (1): 113–120.
- SUBÍAS, L.S. 2004. Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo (excepto fósiles). *Graellsia* **60** (número extraordinario): 3–305. (<http://www.ucm.es/info/zoo/Artropodos/Catalogo.pdf>; accessed in February 2011)