

Aberia caffra Hook.f. & Harv., see *Dovyalis caffra* (Hook.f. & Harv.) Warb.

Acanthocereus pentagonus (L.) Britt. & Rose

FAMILY: Cactaceae

COMMON NAME: barbwire acanthocereus, cactus, dildoe

NATIVE: S. Texas, Mexico

LABORATORY INFESTATION:

Keck & Marshall 1930: fruits were placed in cages and exposed to flies; 4-X-1929, firm, ripe fruits were exposed to fruit flies at 0900 hours; by 1400 hours, 16 oviposition punctures were observed; 9-X-1929, 100 larvae and many unhatched eggs were observed.

LISTING ONLY:

Chilson 1960; DeWoskin 1981; Mitchell et al. 1977; Weems 1981.

Acanthocereus tetragonus (L.) Hummelinck, see *A. pentagonus* (L.) Britt. & Rose

Achras zapota L., see *Manilkara zapota* (L.) Van Royen

Acmena smithii (Poir.) Merrill & L. M. Perry

FAMILY: Myrtaceae

COMMON NAME: lillypilly

NATIVE: Australia

LISTING ONLY:

*Jenkins 1950; *Jenkins & Shedly 1959.

Acokanthera longiflora Stapf

FAMILY: Apocynaceae

COMMON NAME: poison tree, bushman's-poison, poison bush

NATIVE: E. Africa

LISTING ONLY:

DeWoskin 1981; Mitchell et al. 1977; Weems 1981.

Actinidia chinensis Planch.

FAMILY: Actinidiaceae

COMMON NAME: Chinese gooseberry, kiwi berry, yang-tao

NATIVE: China, Taiwan

CULTIVATED: New Zealand

LABORATORY INFESTATION:

Ozaki 1980: in initial tests in small cages, no infestation took place on normal, unbroken fruits; in a larger cage, 10 normal fruits were exposed to flies for 24 h; they eventually developed 0.5 larva per fruit; after 48 h of ex-

posure, five fruits developed 4.6 larvae per fruit; other fruits were also pierced or shaved (or both) to promote oviposition; the number of larvae per fruit ranged from 0.2 to 5.

LISTING ONLY:

*DeWoskin 1981; Lotter 1982.

Adelia segregata Kuntze

FAMILY: Oleaceae

COMMON NAME: Adelia, Florida privet

LABORATORY INFESTATION:

Keck & Marshall 1930: fruits were placed in cages and exposed to flies; infestation was determined by presence of larvae recorded on 26-XII-1929.

LISTING ONLY:

Chilson 1960; DeWoskin 1981; Mitchell et al. 1977; Weems 1981.

Anacardium occidentale L.

FAMILY: Anacardiaceae

COMMON NAME: cashew, maranon

NATIVE: Tropical America

CULTIVATED: India

LISTING ONLY:

Eskafi & Cunningham 1987.

Ananas comosus (L.) Merrill

FAMILY: Bromeliaceae

COMMON NAME: pineapple

NATIVE: Tropical America

CULTIVATED: Hawaii, West Indies

LISTING ONLY:

*Christenson & Stone 1956; DeWoskin 1981;

*French 1907; Mclean 1929; Mitchell et al. 1977;

Orian & Moutia 1960; *Phillips 1946; *Quaint-

ance 1912; Weems 1981.

Ananas sativus Schult., see *A. comosus* (L.) Merrill

Annona sp.

FAMILY: Annonaceae

LISTING ONLY:

Rivnay 1950.

Annona cherimola Mill.

FAMILY: Annonaceae

COMMON NAME: cherimoya, chirimoya, custard apple

NATIVE: Andes of Peru and Ecuador

CULTIVATED: Tropical highlands, subtropics