

References Cited

- Athias-Binche, F.** 1991. Ecology and evolution of phoresy in mites, pp. 27–41. In F. Dusbábek and V. Bukva [eds.], Modern acarology. Academia, Prague.
- Baker, E. W., and C. E. Yunker** 1964. New blattisociid mites (Acarina: Mesostigmata) recovered from neotropical flowers and hummingbirds' nares. Ann. Entomol. Soc. Am. 57(1): 103–126.
- Berry, F., and J. W. Kress.** 1991. *Heliconia*: an identification guide. Smithsonian Institution, Washington, DC.
- Christiansen, K. M., R. K. Colwell, and M. J. Kaliszewski** 1992. Celulose acetate electrophoretic techniques for the genetic analysis of individual ascid mites (Mesostigmata: Ascidae). Int. J. Acarol. 18(2): 97–105.
- Colwell, R. K.** 1973. Competition and coexistence in a simple tropical community. Am. Nat. 107(958): 737–760.
- 1979a.** Toward a unified approach to the study of species diversity, pp. 75–91. In J. F. Grassle, G. P. Patil, W. K. Smith, and C. Taillie [eds.], Ecological diversity in theory and practice. International Cooperative Publishing, Fairland, MD.
- 1979b.** The geographical ecology of hummingbird flower mites in relation to their host plants and carriers, pp. 461–468. In J. G. Rodriguez [ed.], Recent advances in acarology, vol. 2. Academic, New York.
- 1983.** *Rhinoseiulus colwelli* (Acaro Floral del Colibri, Totolate Floral de Colibri, Hummingbird Flower Mite), pp. 767–768. In D. H. Janzen [ed.], Costa Rican natural history. University of Chicago Press, Chicago.