



## **Range Expansion of the Litchi Erinose Mite *Aceria litchii* (Acari: Eriophyidae) in Brazil**

Authors: Fornazier, Maurício José, Martins, David Dos Santos, Fornazier, Débora Lorenção, Azevedo, Letícia Henrique, Jr, José Salazar Zanuncio, et al.

Source: Florida Entomologist, 97(2) : 846-848

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.097.0276>

---

BioOne Complete ([complete.BioOne.org](https://complete.bioone.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## RANGE EXPANSION OF THE LITCHI ERINOSE MITE *ACERIA LITCHII* (ACARI: ERIOPHYIDAE) IN BRAZIL

MAURÍCIO JOSÉ FORNAZIER<sup>1,4</sup>, DAVID DOS SANTOS MARTINS<sup>1</sup>, DÉBORA LORENÇÃO FORNAZIER<sup>2</sup>, LETÍCIA HENRIQUE AZEVEDO<sup>3</sup>, JOSÉ SALAZAR ZANUNCIO JR<sup>1</sup> AND JOSÉ COLA ZANUNCIO<sup>1,4\*</sup>

<sup>1</sup>Instituto Capixaba de Pesquisa, Assistência Técnica e Extensão Rural - Incaper, Rua Afonso Sarlo, 160, Bento Ferreira, 29052-010, Vitória, ES, Brazil

<sup>2</sup>Centro de Ciências Agrárias, Universidade Federal do Espírito Santo, Alto Universitário, Cx. Postal 16, 29500-000, Alegre, ES, Brazil

<sup>3</sup>Escola Superior de Agricultura Luiz de Queiróz. Avenida Pádua Dias, 11, Cx. Postal 9, 13418-900, Piracicaba, SP, Brazil

<sup>4</sup>Departamento de Entomologia, Universidade Federal de Viçosa. 36570-000, Viçosa, MG, Brazil

\*Corresponding author; E-mail: zanuncio@ufv.br

Litchi (*Litchi chinensis* Sonn.) being indigenous to the moist forests of China, belongs to the Sapindaceae family, which has approximately 150 genera and 2,000 species in the tropics and subtropics (Paull & Duarte 2011). *Paullinia cupana* Kunth is the best known species of this family in Brazil. *Litchi chinensis* needs winter cooling to stimulate its flowering, and it is cultivated in most parts of the world including Brazil and other Latin American countries (Diczbalis 2011; Paull & Duarte 2011). Commercial litchi orchards were initiated in Brazil in the 1970-80s. The major cultivar ('Bengal') produces fruits from Nov to Jan (Pimentel & Celim 2013) and its fresh fruits are sold in domestic markets at Christmas.

Fifty-eight pest species have been recorded on litchi trees worldwide, including lepidopterans, scales, stink bugs, fruit flies, and eriophiid mites, and these pest mainly damage the flowers and fruits (Waite 2012). The litchi erinose mite (LEM), *Aceria litchii* (Keifer) (Acari: Eriophyidae) is a major pest of litchi varieties (Menzel 2002; Paull & Duarte 2011). LEM is not visible to the naked eye. LEM is specific to litchi, affecting new shoots on the entire tree during severe infestations (Paull & Duarte 2011), and causing abnormal development and premature defoliation (Waite 2012). Litchi plants with severe erinose symptoms in young and developing leaves were detected in Brazil in Jan 2008 in an orchard of 3,000 thirteen-yr-old litchi plants of the cultivar 'Bengal' in the municipality of Limeira, São Paulo State. A dense population of this mite was detected during the litchi fruiting period, and this represents the first report of *A. litchii* in South America (Picoli et al. 2010; Raga et al. 2010).

The detection of LEM in Espírito Santo State, Brazil (S 20° 20' 26" -W 41° 06' 59", 750 m asl) in Dec 2012 represents its first range expansion in this country. The leaves (Fig. 1A), and mature litchi fruits showed mild erinose mite symptoms, but another orchard nearby had no symptoms. In-

festated litchi leaves with erinose mite symptoms (Fig. 1B) were collected, placed in Kraft paper bags and transported to the regional laboratory of entomology of INCAPER (Instituto Capixaba de Pesquisa, Assistência Técnica e Extensão Rural) from Dec 2012 to Feb 2013 to identify the mite species. These leaves were left for 2 h under direct sunlight to force the mites to leave the erineum (Waite 1992), the latter being an abnormal feltly growth of hairs from the leaf epidermis. Live mites were observed under the microscope, and infested leaves were cut into 4 cm<sup>2</sup> pieces, preserved in 70% alcohol, and sent to Dr. Gilberto José de Moraes of the Laboratory of Entomology and Acarology, Escola Superior de Agricultura Luiz de Queiróz, University of São Paulo (ESALQ-USP), who identified them as *Aceria litchii* (Keifer) (Acari: Eriophyidae) (Fig. 1C). Leaves from trees of the second orchard produced by air-layering were collected, but they showed no LEM or erinose mite symptoms.

Insecticide sprays to control litchi pests, such as *Tessaratomia papillosa* Drury (Hemiptera: Tessaratomidae) (Zeng et al. 2001) can increase LEM populations (PIP 2011). However, no insecticides or fungicides were sprayed in the LEM-infested area. Pruning and burning of affected twigs were the adopted control measures, particularly for severely infested litchi trees. Pruning and burning reduced LEM symptoms up to 3 months when new shoots presented symptoms. Three sprays of wettable sulphur, permitted for use on organic crops, at the 0.4% rate at 15-day-intervals were used. No pesticide is registered to control litchi pests in Brazil, although chemical products are used to control this mite in other countries (Schulte et al. 2007; Brasil 2013).

Predators are important for LEM management, particularly those of the Phytoseiidae family, but Waite (2012) reported them to have low efficiency on *A. litchii*. *Amblyseius* spp., *Euseius* spp., *Iphiseiodes* spp., *Phytoseius* spp., *Okiseius*




Fig. 1. A. Eighteen year old litchi tree damaged by *Aceria litchii* (Acari: Eriophyidae); B. Variations in leaf maturation and damage; and C., *Aceria litchii*. The infested orchard was located in the municipality of Venda Nova do Imigrante, Espírito Santo State, Brazil.

spp., *Typhlodromus* spp. (Acari: Phytoseiidae), and *Agistemus* spp. (Acari: Stigmaeidae) are associated with LEM worldwide, including Brazil (Picoli et al. 2010). The number of species and endemism of predatory Phytoseiidae mites are high in the Neotropical region (Tixier et al. 2008). Detection of predatory mites is important for the biological control of *A. litchii*.

LEM and its damage should be sampled in the litchi orchards in Espírito Santo State, Brazil. The expansion of the LEM on litchi into a second Brazilian state and measures for its management are presented.

#### ACKNOWLEDGMENTS

To the “Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)”, “Coordenação do Aperfeiçoamento de Pessoal de Nível Superior (CAPES)”, “Fundação de Amparo à Pesquisa do Espírito Santo (FAPES)”, and “Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG)” for financial support. To Dr. Gilberto José de Moraes for the mite identification. Global Edico Services edited and proofread this manuscript.

#### SUMMARY

Litchi (*Litchi chinensis* Sonn.) is cultivated in subtropical and tropical regions, and in Brazil, commercial orchards began in the 1970-80s. The litchi erinose mite (LEM), *Aceria litchii* (Keifer) (Acari: Eriophyidae), is a pest of litchi worldwide. In Brazil, LEM was first reported in this crop in São Paulo State in 2008, and Espírito Santo is the second Brazilian state with infestation of this mite discovered in a 18-year-old orchard of 150 plants of the litchi cultivar ‘Bengal’. Cultural and chemical measures for LEM management are presented, and this is the first range expansion of LEM into Espírito Santo State, Brazil.

Key Words: Bengal, Eriophyidae, *Litchi chinensis*, lychee, management, range expansion

#### RESUMO

Lichia (*Litchi chinensis* Sonn.) é cultivada em regiões tropicais e subtropicais e no Brasil pomares comerciais desta planta começaram na década de 1970-80. O ácaro da erinose da lichia (LEM), *Aceria litchii* (Keifer) (Acari: Eriophyidae) é praga importante desta planta no mundo. No Brasil, LEM foi relatado pela primeira vez no Estado de São Paulo em 2008 e o Espírito Santo é o segundo estado brasileiro a confirmar a infestação com este ácaro em pomar de 18 anos, com 150 plantas de lichia, cultivar Bengal. Medidas para o manejo do LEM são apresentadas e este é o primeiro relato da expansão do LEM no Estado do Espírito Santo, Brasil.

Palavras-Chave: Bengal, Eriophyidae, expansão da área, lichia, *Litchi chinensis*, manejo

#### REFERENCES CITED

- BRASIL. 2013. Sistema de agrotóxicos fitossanitários. Available at [http://extranet.agricultura.gov.br/agrofit\\_cons/principal\\_agrofit\\_cons](http://extranet.agricultura.gov.br/agrofit_cons/principal_agrofit_cons). Accessed on 23 May 2013.
- DICZBALIS, Y. 2011. Farm and forestry production and marketing profile for Lychee. 13 pp. Available at <http://www.agroforestry.net/scps>. Accessed on 23 May 2013.
- MENZEL, C. 2002. Major pests and diseases, pp. 74-83 In C. Menzel [ed.], The lychee crop in Asia and the Pacific. FAO. Available at <http://www.fao.org/docrep/005/ac681e/ac681e09.htm>. Accessed on 23 May 2013.

- PAULL, R. E., AND DUARTE, O. 2011. Litchi and longan, pp. 221-251 *In* R. E. Paull and O. Duarte [ed.], Tropical fruits, 2nd ed. CABI. 400 pp.
- PICOLI, P. R. F., VIEIRA, M. R., SILVA, E. A., AND MOTA, M. S. O. 2010. Ácaros predadores associados ao ácaro-da-erinoze da lichia. *Pesqui. Agropecu. Brasileira* 45: 1246-1252.
- PIMENTEL, B. C., AND CELIM, A. 2013. A lichia na Ceagesp de 1999 a 2009. Available at [http://www.hortibrasil.org.br/jnw/index.php?option=com\\_content&view=article&id=942:a-lichia-na-ceagesp-de-1999-a-2009&catid=64:frutas-e-hortalicas-frescas&Itemid=82](http://www.hortibrasil.org.br/jnw/index.php?option=com_content&view=article&id=942:a-lichia-na-ceagesp-de-1999-a-2009&catid=64:frutas-e-hortalicas-frescas&Itemid=82). Accessed on 23 May 2013.
- PIP. 2011. Guide to good crop protection practices for the lychee (*Litchi chinensis*) in ACP countries. Available at: [http://pip.coleacp.org/files/documents/GB-PP-Litchee%2004-2011-11-1-UK\\_0.pdf](http://pip.coleacp.org/files/documents/GB-PP-Litchee%2004-2011-11-1-UK_0.pdf). Accessed on 15 Jul 2013.
- RAGA, A., MINEIRO, J. L. C., SATO, M. E., MORAES, G. J., AND FLECHTMANN, C. H. W. 2010. Primeiro relato de *Aceria litchii* (Keifer) (Prostigmata: Eriophyidae) em plantas de lichia no Brasil. *Rev. Brasileira Frutic.* 32: 628-629.
- SCHULTE, M. J., MARTIN, K., AND SAUERBORN, J. 2007. Efficacy of spiromesifen on *Aceria litchii* (Keifer) in relation to *Cephaluros virescens* Kunze colonization on leaves of litchi (*Litchi chinensis* Sonn.). *J. Plant Dis. Prot.* 114: 133-137.
- TIXIER, M. S., KREITER, S., AND MORAES, G. J. 2008. Biogeographic distribution of the Phytoseiidae (Acari: Mesostigmata). *Biol. J. Linnean Soc.* 93: 845-856.
- WAITE, G. K. 1992. Pest Management in Lychees. Australian Rural Ind. Res. Dev. Corp., Canberra, Australia. 42 pp.
- WAITE, G. 2012. Lychee erinose mite in lychees. Queensland: Dept. Plant Industries and Fisheries. Available at: <http://www.daff.qld.gov.au/plants/fruit-and-vegetables/a-z-list-of-horticultural-insect-pests/lychee-erinose-mite>. Accessed on 19 Aug 2013.
- ZENG, X. N., DENG, D., AND WANG, J. M. 2001. Chlorpyrifos and cypermethrin for the control of litchi stink bug (*Tessarotoma papillosa*). *Acta Hort.* 558: 421-423.