

## **The evolution of sexual size dimorphism: the interplay between natural and sexual selection**

Authors: Castillo, Raúl Cuevadel, and Núñez-Farfán, Juan

Source: Journal of Orthoptera Research, 17(2) : 197-200

Published By: Orthopterists' Society

URL: <https://doi.org/10.1665/1082-6467-17.2.197>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# The evolution of sexual size dimorphism: the interplay between natural and sexual selection

Accepted November 7, 2008

RAÚL CUEVA DEL CASTILLO AND JUAN NÚÑEZ-FARFÁN

(RCC) Laboratorio de Ecología, UBIPRO; Facultad de Estudios Superiores Iztacala, Universidad Nacional Autónoma de México. Apartado Postal 314, Mexico 54090, Estado de México, México. E-mail: rcueva@ecologia.unam.mx

(JN-F) Laboratorio de Genética Ecológica y Evolución, Departamento de Ecología Evolutiva, Instituto de Ecología, Universidad Nacional Autónoma de México. Apartado Postal 70-275, Mexico 04510, Distrito Federal. E-mail: farfan@servidor.unam.mx

## Abstract

Sexual size dimorphism (SSD) in animal species can result from the interplay between natural and sexual selection. In this paper we review the impact of sexual and natural selection on grasshopper body size and the evolution of SSD. Mate choice by females, and natural selection on female fecundity could explain an evolutionary trend to increase SSD in species in which females receive nutritional benefits during mating. In general, sexual selection is stronger in males than females. However, when females receive nutritional resources from males during mating, selection could be stronger in females than males. These resources constitute high energetic costs to males and it is expected that this promotes an increment in male mate selectivity. Higher female-biased SSD might evolve as a result of polyandry in species where males transfer nutritional benefits in the ejaculate. This hypothesis is testable at both macro- and micro-evolutionary levels. Finally, we discuss the relationship between body size and mate-guarding duration and its evolutionary implications and propose future studies to analyze the evolution of SSD and mate-guarding duration in grasshoppers.

## Key words

Orthoptera, katydid, grasshopper, sexual size dimorphism, evolution, body size, sexual selection, mate-guarding, nuptial gift

## Evolution of sexual size dimorphism: effects of natural and sexual selection

The direction and magnitude of selection on body size may differ between sexes and generate sexual size dimorphism (SSD). SSD may result from an interplay between sexual and natural selection (Slatkin 1984, Hedrick & Temeles 1989, Shine 1989, Fairbairn 1997). In insects, body size in both males and females is a target of directional selection. Large females generally have higher fecundity (because of larger clutches), and thus natural selection may favor large female body size (Ridley 1983, Fairbairn 1997). On the other hand, large males often have advantages in male-male competition and female choice (Thornhill & Alcock 1983). If females are larger than males, this suggests that natural selection for high female fecundity could be stronger than sexual selection on males (Ridley 1983, Wiklund & Karlsson 1988). Such differences in selection between males and females can produce SSD. However, the magnitude of SSD is expected to be smaller in those insect taxa where sexual selection favors large males (see Fairbairn & Preziosi 1994). Nevertheless, in insect species where females are larger than males, female size in many cases increases proportionally more as male body size increases (*i.e.*, in different populations or different rearing conditions), thus augmenting SSD (see Teder & Tammaru 2005 for a review).

Why does SSD increase as body size increases in some clades, but decrease in others? Some hypotheses can be offered to explain such observations (Fairbairn 1997, Blanckenhorn & Demont 2004, Fairbairn 2005, Teder & Tammaru 2005, Cueva del Castillo & Gwynne 2007), and these are based on the relative magnitude of sexual selection on females and males. Male fitness is generally related to the number of copulations, whereas female fitness is limited by the amount of resources that can be devoted to produce eggs (Trivers 1972, Andersson 1994). However, male resource investment can alter the intensity of sexual selection pressures on the sexes (Trivers 1972). Thus, food stress can increase sexual selection on females for increasing size; here, selection might occur through female to female competition for access to nurturant males and through male mating preferences, when costly nutrient contributions limit male mating frequency (Rutowski 1982, Gwynne 2004).

Because nutritional resources represent energetic costs to males, it is expected that males will show mate choice, rejecting low-quality females and competing for access to high-quality females (Bonduriansky 2001). The extreme situation can be found in species where there is sex-role reversal. In the katydids *Anabrus simplex* and *Kawanaphila nartee*, a reduction in food availability causes both a decrease in the number of males that are able to produce high quality spermatophylax meals (Gwynne & Simmons 1990; Gwynne 1993, 2001), and an increase in hungry females looking for these mating meals. Females fight for access to males and males prefer large females, rejecting smaller ones (Gwynne 1984, 1993; Simmons & Bailey 1993). Female quality may be "assessed" through body size which is correlated with fecundity. Large females are more fecund (Ridley 1983, Honěk 1993) and may have a higher mating success than smaller ones (Ridley 1983, Bonduriansky 2001, Cueva del Castillo & Núñez-Farfán 2002).

In grasshoppers and other orthopterans, males can transfer nutritional fluids that may increase egg-laying (see Friedel & Gillot 1977, Butlin *et al.* 1987, Muse 1992, Pardo *et al.* 1995, Tregenza & Wedell 1998, Reinhardt *et al.* 1999, Wagner *et al.* 2001), and females can increase their reproductive success by mating repeatedly (see Arnqvist & Nilsson 2000). The potential interaction between natural selection on female fecundity and sexual selection due to male mate choice may explain the results found by Teder and Tammaru (2005) (but see Blanckenhorn & Demont 2004). The six orthopteran species included in Teder and Tammaru's (2005) review show higher intraspecific female-biased SSD, as body size increases across different populations or environments (rearing conditions). Higher female-biased SSD might evolve as a result of polyandry in species where males transfer nutritional benefits in the ejaculate (see Cueva del Castillo & Gwynne 2007).

|                       |      | Male nutrient transference |   |          |   |
|-----------------------|------|----------------------------|---|----------|---|
|                       |      | Absence                    |   | Presence |   |
| Resource availability | Low  | <b>A</b>                   | SS on ♀ +<br>SS on ♂ +++<br>NSF on ♀ +++<br>S on ♂PF 0<br><br>(Moderate SSD - biased to ♀♀) | <b>B</b> | SS on ♀ +++<br>SS on ♂ +<br>NSF on ♀ +++<br>S on ♂PF +++<br><br>(High SSD biased to ♀♀) |
|                       | High | <b>C</b>                   | SS on ♀ +<br>SS on ♂ +++<br>NSF on ♀ +<br>S on ♂PF 0<br><br>(No SSD or SSD biased to ♂♂)    | <b>D</b> | SS on ♀ ++<br>SS on ♂ ++<br>NSF on ♀ +++<br>S on ♂PF ++<br><br>(Low SSD biased to ♀♀)   |

Selection:   +++ high  
                  ++ moderate  
                  + low  
                  0 null

**Fig. 1.** Four ♂♀ potential scenarios for the evolution of sexual size dimorphism (SSD). Taken into consideration are: NSF= natural selection on fecundity; SS = sexual selection on ♀♀ and ♂♂; S on ♂PF = selection on males promoting (female) fecundity. Selection occurs in a variable environment regarding food-resource availability and whether or not males transfer nutritional resources to females during copulation. Predicted SSD are in parentheses.

This hypothesis can be tested at both macro- and microevolutionary scales. For this, it is necessary to compare the evolution of SSD in species where polyandry increases female fecundity *vs* those species where polyandry does not increase female fecundity. It is also possible to compare the magnitude of natural and sexual selection on body size in males and females. The differences in the intensity of sexual selection on females' and males' body size can vary dramatically between them. For instance, Jann *et al.* (2000) found that in the dung fly *Scathophaga stercoraria*, selection on male body size was more than twice the magnitude of selection on female body size; however, we do not know if females receive direct benefits due to mating. In the grasshopper *Leptysmia argentina*, sexual selection was stronger on male femur length than on female femur length, but selection on thorax length was stronger in females than in males (Colombo *et al.* 2004). In the grasshopper *Eyprepocnemis plorans*, somatic condition (body size and somatic mass) and reproductive condition (gonad mass) were positively correlated to female, but not to male, mating success (Martín-Alganza *et al.* 1997). In *S. purpurascens* the magnitude of selection on body size was similar in both sexes (Cueva del Castillo & Núñez-Farfán 1999, 2002). In this latter species, mating duration increases female fecundity, but apparently polyandry does not (Lugo-Olguín & Cueva del Castillo 2007).

**Body size, mate guarding and sperm competition**

Sperm competition is thought to be common in insects and relevant in determining male reproductive success. In katydids, larger males allow more time for ejaculate transfer and thus achieve more fertilizations (Leimar *et al.* 1994, Vahed 1998). However, there are few studies in grasshoppers in relation to sperm competition (see Simmons 2001). Grasshoppers tend to be polygamous, and pre- and postcopulatory mate-guarding behavior has been documented in some species (Parker & Smith 1975, Wickler & Seibt 1985, Muse

& Ono 1996, Cueva del Castillo *et al.* 1999, Zhu & Tanaka 2002). Anecdotic reports of extraordinary female-male guarding periods suggest that long-duration guarding could be a phylogenetically inherited trait in the Pyrgomorphidae (see Descamps & Winterbert 1966). In four species, *Zonocerus elegans*, *Atractomorpha lata*, *Sphenarium purpurascens*, and *Sphenarium magnum* — males can spend long periods mounted on females (*Z. elegans*: up to 45 d, Wickler & Seibt 1985; *A. lata*: up to 16 h, Muse & Ono 1996; *S. purpurascens*: up to 18 d, Cueva del Castillo *et al.* 1999; and *S. magnum*: up to 21 d Cueva del Castillo, pers. obs.). In *S. purpurascens*, the duration of guarding and the number of copulations are positively related to female body size (Cueva del Castillo 2003). Interestingly, *A. lata* is the smallest of the four species and has the shortest guarding duration, whereas *Z. elegans* is the largest species and has the longest guarding duration. Moreover, it seems that males might change their guarding duration according to the potential risk of sperm competition and paternity payoff. For instance, males of *S. purpurascens* invest more time guarding and copulating with large females or females that have been previously mated by other males (see Cueva del Castillo 2003, Lugo-Olguín & Cueva del Castillo 2007). However, before accepting this interpretation, it is necessary to know to what extent guarding behavior is related to the likelihood of paternity in pyrgomorphids.

Long-duration guarding could be adaptive from the female's perspective if during the association, as result of multiple copulations, there is a transference of nutritional resources that increase females' longevity and/or fecundity (see Butlin *et al.* 1987; Muse 1992, 2002). Perhaps larger males contribute more to female fecundity, transferring larger nutritional donations than smaller ones. Nevertheless, this hypothesis remains to be tested as well.

Guarding duration can be affected by several factors, including: i) energetic and predation costs for each sex, ii) operational sex ratio, and iii) body size (see Alcock 1994). The prolonged guarding periods in some members of the Pyrgomorphidae open an opportunity to

study in detail, the energetic and predatory costs of guarding. For instance in *S. purpurascens*, males can feed only in an opportunistic way during guarding (e.g., if they get positioned close to plant leaves and no potential rival males are nearby). Furthermore, because optimal guarding duration can differ between females and males, it may give rise to a conflict of interest between sexes.

### Future studies

Several topics of the evolution of mating systems deserve further study in grasshoppers. A complete understanding of the evolution of body size and SSD requires the simultaneous analysis of the impact of sexual and natural selection on female and male body size (Fairbairn *et al.* 2007). Particularly interesting is the study of the consequences of 1) the availability of environmental resources, and 2) the presence/absence of the transference of nutritional resources during copulation, on the evolution of body size and SSD. Considering a discrete variation in resource availability (low and high), four potential scenarios can be analyzed to predict the evolution of SSD (Fig. 1). In the A scenario, the reproductive cost is higher in females than males and selection to increase fecundity is expected to be greater than sexual selection on males; here, moderate SSD biased toward females is expected. The B scenario is very interesting because the low availability of resources promotes sex-role reversal, with female-female competition, selection on female fecundity and male mate choice; accordingly, high SSD biased to females is expected. In the C scenario, sexual selection on males is higher than sexual selection on females and selection on female fecundity; very low or even no SSD biased toward females is expected but SSD biased toward males is possible. Finally, in D scenario reciprocal sexual selection plus selection on females' fecundity predicts moderate SSD biased toward females (Fig. 1).

Finally, much of the information regarding mating system evolution and SSD in grasshoppers has derived from Holarctic species. In contrast, the diverse contingent of tropical grasshopper species is still poorly known. Due to their diversity and abundance, tropical grasshopper species remain as potential and valuable models to test these evolutionary hypotheses.

### Acknowledgements

The authors are indebted to Douglas Whitman, Glenn Morris and two anonymous reviewers for their helpful discussion and critical comments on this manuscript.

### References

- Alcock J. 1994. Post-insemination association between males and females in insects: the mate-guarding hypothesis. *Annual Review of Entomology* 39: 1-21.
- Andersson M. 1994. *Sexual Selection*. Princeton University Press, Princeton, NJ.
- Arnqvist G.N., Nilsson T. 2000. The evolution of polyandry: multiple mating and female fitness in insects. *Animal Behavior* 60: 145-164.
- Blanckenhorn W.U., Demont M. 2004. Bergmann and converse Bergmann latitudinal clines in arthropods: two ends of a continuum? *Integrative and Comparative Biology* 44: 413-424.
- Bonduriansky R. 2001. The evolution of male mate choice in insects: a synthesis of ideas and evidence. *Biological Reviews* 76: 305-339.
- Butlin R.K., Woodhatch C.W., Hewitt G.M. 1987. Male spermatophore investment increases female fecundity in a grasshopper. *Evolution* 41: 221-225.
- Colombo P.C.S., Pensel S., Isabel R.M. 2004. Chromosomal polymorphism, morphometric traits and mating success in *Leptysma argentina* Bruner (Orthoptera). *Genetica* 121: 25-31.
- Cueva del Castillo R. 2003. Body size and multiple copulations in a neotropical grasshopper with an extraordinary mate-guarding duration. *Journal of Insect Behavior* 16: 503-522.
- Cueva del Castillo R., Gwynne D.T. 2007. Increase in song frequency decreases spermatophore size: correlative evidence of a macroevolutionary trade-off in katydids (Orthoptera: Tettigoniidae). *Journal of Evolutionary Biology* 20: 1028-1036.
- Cueva del Castillo R., Núñez-Farfán J. 1999. Sexual selection on maturation time and body size in *Sphenarium purpurascens* (Orthoptera: Pyrgomorphidae): correlated response to selection. *Evolution* 53: 209-215.
- Cueva del Castillo R., Núñez-Farfán J. 2002. Female mating success and risk of prereproductive death in a protandrous grasshopper. *Oikos* 96: 217-224.
- Cueva del Castillo R., Núñez-Farfán J., Cano-Santana Z. 1999. The role of body size in mating success of *Sphenarium purpurascens* in central Mexico. *Ecological Entomology* 24: 146-155.
- Descamps M., Winterbert D. 1966. Pyrgomorphidae et Acrididae de Madagascar. Observations biologiques et diagnoses (Orth. Acridoidea). *Revista Española de Entomología* 42: 41-263.
- Fairbairn D.J. 1997. Allometry for sexual size dimorphism: patterns and process in the coevolution of body size in females and males. *Annual Review of Ecology and Systematics* 28: 659-687.
- Fairbairn D.J. 2005. Allometry for sexual size dimorphism: testing two hypotheses for Rensch's rule in the water strider *Aquarius remigis*. *American Naturalist* 166: S69-S84.
- Fairbairn D.J., Preziosi R.F. 1994. Sexual selection and the evolution of allometry for sexual size dimorphism in the water strider, *Aquarius remigis*. *American Naturalist* 144: 101-118.
- Fairbairn D.J., Blanckenhorn W.U., Székely T. 2007. *Sex, Size & Gender Roles*. Evolutionary Studies of Sexual Size Dimorphism. Oxford University Press, Oxford.
- Friedel T., Gillot C. 1977. Contribution of male-produced protein to vitellogenesis in *Melanoplus sanguinipens*. *Journal of Insect Physiology* 23: 145-151.
- Gwynne D.T. 1984. Sexual selection and sexual differences in mormon crickets (Orthoptera: Tettigoniidae, *Anabrus simplex*). *Evolution* 38: 1011-1022.
- Gwynne D.T. 1993. Food quality controls sexual selection in mormon crickets by altering male mating investment. *Ecology* 74: 1406-1413.
- Gwynne D.T. 2001. *Katydid and Bush-crickets: Reproductive Behavior and Evolution of the Tettigoniidae*. Comstock Pub. Associates, Ithaca, NY.
- Gwynne D.T. 2004. Sexual differences in response to larval food stress in two nuptial feeding orthopterans: implications for sexual selection. *Oikos* 105: 619-625.
- Gwynne D.T., Simmons L.W. 1990. Experimental reversal of courtship roles in an insect. *Nature* 346: 172-174.
- Hedrick A.V., Temeles E.J. 1989. The evolution of sexual dimorphism in animals: hypotheses and tests. *Trends in Ecology and Evolution* 4: 136-138.
- Honek A. 1993. Intraspecific variation in body size and fecundity in insects: a general relationship. *Oikos* 66: 483-492.
- Jann P., Blanckenhorn W.U., Ward P.I. 2000. Temporal and microspatial variation in the intensities of natural and sexual selection in the yellow dung fly *Scathophaga stercoraria*. *Journal of Evolutionary Biology* 13: 927-938.
- Leimar O., Karlsson B., Wiklund C. 1994. Unpredictable food and sexual size dimorphism in insects. *Proceedings Royal Society of London, Series B*. 258: 121-125.
- Lugo-Olguín S.D., Cueva del Castillo R. 2007. Multiple copulation, female fecundity and evaluation of sperm competition risk in a protandrous grasshopper. *Annals Entomological Society of America* 100: 591-595.
- Martín-Alganza A., López León M.D., Cabrero J., Camacho J.P.M. 1997. Somatic condition determines female mating frequency in a field population of the grasshopper *Epyreprocnemis plorans*. *Heredity* 79: 524-530.

- Muse W.A. 1992. Fecundity-enhancing substance in the accessory reproductive gland of adult grasshopper, *Zonocerus variegatus* L. (Orthoptera: Acridoidea, Pyrgomorphidae). *Journal of Agricultural Science and Technology* 2: 92-94.
- Muse W.A., Ono T. 1996. Copulatory behavior and postcopulatory mate guarding in a grasshopper *Atractomorpha lata* Motschulsky (Orthoptera: Tetrigidae) under laboratory conditions. *Applied Entomology and Zoology* 31: 233-241.
- Muse W.A. 2002. Morphology of the male reproductive system and the nature of secretions of the accessory glands and seminal vesicles of adult *Atractomorpha lata* Motschulsky (Orthoptera: Acrididae). *Formosan Entomology* 22: 249-294.
- Pardo M.C., López-León M.D., Hewitt G.M., Camacho J.P.M. 1995. Female fitness is increased by frequent mating in grasshoppers. *Heredity* 73: 654-660.
- Parker G.A., Smith L.A. 1975. Sperm competition and the evolution of the precopulatory passive phase behaviour in *Locusta migratoria migratorioides*. *Journal of Entomology* 49: 155-171.
- Reinhardt K., Köhler G., Schumacher J. 1999. Females of the grasshopper *Chorthippus parallelus* (Zett.) do not remate for fresh sperm. *Proceedings Royal Society of London, Series B*. 266: 2003-2009.
- Ridley M. 1983. *The Explanation of Organic Diversity: the Comparative Method and Adaptation for Mating*. Clarendon Press, Oxford.
- Rutowski R.L. 1982. Epigamic selection by males as evidenced by courtship partner preferences in the checkered white butterfly (*Pieris protodice*). *Animal Behavior* 30: 108-112.
- Shine R. 1989. Ecological causes for the evolution of sexual dimorphism: a review of the evidence. *Quarterly Review of Biology* 64: 419-461.
- Simmons L.W. 2001. *Sperm Competition and its Evolutionary Consequences in the Insects*. Princeton University Press, Princeton, NJ.
- Simmons L.W., Bailey W.J. 1993. Agonistic communication between males of a zaprochiline katydid (Orthoptera: Tettigoniidae). *Behavioral Ecology* 4: 364-368.
- Slatkin M. 1984. Ecological causes of sexual dimorphism. *Evolution* 38: 622-630.
- Teder T., Tammaru T. 2005. Sexual size dimorphism within species increases with body size in insects. *Oikos* 108: 321-334.
- Thornhill R., Alcock J. 1983. *The Evolution of Insect Mating Systems*. Harvard University Press, Cambridge, MA.
- Tregenza T., Wedell N. 1998. Benefits of multiple mates in the cricket *Gryllus bimaculatus*. *Evolution* 52: 1726-1730.
- Trivers R.L. 1972. Parental investment and sexual selection, pp. 136-179. In: Cambell B. (Ed.) *Sexual Selection and Descent of Man*. Aldine, Chicago.
- Vahed K. 1998. The function of nuptial feeding in insects: review of empirical studies. *Biological Reviews* 73: 43-78.
- Wagner W.E. Jr., Kayleen R.J., Tucker K.R., Harper C.J. 2001. Females receive a life-span benefit from male ejaculate in a field cricket. *Evolution* 55: 994-1001.
- Wickler W., Seibt U. 1985. Reproductive behavior in *Zonocerus elegans* (Orthoptera: Pyrgomorphidae) with special reference to nuptial gift guarding. *Zeitschrift für Tierpsychologie* 69: 203-223.
- Wiklund C., Karlsson B. 1988. Sexual size dimorphism in relation to fecundity in some Swedish satyrid butterflies. *American Naturalist* 131: 132-138.
- Zhu D-H., Tanaka S. 2002. Prolonged precopulatory mounting increases the length of copulation and sperm precedence in *Locusta migratoria* (Orthoptera: Acrididae). *Annals Entomological Society of America* 95: 370-373.