

## **Functional and Numerical Responses of the Predatory Mite, *Neoseiulus longispinosus*, to the Red Spider Mite, *Oligonychus Coffeae*, Infesting Tea**

Authors: Rahman, Vattakandy jasin, Babu, Azariah, Roobakkumar, Amsalingam, and Perumalsamy, Kandasamy

Source: Journal of Insect Science, 12(125) : 1-12

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.012.12501>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Functional and Numerical Responses of the predatory mite, *Neoseiulus longispinosus*, to the red spider mite, *Oligonychus coffeae*, infesting tea

Vattakandy Jasin Rahman<sup>a\*</sup>, Azariah Babu<sup>#b</sup>, Amsalingam Roobakkumar<sup>c</sup>, and Kandasamy Perumalsamy<sup>d</sup>

Entomology Division, UPASI Tea Research Foundation Tea Research Institute, Nirar Dam (BPO), Valparai 642 127 Coimbatore (Dt), Tamil Nadu, India

<sup>#</sup>Current address: Department of Entomology, Tocklai Experimental Station, Tea Research Association, Jorhat-785 008, Assam, India

### Abstract

Functional and numerical responses of the predatory mite, *Neoseiulus longispinosus* (Evans) (Acari: Phytoseiidae) to the red spider mite, *Oligonychus coffeae* Nietner (Acari: Tetranychidae), infesting tea were determined in a laboratory on leaf discs. Prey consumption increased with increases in temperature and prey density. Handling time decreased and successful attack rate increased with increased temperature. *N. longispinosus* was more voracious on larvae and nymphs than on adults of *O. coffeae*. Handling time was higher on adult females than on larvae. Rate of predation leveled off at temperatures greater than 25° C. Functional responses to prey density at six temperatures and to each life stage of *O. coffeae* approximated the Holling type II model. The oviposition rate increased with prey consumption and temperature. On average, a predator consumed 1.62 adult female prey for every egg it laid. With a fixed number of prey available, predation rate per predator decreased with increased predator density.

**Keywords:** biocontrol, handling time, oviposition, pest, predation

**Correspondence:** <sup>a</sup> [jasinrahman@gmail.com](mailto:jasinrahman@gmail.com), <sup>b</sup> [azariah.babu@gmail.com](mailto:azariah.babu@gmail.com), <sup>c</sup> [roobakkumar@gmail.com](mailto:roobakkumar@gmail.com), <sup>d</sup> [insectakperumalsamy@gmail.com](mailto:insectakperumalsamy@gmail.com), \*Corresponding author

**Editor:** TX Liu was editor of this paper.

**Received:** 13 May 2011, **Accepted:** 31 May 2012

**Copyright :** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 12, Number 125

#### Cite this paper as:

Rahman VJ, Babu A, Roobakkumar A, Perumalsamy K. 2012. Functional and numerical responses of the predatory mite, *Neoseiulus longispinosus*, to the red spider mite, *Oligonychus coffeae*, infesting tea. *Journal of Insect Science* 12:125. Available online: <http://www.insectscience.org/12.125>

## Introduction

Functional and numerical responses give data on the biocontrol efficiency of natural enemies on a particular pest. The functional response of predators describes the relationship between the numbers of prey attacked at different prey densities. Numerical response is defined as the change in a predator's reproductive output at varying prey densities. Since the functional response is an important aspect in the dynamics of a predator-prey relationship and a major component of a population model, it has been used to predict the mechanisms underlying predator-prey behavior to improve the practical predictive potential of predator candidates for biocontrol (Sepúlveda and Carrillo 2008). Holling (1959a; 1959b; 1961) described three types of functional response curves: (1) a linear rise in prey consumption and the rate of reproduction to a plateau (type I), (2) a cyrtoid curve rising at a decreasing rate to a plateau (type II), and (3) a sigmoid curve with a positive accelerating rate up to an inflection point and thereafter a diminishing rate up to the plateau (type III). The basic factors that affect the functional response are: time of exposure of predator to prey, search rate, identification, capture, and prey consumption (Holling 1959b). The ability of the predator to control prey population is influenced by several factors, such as developmental response, utilization of alternative prey, behavioral patterns in response to the predator's own density, and intra-guild interaction (Lester and Harmsen 2002). The present study reports the functional and numerical responses of *Neoseiulus longispinosus* Evans (Acari: Phytoseiidae) on *Oligonychus coffeae* Nietner (Acari: Tetranychidae) in a laboratory. A

laboratory experiment was also conducted to study the effect of mutual interference of predators on searching efficiency in the same feeding area.

While considering the pest scenario of tea in south India, the red spider mite, *O. coffeae* deserves special attention because of its widespread occurrence in all the tea growing areas causing about 18% crop loss (Muraleedharan et al. 2005). High incidence of this mite may occur in dry weather. *O. coffeae* infests the upper-surface of leaves, and, as a result of feeding on chlorophyll, the infested plants defoliate, causing the death of bushes and losses in crops. There is an increasing trend of switching over to integrated pest management practices in tea plantations because of the negative effects of synthetic pesticides used for the control of tea pests. Continuous use of synthetic pesticides causes a disruption of the natural balance between crop, pests and predators, the proliferation of pesticide resistant generations of pests, and undesirable residues in the tea (Rahman et al. 2012). Recent surveys conducted in the south Indian tea plantations (Vandiperiyar and Munnar (Idukki, Kerala), Coonoor and Kotagiri (The Nilgiris, Tamil Nadu), and Anamallais (Coimbatore, Tamil Nadu)) revealed the presence of a predatory mite, *N. longispinosus*, feeding on all stages of *O. coffeae*. Being major predators of Tetranychids and Eriophyids, Phytoseiid mites receive attention in agricultural pest control. The predatory mite *Scapulaseius newsami* (Evans) was introduced from Malaysia to India in 1961 for the control of *Calcarus carinatus*, *Brevipalpus californicus*, and *Oligonychus coffeae* in tea (Sankaran 1974). Zhang et al. (1999) reported the control potential of *N. longispinosus* on

*Schizotetranychus nanjingensis*, a pest of bamboo. *N. longispinosus* was evaluated as a bio control agent (Zhang et al. 1998) and a mass release was done in bamboo forests (Zhang et al. 2000). It was reported as a promising biocontrol agent against citrus yellow mite, *Eotetranychus cendanae* (Thongtab et al. 2001), and two spotted spider mite, *Tetranychus urticae*, on strawberry (Kongchuensin et al. 2001).

Since there is little information on the predator-prey relationship between *N. longispinosus* and *O. coffeae*, an attempt was made to document the functional and numerical responses as well as the mutual interference response of *N. longispinosus* with *O. coffeae* as the prey. Data on functional responses were fitted to Holling's disc equation (Holling 1959a). Data on the numerical responses and oviposition at different temperatures were fitted to regression equations. The data on the response of predator to predator density were fitted to Hassell and Varley's (1969) equation. The values of each experiment can be retrieved from these equations.

## Materials and Methods

### Stock colonies of *O. coffeae* and *N. longispinosus*

Both *O. coffeae* and *N. longispinosus* were collected from the United Planters' Association of Southern India tea experimental farm (Valparai, Coimbatore, Tamil Nadu, India) and used to start the stock colony. Adult red spider mites were transferred onto healthy tea leaves placed on a moist cotton pad, which was laid on the top of a sponge (0.5 inch thick) in plastic trays (42 x 30 x 6.5 cm). Water was added to the rearing unit when necessary to keep the cotton moist and thereby to prevent the leaves from drying.

Deteriorating leaves with red spider mite were cut and placed dorsal surface down on new leaves arranged as described previously to maintain the colony. Gravid females of *N. longispinosus* were collected from the red spider mite infested fields and reared on infested tea leaves placed on a moist cotton pad, which was kept in Petri plates (9 cm diameter). Cotton was kept oversaturated with water to prevent the possible escape of predatory mite. Petri plates were examined every two days to check over-population of the predator, and the leaves without enough spider mites to sustain the predatory mites were cut into small pieces and placed on new red spider mite infested tea leaves in Petri plates. Both the spider mite and predatory mite colony was maintained at  $25 \pm 2^\circ \text{C}$  and  $75 \pm 10\% \text{RH}$ .

### General experimental methodology

The experimental arenas consisted of freshly excised tea leaf discs (2 cm in diameter) placed on moist cotton in Petri plates. Experiments on the functional response to different prey stages, the numerical response to different prey densities, and the response of predators to predator density were conducted at  $25^\circ \text{C}$  and  $75 \pm 5\% \text{RH}$ . The photoperiod for all experiments was 14:10 L:D. Consumed prey in the arena of experiments on functional response were removed at each observation without destructing the web formed by spider mite, and the constant number of prey was maintained by adding new ones till the termination of the experiment. All experiments were conducted for seven consecutive days, and the mean values of all replications on each day were summed and averaged to get the final results of the experiment. The data from the first day of each experiment were excluded from the final analysis to dismiss the possibility of unusual response of the sample of predatory mites to

an abrupt change in temperature, life stage of the prey, and prey density from that of stock colony.

### **Functional response at different temperatures**

Adult females of *N. longispinosus* were transferred from the stock colony onto an excised tea leaf with prey mites and allowed to lay eggs for 12 hours. Then, adults were removed, and the eggs of *N. longispinosus* were allowed to complete their post-embryonic development on the leaf. After mating, females were separated and used in the bioassays when they were 6 days old after hatching. They were starved for 24 hours on tea leaves arranged as described for the rearing of predatory mite. Each leaf disc was then inoculated with 1, 5, 10, 15, 20, 25, or 30 adult females of *O. coffeae*, and a single gravid female of *N. longispinosus* was introduced on each leaf disc. Experiments were conducted at 10, 15, 20, 25, 30, and 35° C with five replications for each experiment. The number of prey consumed per predatory mite was recorded every 24 hours for seven consecutive days.

### **Functional response to different prey stages**

The experimental methods and density of prey were similar to the above except that the functional responses of different life stages of *O. coffeae* (egg, larva, protonymph, deutonymph, and adult female) at 25° C were observed. Experiments on each life stage were replicated five times. The number of prey consumed per predatory mite was recorded every 24 hours for seven consecutive days.

### **Numerical response to different prey densities**

The numerical response of *N. longispinosus* to various densities of *O. coffeae* was studied in terms of the rate of oviposition of an

individual predator to the rate of predation. A six-day-old gravid female predatory mite was starved for 24 hours and confined on a leaf disc with adult female *O. coffeae* at a density of 1, 3, 6, 9, or 12. The number of prey consumed and number of eggs laid by the predator were counted everyday for seven consecutive days with six replicates per prey density. Consumed prey were replaced daily to maintain the original number, and the eggs laid by the predator were removed every day.

### **Numerical response at different temperatures**

Ten adult red spider mites were confined in the experimental arena. A six-day-old gravid female predator was introduced onto the leaf. Experiments were conducted at 10, 15, 20, 25, 30, and 35° C. The number of eggs laid by the predator was recorded everyday for seven consecutive days with six replicates for each temperature. Dead prey were replaced with new ones every day.

### **Response of predator to predator density**

The influence of the mutual interference of predators on prey consumption was evaluated in the laboratory. The number of prey was fixed as 40 females of *O. coffeae* per leaf disc. Adult female predatory mites were introduced to different leaf discs at the density of 1, 3, 5, 7, or 9. The number of prey consumed was counted 24 hours after the start of the experiment. There were ten replicates for each predator density.

### **Data analysis**

Data were analyzed in Microsoft Excel (<http://www.microsoft.com/>). The functional responses were determined by fitting the data to the Holling disc equation (Holling 1959a),  $N_a/P = aNT/1 + aT_hN$ , where  $N_a$  is the number of successful attacks per predator ( $P$ ) during a specific time period ( $T$ ), which in this is case

**Table 1.** Functional response of *Neoseiulus longispinosus* adult females to *Oligonychus coffeae* adults at different temperatures.

Parameters*	10° C	15° C	20° C	25° C	30° C	35° C
<i>Na</i>	0.563 <i>N</i> / 1 + 0.174 <i>N</i>	0.687 <i>N</i> / 1 + 0.196 <i>N</i>	0.991 <i>N</i> / 1 + 0.277 <i>N</i>	1.230 <i>N</i> / 1 + 0.266 <i>N</i>	1.187 <i>N</i> / 1 + 0.204 <i>N</i>	1.183 <i>N</i> / 1 + 0.202 <i>N</i>
<i>Th</i>	0.309	0.285	0.28	0.216	0.172	0.171
<i>a</i>	0.563	0.687	0.991	1.23	1.187	1.183
<i>a/Th</i>	1.819	2.415	3.546	5.685	6.892	6.912
<i>K</i>	3.232	3.514	3.578	4.623	5.807	5.845
<i>R</i> <sup>2</sup>	0.99	0.96	0.92	0.93	0.98	0.97

\**Na* = number of prey consumed; *N* = initial prey density; *Th* = handling time (day); *a* = attack rate; *K* = maximum predation rate; *R*<sup>2</sup> = coefficient of determination.

1 day, *N* is the initial density of the prey, and *a* and *T<sub>h</sub>* are the rate of successful attacks and the time required to handle the prey respectively. Handling time is defined as the time that the predator requires to pursue, kill, and digest the prey (Holling 1963). The parameters *a* and *T<sub>h</sub>* were calculated using a linear regression technique where 1/*Na* was regressed on 1/*N*. *a* is the reciprocal of the slope and *T<sub>h</sub>* is the intercept. The *a/Th* value indicates the effectiveness of predation. Maximum predation rate (*K*) was calculated as *T/Th*.

The results of numerical responses and oviposition at different temperatures were fitted to regression equations. Different regression curves tested to fit the data are presented in this paper. The regression model whose *R*<sup>2</sup> value was closer to 1 was selected to fit the data.

Since the relationship between searching efficiency and predator density is linear, the data of mutual interference of predators were fitted to Hassell and Varley’s (1969) empirical model,  $\log E = \log Q - m \log P$ ; where *E* is the measure of searching efficiency, *P* is the predator density, *Q* is the quest constant (the value of *E* when *P* = 1), and *m* is the interference constant (the slope of the regression of  $\log E$  on  $\log P$ ). Searching efficiency (*E*, a measure of the per capita predation when there is mutual interference between predators in the same feeding area) was calculated as  $1/P \log_e [N/(N - N_a)]$ , where *N* is the initial prey density, and *N<sub>a</sub>* is the

number of prey consumed. Though a regression analysis of  $\log E$  on  $\log P$  to obtain *m* is not valid since *P* is used in the calculation of *E*, *m* could be significant if the corresponding value of *r* (regression coefficient) is significant (Varley et al. 1973).

Results

Functional response at different temperatures

The functional responses of predators at six different temperatures approximated the Holling type II model (Table 1) by a cyrtoid curve rising at a decreasing rate to a plateau. As the temperature increased, handling time (*Th*) decreased and successful attack rate (*a*) increased while a leveled off at temperatures greater than 25° C. The higher values of *a/Th* recorded at the temperatures above 25° C indicate that *N. longispinosus* was more effective against *O. coffeae* at those temperatures.

Functional responses to different prey stages

A cyrtoid curve was observed in the case of functional responses of predators to different prey stages of *O. coffeae*; therefore, it was fitted to the Holling type II model (Table 2). *a/Th* values indicate that *N. longispinosus* was most effective against larvae and nymphs of *O. coffeae*.

Numerical responses to different prey densities

**Table 2.** Functional response of *Neoseiulus longispinosus* adult females to different stages of *Oligonychus coffeae*.

Parameters*	Egg	Larva	Protonymph	Deutonymph	Adult female
<i>Na</i>	0.826 <i>N</i> / 1 + 0.044 <i>N</i>	1.012 <i>N</i> / 1 + 0.015 <i>N</i>	1.022 <i>N</i> / 1 + 0.036 <i>N</i>	1.032 <i>N</i> / 1 + 0.047 <i>N</i>	1.211 <i>N</i> / 1 + 0.237 <i>N</i>
<i>Th</i>	0.053	0.015	0.035	0.046	0.196
<i>a</i>	0.826	1.012	1.022	1.032	1.211
<i>a/Th</i>	15.59	67.47	28.878	22.43	6.181
<i>K</i>	18.87	66.67	28.25	21.74	5.102
<i>R</i> <sup>2</sup>	0.99	0.99	0.99	0.98	0.96

\**Na* = number of prey consumed; *N* = initial prey density; *Th* = handling time (day); *a* = attack rate; *K* = maximum predation rate; *R*<sup>2</sup> = coefficient of determination.

The number of eggs laid per predator per day (y) was found to increase, with a gradual leveling off, with prey density (x) (Figure 1). The relationship can be expressed by the equation  $y = -0.023x^2 + 0.537x - 0.574$ , ( $R^2 = 0.992$ ) (Figure 2). At the density of 12 *O. coffeae* females per leaf disc, *N. longispinosus* laid a maximum of 2.52 eggs per day. No eggs were laid at the density of 1 red spider mite per leaf.

Oviposition rate (y) increased linearly with predation rate (x) (Figure 1); this relationship can be expressed by the equation,  $y = 0.725x - 0.26$ , ( $R^2 = 0.986$ ) (Figure 3).

**Numerical response at different temperatures**

The rate of oviposition increased with an increase in temperature (Figure 4). This relationship can be expressed by the equation  $y = 0.003x^2 + 0.033x - 0.332$ , ( $R^2 = 0.987$ ), where y is the number of eggs laid by a predator per day and x is the temperature (Figure 5).

**Response of predators to predator density**

Predation rate per predator decreased with an increase in predator density with a constant number of prey (40) available. The per capita rate of predation (per day) of *N. longispinosus* females on *O. coffeae* females decreased from 5.5 to 2.01 in relation to the predator density increasing from 1 to 9. The relationship between searching efficiency and predator density was found to be linear, which can be

expressed by the equation  $\log E = \log 0.148 - 1.481 \log P$  ( $R^2 = 0.993$ ).

**Discussion**

In this study, the predatory mite *N. longispinosus* was found to consume all stages of *O. coffeae*. The maximum predation rate is limited by an upper asymptote defined by the ratio  $T/Th$  (*K*) (Hassell 1978), which was higher at 30 and 35° C for *N. longispinosus*. This result is desirable for the biocontrol of *O. coffeae* because the maximum daily temperature in the summer, during which *O. coffeae* is present, fluctuates around 30° C in south Indian tea plantations. When tested with different prey stages, larvae and nymphs of *O. coffeae* were more consumed than other stages; therefore, *K* value was higher for larvae and nymphs. Blackwood et al. (2001) observed a strong preference of *N. longispinosus* for the egg stage of *T. urticae*. However, Ibrahim and Palacio (1994) noted that *N. longispinosus* preferred larvae and nymphs of *Tetranychus urticae* to eggs. The predatory mites *N. californicus* (Canlas et al. 2006) and *Kampimodromus aberrans* (Kasap and Atlihan 2010) also showed preference for larvae when fed *T. urticae*.

The preference for younger stages will be helpful in preventing the proliferation of pest population. Predation rate was found to increase with increased density of prey. The reason for increased predation rate at higher prey densities may be the interference or



disturbance by the prey as noticed in *Amblyseius largoensis* (Muma), *Euseius corcordis* (Chant), and *Galendromus* (= *Typhlodromus*) *helveolus* (Chant) (Sandness and Mc Murtry 1970). At higher prey densities, predators may spend less time on individual prey because prey accidentally bumping into a feeding predator may cause the predator to abandon the prey it is eating and attack another. The bumping of prey into each other may cause prey to move, thereby increasing the chance of an encounter with a predator. This higher prey density results in wasteful killing (Metz et al. 1988), as a predator may partially eat multiple prey instead of eating one whole prey. Predation rate reaches a plateau at higher prey densities due to factors such as satiation (Sabelis 1985). The number of prey consumed by *N. longispinosus* increased with increased density of *T. urticae* and leveled off at a prey density of 40 per predator (Ibrahim and Rahman 1997). The density dependant interference or defensive behavior of spider mites may also have caused the reduced rate of successful attacks.

The oviposition rate of *N. longispinosus* appears to depend on temperature and the quality of prey taken as food. The rate of oviposition increased with increased prey consumption. Oviposition correlates with predation because phytoseiid mites allocate a major fraction of food ingested to egg production (Sabelis and Janssen 1994). At the density of one prey per leaf, predators survived but hardly laid any eggs. On average, one egg was laid for every 1.62 prey eaten. The increase in oviposition in response to the increase in prey density adds to the effectiveness of biological control by causing an increase in predator population. The rate of oviposition increased with an increase in temperature from 10 to 30° C. In a previous

study, it was shown that *N. longispinosus* could not survive at temperatures above 35° C (unpublished data).

An increase in the population size of a predator may cause a decrease in prey consumption. Mutual interference denotes the adverse influence of predator density on the instantaneous success of individual predator (Arditi and Akcakaya 1990). However, spatial complexity and aggregation also have roles in predator-prey interactions (Hassell and May 1973; Free et al. 1977). Under field conditions, interference probably leads to the emigration of individual predators from regions of high parasite density (Rogers and Hassell 1974). Begon et al. (1996) studied mutual interference with an unlimited number of prey. But 40 prey is large for predatory mites as far as their predation rate is concerned. So the effects of mutual interference will not be mixed up with the effects of competition for food. In the present study, predator interference affected the rate of prey consumption. However, the natural dispersal of predators to explore for more prey will reduce this interference behavior among predators in field.

Results reported here form a database for further studies on the feasibility to incorporate *N. longispinosus* in integrated pest management programs against *O. coffeae* in tea.

## Acknowledgement

The financial support for this work was provided by the Tea Board, Ministry of Commerce and Industry, Government of India, under XI five-year plan. Thanks are due to Dr. K. Ramaraju, Professor of Entomology, Tamil Nadu Agricultural University, and Dr. Mary Anithalatha Sadanandan, Head of the



Post Graduate and Research Department of Zoology, Malabar Christian College, Calicut, for confirming the taxonomic identity of the predatory mite.

## References

- Arditi R, Akcakaya HR. 1990. Underestimation of mutual interference of predators. *Oecologia* 83: 358-361
- Begon M, Harper JL, Townsend CR. 1996. *Ecology: Individuals, Populations, Communities*, third edition. Blackwell Scientific Publications.
- Blackwood JS, Schausberger P, Croft BA. 2001. Prey- stage preference in generalist and specialist Phytoseiid mites (Acari: Phytoseiidae) when offered *Tetranychus urticae* (Acari: Tetranychidae) eggs and larvae. *Environmental Entomology* 30: 1103-1111
- Canlas LJ, Amano H, Ochiai N, Takeda M. 2006. Biology and predation of the Japanese strain of *Neoseiulus californicus* (McGregor) (Acari: Phytoseiidae). *Systematic and Applied Acarology* 11: 141-157
- Free CA, Beddington JR, Lawton JH. 1977. On the inadequacy of simple models of mutual interference for parasitism and predation. *Journal of Animal Ecology* 46: 543-554
- Hassell MP. 1978. *Dynamics of arthropod predator-prey systems*. Princeton University Press.
- Hassell MP, May RM. 1973. *Stability in insect host-parasite models*. *Journal of Animal Ecology* 42: 693-726
- Hassell MP, Varley GC. 1969. New inductive population model for insect parasites and its bearings on biological control. *Nature* 223: 1133-1137
- Holling CS. 1959a. The components of predation as revealed by a study of small mammal predation of European pine saw fly. *Canadian Entomologist* 91: 293-320
- Holling CS. 1959b. Some characteristics of simple types of predation and parasitism. *Canadian Entomologist* 91: 385-398
- Holling CS. 1961. Principles of insect predation. *Annual Review of Entomology* 6: 163-182
- Holling CS. 1963. An experimental component analysis of population process. *Memoirs of Entomological Society of Canada* 32: 22-32
- Ibrahim YB, Palacio VB. 1994. Life history and demography of the predatory mite, *Amblyseius longispinosus* Evans. *Experimental and Applied Acarology* 18: 361-369
- Ibrahim YB, Rahman RBA. 1997. Influence of prey density, species and developmental stages on the predatory behaviour of *Amblyseius longispinosus* (Acari: Phytoseiidae). *Entomophaga* 42: 319-327
- Kasap I, Atlihan R. 2010. Consumption rate and functional response of the predaceous mite *Kampimodromus aberrans* to two-spotted spider mite *Tetranychus urticae* in the laboratory. *Experimental and Applied Acarology* 53: 253-261
- Kongchuensin M, Charanasri V, Kulpiyawat T, Khantonthong P. 2001. Biological control

of two spotted spider mite in strawberries by the predatory mite, *Amblyseius longispinosus* (Evans) (Acari: Phytoseiidae). In: Halliday RB, Walter DE, Proctor HC, Norton RA, Colloff MJ, Editors. *Acarology: Proceedings of the 10<sup>th</sup> International Congress*. pp. 513-517.

Lester P, Harmsen R. 2002. Functional and numerical responses not always indicate the most effective predator for biological control: an analysis of two predators in a two-prey system. *Journal of Applied Ecology* 39: 455-468

Metz JAJ, Sabelis MW, Kuchlein JH. 1988. Sources of variation in predation rate at high prey densities: an analytic model and a mite example. *Experimental and Applied Acarology* 5: 187-205

Muraleedharan N, Sudarmani DNP, Selvasundaram R. 2005. Bioecology and management of the red spider mite infesting tea in south India. In: *Proceedings of International Symposium on Innovation in Tea Science and Sustainable Development In Tea industry*. pp. 756-766. China Tea Science Society.

Rahman VJ, Babu A, Roobakkumar A, Perumalsamy K, Vasanthakumar D, Subramaniam MSR. 2012. Efficacy, prey stage preference and optimum predator-prey ratio of the predatory mite, *Neoseiulus longispinosus* Evans (Acari: Phytoseiidae) to control the red spider mite, *Oligonychus coffeae* Nietner (Acari: Tetranychidae) infesting tea. *Archives of Phytopathology and Plant Protection* 45: 699-706

Rogers DJ, Hassell MP. 1974. General models for insect parasite and predator searching

behavior: Interference. *Journal of Animal Ecology* 43: 239-253

Sabelis MW. 1985. Predation on spider mites. In: Helle W, Sabelis MW, Editors. *Spider mites. Their Biology, Natural Enemy and Control*. pp. 103-127. Elsevier.

Sabelis MW, Janssen A. 1994. Evolution of life-history patterns in the Phytoseiidae. In: Houck MA, Editor. *Mites*. pp. 70-98. Chapman & Hall.

Sandness JN, Mc Murtry JA. 1970. Functional response of three species of Phytoseiidae (Acarina) to prey density. *Canadian Entomologist* 102: 692-704

Sankaran T. 1974. Natural enemies introduced in recent years for biological control of agricultural pests in India. *Indian Journal of Agricultural Sciences* 44: 425-433

Sepuúlveda F, Carrillo R. 2008. Functional response of the predatory mite *Chileseius camposi* (Acari: Phytoseiidae) on densities of its prey, *Panonychus ulmi* (Acarina: Tetranychidae). *Revista de Biología Tropical* 56: 1255-1260

Thongtab T, Chandrapatya A, Baker GT. 2001. Biology and efficacy of the predatory mite, *Amblyseius longispinosus* (Evans) (Acari: Phytoseiidae) as a biological control agent of *Eotetranychus cendanai* Rimando (Acari: Tetranychidae). *Journal of Applied Entomology* 125: 543-549


Varley GC, Gradwell GR, Hassell MP. 1973. *Insect Population Ecology: An Analytical Approach*. Blackwell Scientific Publication.

Zhang YX, Zhang ZQ, Lin JZ, Liu Q. 1998. Predation of *Amblyseius longispinosus* (Acari:

Phytoseiidae) on *Aponychus corpuzae* (Acari: Tetranychidae). *Systematic and Applied Acarology* 3: 53-58

Zhang YX, Zhang ZQ, Ji J, Lin JZ. 1999. Predation of *Amblyseius longispinosus* (Acari: Phytoseiidae) on *Schizotetranychus nanjingensis* (Acari: Tetranychidae), a spider mite injurious to bamboo in Fujian, China. *Systematic and Applied Acarology* 4: 63-68

Zhang YX, Zhang ZQ, Lin JZ, Ji J. 2000. An overview of occurrence, development and damage of bamboo mites and their integrated management in Fujian, China. *Systematic and Applied Acarology* (special publication) 4: 9-17



**Figure 1.** Numerical responses (mean  $\pm$  SE) of *Neoseiulus longispinosus* female to different density of *Oligonychus coffeae* females. High quality figures are available online.

