

Geographic variation in body mass of rock ptarmigan *Lagopus mutus* in the Alps and the Pyrenees

Authors: Brenot, Jean-François, Ellison, Laurence, Rotelli, Luca, Novoa, Claude, Calenge, Clément, et al.

Source: *Wildlife Biology*, 11(4) : 281-285

Published By: Nordic Board for Wildlife Research

URL: [https://doi.org/10.2981/0909-6396\(2005\)11\[281:GVIBMO\]2.0.CO;2](https://doi.org/10.2981/0909-6396(2005)11[281:GVIBMO]2.0.CO;2)

BioOne Complete (complete.bioone.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Geographic variation in body mass of rock ptarmigan *Lagopus mutus* in the Alps and the Pyrenees

Jean-François Brenot, Laurence Ellison, Luca Rotelli, Claude Novoa, Clément Calenge, Patrick Léonard & Emmanuel Ménoni

Brenot, J-F., Ellison, L., Rotelli, L., Novoa, C., Calenge, C., Léonard, P. & Ménoni, E. 2005: Geographic variation in body mass of rock ptarmigan *Lagopus mutus* in the Alps and the Pyrenees. - *Wildl. Biol.* 11: 281-285.

We compared the body mass of adult rock ptarmigan *Lagopus mutus* shot in autumn in the French Pyrenees (Ariège), the French Alps (Hautes-Alpes) and the Italian Alps (Province of Verbania). Mean body mass of adults was greater in the Ariège than in the Hautes-Alpes both for cocks (466 vs 433 g) and hens (429 vs 406 g), with the body mass of Italian birds being intermediate (cocks 448 g, hens 418 g). The relatively large mass of rock ptarmigan in the Pyrenees contrasts with the geographic trends found in capercaillie *Tetrao urogallus* and grey partridge *Perdix perdix* in France. For the last two species, birds in the Pyrenees are smaller than those further north. The site differences that we observed in body mass of rock ptarmigan may be related to differences in availability of high quality foods, particularly ericaceous shrubs. However, at the continental scale within Europe, the body mass of rock ptarmigan appears to be greatest in northern latitudes.

Key words: autumn, body mass, France, geographic variation, Italy, *Lagopus mutus*, rock ptarmigan

Jean-François Brenot & Claude Novoa, Office National de la Chasse et de la Faune Sauvage, Espace Alfred Sauvy, F-66500 Prades, France - e-mail: prades@oncfs.gouv.fr

Laurence Ellison, Dolgue, F-34270 Claret, France - e-mail: larry.ellison@wanadoo.fr

Luca Rotelli, University of Bern, Zoological Institute, Conservation Biology, Baltzerstrasse 6, CH- 3012 Bern, Switzerland - e-mail: luca.rotelli@libero.it

Clément Calenge, LBBE - UMR CNRS 5558, Université Claude Bernard Lyon 1, 43 boulevard du 11 novembre 1918, F-69622 Villeurbanne cedex, France - e-mail: calenge@biomserv.univ-lyon1.fr

Patrick Léonard, Office National de la Chasse et de la Faune Sauvage, Micropolis, F-05000 Gap, France - e-mail: p.leonard@oncfs.gouv.fr

Emmanuel Ménoni, Office National de la Chasse et de la Faune Sauvage, Impasse de la Chapelle, F-31800 Villeneuve de Rivière, France - e-mail: e.menoni@oncfs.gouv.fr

Corresponding author: Jean-François Brenot

Article accepted for publication in Game and Wildlife Science (GWS) prior to the merger between WILDLIFE BIOLOGY and GWS. Therefore, this article has not been peer-reviewed by WILDLIFE BIOLOGY Associate Editors

Data on body mass in tetraonids are useful for investigating taxonomic relationships (Kahn et al. 1999), and for explaining the evolution of life history traits, including social organisation and demographic parameters (Wiley 1974, Trouvilliez et al. 1987, Sæther & Andersen 1988, Stearns 1992). In grouse, body mass may also be

Table 1. Models employed to first test the effect of date of shooting (D) on body mass of rock ptarmigan mass and then the effects of sex (S) and location (L). Y(L) = year (nested in location), + = additive effect, * = interactions.

Model	Tested Effect	F	df	P
S + L + S*L + D + S*D + L*D + S*L*D	S*L*D	0.07	2, 165	0.931
S + L + S*L + D + S*D + L*D	L*D	0.23	2, 167	0.794
S + L + S*L + D + S*D	S*D	0.0001	1, 169	0.994
S + L + S*L + D	D	1.11	1, 170	0.293
S + L + S*L + Y(L) + S*Y(L)	S*Y(L)	0.63	14, 137	0.836
S + L + S*L + Y(L)	Y(L)	1.59	20, 151	0.061
S + L + S*L	S * L	0.46	2, 171	0.634
S + L	S	39.08	1, 173	< 0.001
S + L	L	10.89	2, 173	< 0.001

helpful in interpreting the effects of environmental stress on individuals (Sherfy & Pekins 1994). Such studies require accurate information on both inter- and intraspecific variation in body mass. In this paper we report on differences in body mass of two subspecies of rock ptarmigan in three areas: *Lagopus mutus pyrenaicus* in the French Pyrenees and *L. m. helveticus* in the French and Italian Alps.

Material and methods

Data collection

In France, we examined 53 adult rock ptarmigan shot by hunters during 1994-1999 in four communes in the Pyrenees, department of the Ariège (~42°30'N, 1°30'E) and 51 birds shot during 1979-1996 in nine communes in the Alps, department of the Hautes-Alpes (~44°30'N, 6°45'E). In Italy, the sample included 73 adults shot during 1997-2001 in two hunting districts in the Province of Verbania (~46°00'N, 8°30'E). We did not consider juveniles (3-4 months old) because they only occurred in small numbers in the shooting bags.

Dates of shooting varied from 15 September to 12 October in the Ariège and from 9 September to 11 November in the Hautes-Alpes. Shooting dates in Italy were 1 October to 28 November.

In France, we determined the sex of most birds by internal examination of the gonads. Age (adult or juvenile) was established by the presence or absence of the bursa of Fabricius. In the case of 18 birds not examined internally, age was determined by colour of primary wing feathers (Ellison & Léonard 1996), and sex was assessed by external plumage characteristics, including the colour and pattern of barring on body feathers. Jean-François Brenot verified this method of sex determination by examining the gonads of several birds.

In Italy, birds were not examined internally. Sex was determined by the presence or absence of the black loreal

stripe. This eye stripe appears in late September in males and is a reliable indicator of sex from October to May (Weeden 1964). Age was assessed from the wear and pigmentation of the two outermost primary wing feathers (Bocca 1984).

In all locations, body mass of birds, including the crop and its contents, was measured to the nearest 5 g either by us or by field assistants that we had trained.

Data analysis

We analysed the variations in body mass using linear models (McCullagh & Nelder 1989). Because birds might gain or lose mass during autumn, we studied the effect of date of shooting on mass by fitting the model: $\text{Mass} = \text{Sex} + \text{Location} + \text{Sex} * \text{Location} + \text{Date} + \text{Sex} * \text{Date} + \text{Location} * \text{Date} + \text{Sex} * \text{Location} * \text{Date}$. We then tested the effects written in italics with an analysis of variance, beginning with the interactions. In a second analysis, we studied the effects of sex and location by fitting the model: $\text{Mass} = \text{Sex} + \text{Location} + \text{Sex} * \text{Location}$




Figure 1. Body mass (in g) of adult rock ptarmigan in the French Pyrenees (Ariège), the Italian Alps (province of Verbania) and the French Alps (Hautes-Alpes) with error bars indicating ± 1 standard error.

Table 2. Mean body mass (in g), standard error (SE) and range of adult rock ptarmigan in the French Pyrenees (Ariège), the French Alps (Hautes-Alpes) and the Italian Alps (Province of Verbania).

Sex	Ariège				Hautes-Alpes				Italy			
	Mean	SE	Range	N	Mean	SE	Range	N	Mean	SE	Range	N
Cocks	466	5.50	400-540	32	433	5.92	360-475	22	448	4.36	385-520	56
Hens	429	7.93	350-485	21	406	5.47	350-460	29	418	5.97	385-470	17

+ Year(Location) + Sex*Year(Location). Note that Year was nested within Location, i.e. it was assumed that a given year did not have the same effect in different locations. In this model, we first tested all interactions and then the main effects. We conducted all analyses using R software (R Development Core Team 2004).

Results

In the first analysis, after we withdrew the non-significant interaction terms, the effect of date was not significant (Table 1). Because mass showed no trend during the hunting season, we did not retain date in the second analysis. In the second analysis, the effect of year within location was nearly significant (see Table 1), but we removed this effect from the model because the P value of 0.061 was caused by one unusually small hen (350 g), which was the only hen weighed in 1998 in the Ariège. We also removed the other non-significant interaction terms from the initial model, allowing us to test for sex and location, both of which were highly significant (see Table 1). In the model Sex + Location, variances were homogeneous in cocks and hens, as well as in the three locations. Residuals of the model were also normally distributed. We conclude that body mass of both cocks and hens is greater in the Ariège than in the Hautes-Alpes, with birds in Italy occupying an intermediate position (Fig. 1, Table 2). The body mass of cocks exceeded that of hens by about 30 g in all three areas.

Discussion

Our data on mean body mass of adult rock ptarmigan in the Hautes-Alpes are similar to those reported by Couturier (1964) for adults shot in autumn in the French Alps (Table 3). Mass of birds in the Italian Province of Verbania is similar to that of ptarmigan shot about 120 km to the east in the Italian Province of Sondrio (Scherini 1984; see Table 3). The Italian ptarmigan appear to be about 10-20 g heavier than those in the Hautes-Alpes. Couturier (1964) reported that rock ptarmigan were heavier in the Pyrenees than in the French Alps, and our results suggest a difference of about 20-30 g. Nevertheless, we believe that other populations in both mountain ranges should be sampled before concluding that this difference characterises the subspecies *L. m. pyrenaicus* and *L. m. helveticus*.

Within subspecies of rock ptarmigan and spruce grouse *Dendragapus canadensis* in North America, birds of northern populations tend to be heavier or larger in body dimensions than birds of southern populations (Browning 1979, Holder & Montgomerie 1993, Boag & Schroeder 1992). These trends coincide with Bergmann's rule, which states that mass is higher in cooler climates because small animals lose heat faster than large animals. However there are exceptions in tetraonids. For example, the opposite latitudinal trend occurs in ruffed grouse *Bonasa umbellus* (Rusch et al. 2000). Likewise, in our study, birds in the Ariège are heavier than those further north in the Hautes-Alpes and in Italy, despite

Table 3. Mean body mass (in g) and range (figures in parentheses) of adult cocks and hens of four subspecies of rock ptarmigan in autumn in Europe.

Subspecies	Area	Reference	Months	Adult cocks	Adult hens
<i>L. m. pyrenaicus</i>	French Pyrenees (Ariège)	This study	September - October	466 (400-540)	429 (350-485)
<i>L. m. helveticus</i>	French Alps (Hautes-Alpes)	This study	September - early November	433 (360-475)	406 (350-460)
<i>L. m. helveticus</i>	French Alps	Couturier (1964)	Most in October - early November	428 (375-515) N = 39	401 (347-471) N = 22
<i>L. m. helveticus</i>	Italian Alps (Verbania)	This study	October- November	448 (385-520)	418 (385-470)
<i>L. m. helveticus</i>	Italian Alps (Sondrio)	Scherini (1984)	September-November	449 (360-520) N = 42	414 (350-455) N = 26
<i>L. m. millaisi</i>	Scotland	Watson (1987)	September-November	530 ¹	490 ¹
<i>L. m. hyperboreus</i>	Norway	Mortensen et al. (1985)	September-November	514 ²	465 ²

¹ Estimated from graph of Figure 1 in Watson (1987).

² Estimated from graph of Figure 1 in Mortensen et al. (1985).

the generally colder temperature in the Alps (Izard et al. 1985). On the other hand, the greater mass of *L. m. mil-laisi* in Scotland and of *L. m. hyperboreus* in Norway (see Table 3) suggests that at a continental scale in Europe, mass may increase with latitude.

The relatively large mass of rock ptarmigan in the Pyrenees does not correspond to the geographical trends observed in two other galliformes in France. The subspecies of capercaillie inhabiting the Pyrenees *Tetrao urogallus aquitanicus* is smaller than *T. u. major* living further north in the Jura and the Vosges (Ménoni & Corti 2000). Similarly, the grey partridge of the Pyrenees *Perdix perdix hispaniensis* is smaller than the subspecies *P. p. perdix* inhabiting the plains further north in France (Lescourret et al. 1987). The relationship in the partridges cannot be a simple matter of temperature because the climate is colder in the Pyrenean Mountains than in the lowland plains.

To explain the differences reported here, we believe that mass will have to be measured in many populations in the Pyrenees and the Alps because both environment (e.g. soil fertility, food quality, temperature) and genetics likely contribute to variation in body mass. Indeed habitat quality may contribute to large body size in the Pyrenees. Nearly all birds from the Ariège were shot in the commune of Aston, where summer and autumn habitats are comprised of areas with a rich ground cover of herbaceous and ericaceous plants, including bilberry *Vaccinium myrtillus*, bog whortleberry *V. uliginosum* and heather *Calluna vulgaris*. Bilberry is a nutrient-rich food (Moss & Watson 1984) that, if available, is selected by rock ptarmigan in all seasons (Watson 1964, Martinez 1985, Boudarel & Garcia-Gonzalez 1991). Bog whortleberry is the only ericaceous plant readily available to rock ptarmigan in the Hautes-Alpes (Bernard 1982), where the habitat tends to be composed of a sparse cover of sedges, grasses and decumbent woody plants, for example mountain azalea *Loiseleuria procumbens*, along with large expanses of stony ground (scree) and boulder fields. These habitats are also often heavily grazed by sheep, which may reduce food available to ptarmigan. On the average, habitats in the Italian Province of Verbania tend to be of intermediate quality. Birds were harvested on both poor areas resembling those in the Hautes-Alpes and on better areas where the vegetation is dominated by bilberry, whortleberry, willow *Salix* spp. and rhododendron *Rhododendron ferrugineum*.

Finally, it would be interesting to investigate in further studies if differences in body mass of rock ptarmigan in the Pyrenees and the Alps are correlated with demographic parameters. Across species of tetraonids, incubation period and clutch mass (egg mass times

clutch size), but not clutch size, increase with body mass (Sæther & Andersen 1988) and within galliformes low population turnover rates, and thus low sustained harvests, tend to be associated with high body mass (Trouvilliez et al. 1987).

Acknowledgements - we thank the numerous hunters in France and Italy who presented their birds for examination. We appreciate the constructive comments of Ariane Bernard-Laurent on an earlier draft of the manuscript, and those of three reviewers.

References

- Bernard, A. 1982: Note sur le régime alimentaire automnal du lagopède alpin (*Lagopus mutus* Montin) dans les Alpes du Sud. - Bulletin Mensuel Office National de la Chasse 55: 20-23. (In French).
- Boag, D.A. & Schroeder, M.A. 1992: Spruce grouse. - In: Poole, A., Stettenheim, P. & Gill, F. (Eds.); The Birds of North America, No. 5. The Academy of Natural Sciences, Philadelphia, The American Ornithologists' Union, Washington, D.C., 28 pp.
- Bocca, M. 1984: Pernice bianca, fagiano di monte, coturnice. Determinazione del sesso e dell'età dei capi abbattuti durante la stagione venatoria autunnale: Criteri generali per la determinazione dell'età. Regione Autonoma Valle d'Aosta. - Assessorato Agricoltura, Foreste e Ambiente Naturale. Servizio tutela dell'ambiente naturale e delle foreste, 4 pp. (In Italian).
- Boudarel, P. & Garcia-Gonzalez, R. 1991: Approche du régime alimentaire du lagopède alpin (*Lagopus mutus pyrenaicus*) dans les Pyrénées occidentales - printemps, été et automne. - Acta biologica montana 10: 11-23. (In French).
- Browning, M.R. 1979: Distribution, geographic variation, and taxonomy of *Lagopus mutus* in Greenland and northern Canada. - Dansk ornitologisk Forenings Tidsskrift 73: 29-40.
- Couturier, M. 1964: Le gibier des montagnes françaises. Deuxième édition. - Bellegarde, Arthaud, 471 pp. (In French).
- Ellison, L. & Léonard, P. 1996: Validation d'un critère d'âge chez le lagopède alpin (*Lagopus mutus*) et sexe et âge ratios dans des tableaux de chasse des Alpes et des Pyrénées. (In French with an English summary: Validation of determining age of rock ptarmigan (*Lagopus mutus*) from wings, and sex and age ratios in shooting bags in the Alps and Pyrenees.) - Gibier Faune Sauvage 13: 1495-1509.
- Holder, K. & Montgomerie, R. 1993: Rock ptarmigan. - In: Poole, A. & Gill, F. (Eds.); The Birds of North America, No. 51. The Academy of Natural Sciences, Philadelphia, The American Ornithologists' Union, Washington, D.C., 24 pp.
- Izard, M., Casanova, H., Devau, B. & Pautou, G. 1985: Con-

- tinentalité et notion de zone interne dans les Pyrénées. (In French with an English summary: Continental climate and the notion of an internal zone in the Pyrenees.) - *Acta Oecologiae Generalis* 6: 317-344.
- Kahn, N.W., Braun, C.E., Young, J.R., Wood, S., Mata, D.R. & Quinn, T.W. 1999: Molecular analysis of genetic variation among large- and small-bodied sage grouse using mitochondrial control-region sequences. - *Auk* 116: 819-824.
- Lescourret, F., Birkan, M. & Novoa, C. 1987: Aspects particuliers de la morphologie de la perdrix grise des Pyrénées, *Perdix perdix hispaniensis* R., et comparaison avec la perdrix grise de Beauce, apparentée à *Perdix perdix* L. (In French with an English summary: Morphological characteristics of the Pyrenean grey partridge, *Perdix perdix hispaniensis* R., compared to the Beaucean grey partridge, related to *Perdix perdix* L.) - *Gibier Faune Sauvage* 4: 49-66.
- Martinez, C. 1985: Autumn food of the ptarmigan (*Lagopus mutus* Montin, 1776) in the Spanish Central Pyrenees. - *Doñana, Acta Vertebrata* 12: 174-181.
- McCullagh, P. & Nelder, J.A. 1989: Generalized linear models. Second Edition. - Chapman & Hall, London, 511 pp.
- Ménoni, E. & Corti, R. 2000: Le grand tétras. - Brochure technique, Office National de la Chasse, Paris, 36 pp. (In French).
- Mortensen, A., Nordøy, E.S. & Blix, A.S. 1985: Seasonal changes in the body composition of the Norwegian rock ptarmigan *Lagopus mutus*. - *Ornis Scandinavica* 16: 25-28.
- Moss, R. & Watson, A. 1984: Maternal nutrition, egg quality and breeding success of Scottish ptarmigan *Lagopus mutus*. - *Ibis* 126: 212-220.
- R Development Core Team 2004: R: A language and environment for statistical computing. - R Foundation for Statistical Computing, Vienna, Austria. Available at: <http://www.R-project.org>
- Rusch, D.H., Destefano, S., Reynolds, M.C. & Lauten, D. 2000: Ruffed grouse. - In: Poole, A. & Gill, F. (Eds.); *The Birds of North America*, No. 515. Philadelphia, Pennsylvania, 28 pp.
- Sæther, B-E. & Andersen, R. 1988: Ecological consequences of body size in grouse Tetraonidae. - *Fauna Norvegicus Series C, Cinclus* 11: 19-26.
- Scherini, G.C. 1984: La pernice bianca. Studio dell'alimentazione della pernice bianca in Provincia di Sondrio (Alpi Retiche e Orobie). (In Italian with an English summary: The rock ptarmigan: food habits in the Province of Sondrio.) - *Unione Nazionale Cacciatori Zona Alpi*, Vicenza, 52 pp.
- Sherfy, M.H. & Pekins, P.J. 1994: The influence of season, temperature, and absorptive state on sage grouse metabolism. - *Canadian Journal of Zoology* 72: 898-903.
- Stearns, S.C. 1992: The evolution of life histories. - Oxford University Press, Oxford, 249 pp.
- Trouvilliez, J., Gaillard, J-M., Allaine, D. & Pontier, D. 1987: Stratégies démographiques et gestion des populations chez les oiseaux: particularités des galliformes. (In French with an English summary: Demographic strategies and management of bird populations: characteristics of galliformes.) - *Colloque Galliformes de Montagne*. Office National de la Chasse & Service de la Recherche, des Études et du Traitement de l'Information sur l'Environnement, Grenoble, pp. 37-51.
- Watson, A. 1964: The food of rock ptarmigan (*Lagopus mutus*) in Scotland. - *The Scottish Naturalist* 71: 60-66.
- Watson, A. 1987: Weight and sexual cycle of Scottish rock ptarmigan. - *Ornis Scandinavica* 18: 231-232.
- Weeden, R.B. 1964: Spatial separation of sexes in rock and willow ptarmigan in winter. - *Auk* 81: 534-541.
- Wiley, R.H. 1974: Evolution of social organization and life-history patterns among grouse. - *Quarterly Review of Biology* 49: 201-227.