

TWO THEILERIA CERVI SSU RRNA GENE SEQUENCE TYPES FOUND IN ISOLATES FROM WHITE-TAILED DEER AND ELK IN NORTH AMERICA

Authors: Chae, Joon-seok, Waghela, Suryakant D., Craig, Thomas M., Kocan, Alan A., Wagner, Gerald G., et al.

Source: Journal of Wildlife Diseases, 35(3) : 458-465

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-35.3.458>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

TWO *THEILERIA CERVI* SSU rRNA GENE SEQUENCE TYPES FOUND IN ISOLATES FROM WHITE-TAILED DEER AND ELK IN NORTH AMERICA

Joon-seok Chae,^{1,2} Suryakant D. Waghela,¹ Thomas M. Craig,¹ Alan A. Kocan,³ Gerald G. Wagner,¹ and Patricia J. Holman^{1,4}

¹ Texas Agricultural Experiment Station, Department of Veterinary Pathobiology, The Texas Veterinary Medical Center, Texas A&M University, College Station, Texas 77843-4467, USA

² Current address: Department of Medicine and Epidemiology, University of California, Davis, California 95616, USA

³ Department of Infectious Diseases and Physiology, College of Veterinary Medicine, Oklahoma State University, Stillwater, Oklahoma 74074, USA

⁴ Corresponding author (e-mail: pholman@cvm.tamu.edu)

ABSTRACT: Two *Theileria cervi* SSU rRNA gene sequence Types, F and G, from white-tailed deer (*Odocoileus virginianus*) and elk (*Cervus elaphus canadensis*) isolates in North America were confirmed. Previously, nucleotide sequencing through a single variable (V4) region showed the presence of SSU rRNA gene Types F and G in *T. cervi* isolates from white-tailed deer and an elk. In this study, both sequence types were found in four *T. cervi* isolates (two from deer and two from elk). Microheterogeneity only appeared in the Type G gene, resulting in Subtypes G1, G2 and G3. Subtype G1 was found in two elk and one white-tailed deer *T. cervi* isolate; Subtypes G2 and G3 were found in a white-tailed deer *T. cervi* isolate. The Type F SSU rRNA genes were identical in nucleotide sequence in both elk and white-tailed deer *T. cervi* isolates. The high degree of conservation in the Type F variable regions may be exploited to design specific oligonucleotide primers for parasite detection by the polymerase chain reaction in cervine or tick hosts.

Key words: *Cervus elaphus canadensis*, elk, gene amplification, nucleotide sequence, *Odocoileus virginianus*, small subunit ribosomal RNA gene, *Theileria cervi*, white-tailed deer.

INTRODUCTION

Theileria spp., the causative agents of theileriosis, are cosmopolitan hemoprotozoan parasites of domestic and wild mammals. The prevalence of the organism is dependent upon the geographic range of the vector tick. Some *Theileria* spp., such as *Theileria parva* and *Theileria annulata* of cattle in Africa, are highly pathogenic and infections may result in devastating losses. Other less pathogenic species, such as *Theileria sergenti* of cattle in the Middle East, cause mild to moderate clinical theileriosis. In the United States, the *Theileria* spp. infecting domestic and wild ruminants are generally considered benign. Specifically, *Theileria cervi* infections in deer are generally considered nonpathogenic, although clinical disease does occur in hosts debilitated by other parasitic burdens or malnutrition (Kocan and Kocan, 1991).

In 1962 the *Theileria* sp. found in white-tailed deer (*Odocoileus virginianus*) in the

United States was designated *Theileria cervi* (Schaeffler, 1961, 1962). *Theileria cervi* has since been identified in white-tailed deer in Texas, Oklahoma, Missouri, Arkansas, and Alabama (reviewed by Kingston, 1981). *Theileria* sp. organisms indistinguishable from *T. cervi* also have been reported in mule deer (*Odocoileus hemionus*) and in exotic axis (*Axis axis*) and sika (*Cervus nippon nippon*) deer in Texas (Laird et al., 1988; Waldrup et al., 1989). The primary tick vector, *Amblyomma americanum*, is widespread throughout the southeastern USA, and *T. cervi* probably occurs throughout the range of the vector tick (Kocan and Kocan, 1991).

Identification of the benign *Theileria* spp. of both deer and cattle in the USA has been primarily based on the mammalian host of origin. Not only are *T. cervi* and the bovine *Theileria* sp. morphologically indistinguishable, but serologic cross-reactions occur and infections with either parasite result in mild or non-existent clin-

TABLE 1. Summary of *Theileria cervi* isolates including year of acquisition, mammalian host source, geographic origin, health status of host regarding clinical signs attributable to *Theileria cervi* infection, Giemsa stained blood film observations, and SSU rRNA gene types identified in each.

Isolate	Year	Source	Clinical signs	Parasite identified by	
				Giemsa stain	SSU rRNA type
CNELK	1994	Free-ranging elk, Canada	No	<i>Theileria</i> sp.	<i>Theileria cervi</i> F, G, G1
OKELK1	1998	Free-ranging elk, Oklahoma	No	<i>Theileria</i> sp.	<i>T. cervi</i> F, G1
USWTD1	1994	Farmed WTD ^a , Texas	Yes	<i>T. cervi</i>	<i>T. cervi</i> F, G, G1
USWTD2	1996	Farmed WTD, Texas	Yes	<i>T. cervi</i>	<i>T. cervi</i> F
USWTD3	1998	Wild-caught WTD, Oklahoma	No	<i>T. cervi</i>	<i>T. cervi</i> F, G2, G3

^a WTD = white-tailed deer.

ical signs in an otherwise healthy host population. Attempts at tick or blood transfer of *T. cervi* from white-tailed deer to domestic ruminants have not been successful (reviewed by Kingston, 1981). To date, similar transfer experiments of the bovine *Theileria* spp. into cervine hosts have not been reported.

Previously, SSU rRNA gene V4 variable region sequence analyses distinguished genes amplified from white-tailed deer and elk *T. cervi* isolates (sequence Types F and G) from genes of bovine *Theileria* spp. isolates (Chae et al., 1998). DNA probes to identify *Theileria* spp. have been designed based on specific SSU rRNA gene sequences (Allsopp et al., 1993). In the present study, entire Type F and G (and variations of G) SSU rRNA genes from white-tailed deer and elk *Theileria* sp. isolates were sequenced to confirm genes or gene regions specific to *T. cervi*.

MATERIALS AND METHODS

A male white-tailed deer housed at Oklahoma State University (Stillwater, Oklahoma, USA) was found by routine examination of a Giemsa-stained blood smear to be infected with *T. cervi* (designated USWTD3). The animal originated from Payne County (35°50'N, 97°00'W), Oklahoma, and was < 1-yr-old when the blood sample was drawn. During a routine epidemiological survey, a 2-yr-old bull elk (*Cervus elaphus canadensis*) residing at the Spavinaw Game Management Area (Spavinaw, Oklahoma; 36°20'N, 95°60'W) was found infected with *Theileria* sp. organisms (OKELK1). Two additional *T. cervi* isolates (USWTD1 and USWTD2) from Texas white-tailed deer and a Canadian elk *Theileria* sp. isolate (CNELK)

used in this study have been previously described (Chae et al., 1998). A summary of the *Theileria* spp. isolates used in this study is given in Table 1.

DNA extraction and SSU rRNA gene amplification procedures for the USWTD1, USWTD2 and CNELK isolates have been previously reported (Allsopp et al., 1989; Chae et al., 1998). DNA purification and SSU rRNA gene amplification from the USWTD3 and the OKELK1 *Theileria* isolates were similarly executed. Appropriately sized amplicons were confirmed by electrophoresis in ethidium bromide stained 1% agarose gels which were then viewed by UV transillumination. The amplification product was directly ligated into the plasmid vector pCR® 2.1-TOPO and INVαF' One Shot® *Escherichia coli* transformed according to manufacturer's recommendations (TOPO TA Cloning Kit®; Invitrogen, San Diego, California, USA). Transformed clones were color-selected and tested by colony amplification to ensure that the correct-sized insert was present. A portion of an isolated colony for each clone to be tested was mixed into 4 µl of molecular reagent grade water (Sigma®, St. Louis, Missouri, USA) and 2 µl then added to amplification reaction mix to a final volume of 10 µl. Each 10 µl amplification reaction contained 1 pmol each of T7 and M13 reverse primer, 5 mM KCl, 1 mM Tris-HCl, pH 8.3, 1.5 mM MgCl₂ and 0.2 U TAQ polymerase. The amplification conditions initiated with denaturation at 96 C for 5 min, followed by 30 cycles of 92 C for 30 sec, 55 C for 30 sec and 72 C for 30 sec and terminated with a final extension at 72 C for 10 min and hold at 4 C until use. The insert amplicons were confirmed by electrophoresis in ethidium bromide stained 1% agarose gels viewed by UV transillumination.

Plasmid DNA for each selected clone was purified from overnight broth cultures by a modified alkaline lysis procedure (QIAprep Spin Miniprep Kit; Qiagen®, Valencia, Califor-

nia, USA). The plasmid DNA samples were then used to identify both Type F and Type G clones for further study. Sequencing reactions for each plasmid DNA sample (Dye Terminator Cycle Sequencing Ready Reaction; PE Applied Biosystem, Norwalk, Connecticut, USA) used primer 528F to determine the SSU rRNA gene V4 variable region nucleotide sequence as previously described (Chae et al., 1998). Sequencing was carried out in either an ABI PRISM Model 373A or ABA Model 377 automated sequencer with Version 1.2.2 or Version 2.1.1 software, respectively (Gene Technologies Laboratory, Institute of Developmental and Molecular Biology, Department of Biology, Texas A&M University, College Station, Texas, USA).

Plasmid clones containing the entire SSU rRNA genes from USWTD1, USWTD2, and CNELK from previous work (Chae et al., 1998) were used for complete gene sequencing for these isolates. Since previously only one plasmid clone containing the WTD2 SSU rRNA gene insert was obtained (Chae et al., 1998), transformation of the original ligated plasmid into INVαF⁺ One Shot[®] *E. coli* was repeated. Putative transformed colonies were selected and tested by colony amplification as described above. Those with the correct-sized insert were then checked by sequence analysis of the SSU rRNA gene V4 variable region.

The complete forward and reverse strands of clones of SSU rRNA gene Type F in USWTD1, USWTD2, and CNELK were sequenced. The Type G genes from CNELK and USWTD1, Subtype G1 (USWTD1 and CNELK) and Subtype G2 genes (USWTD3) also were sequenced. A primer complementary to the T7 promotor region of the plasmid vector (Stratagene[®], La Jolla, California, USA) and a series of previously described internal primers (Elwood et al., 1989) were used for automated sequencing reactions and sequencing was carried out as described above. The CLUSTAL W (Version 1.60) multiple sequence alignment program (Thompson et al., 1994) and MACAW multiple alignment construction and analysis workbench (Version 2.05 Win 32i; Schuler et al., 1991) were used to facilitate sequence alignment and comparison. Sequence Types F and G and Subtype G1 also were aligned with corresponding *Theileria* spp. SSU rRNA gene sequences from the GenBank database including *T. buffeli* (GenBank Accession No. Z15106; Allsopp et al., 1994), *Theileria annulata* (GenBank Accession No. M64243, M34845; Gajadhar et al., 1991), *Theileria parva* 18S rRNA gene (GenBank Accession No. L02366; Allsopp et al., 1993), *T. parva* 16S rRNA gene (GenBank Accession No. L28999, Kibe et al., 1994), *Theileria taurotragi* (GenBank Acces-

sion No. L19082; Allsopp et al., 1994) and identity values were obtained by the ALIGN program (GeneStream, Centre de Recherche en Biochimie Macromoléculaire, Montpellier, France).

Nucleotide sequence data reported in this paper have been submitted to the GenBank[®] data base with the accession numbers U97054, U97055, U97056, AF86804, AF86805.

RESULTS

Amplification with SSU rRNA gene primers A and B resulted in a single band of approximately 1.8 kb for the *Theileria* spp. isolates USWTD3 and OKELK1 as observed by ethidium bromide stained agarose gel electrophoresis (not shown). *Theileria cervi* SSU rRNA gene sequence Types F (GenBank Accession No. U97054) and G Subtypes G1, G2 and G3 (GenBank Accession Nos. U97056, AF086804, and AF086805, respectively) were identified from these isolates (Table 1). Three additional plasmid clones from USWTD2 were identified with the correct-sized insert. These inserts were determined to be Type F by SSU rRNA gene V4 variable region sequence analysis.

Entire SSU rRNA gene sequences for Types F (USWTD1, USWTD2, and CNELK) and G (GenBank Accession No. U97055) (USWTD1 and CNELK) and Subtypes G1 (CNELK and USWTD1) and G2 (USWTD3) were obtained (Fig. 1). Partial gene sequences were obtained for Type F (USWTD3 and OKELK1) and Subtype G3 (USWTD3). USWTD3 and OKELK1 Type F sequences (3 clones checked for each) were identical to the V4 variable region sequences of USWTD1, USWTD2, and CNELK. Subtype G3 was found in three clones from the USWTD3 isolate, but as it differed from G2 only in one nucleotide position (635) in the V4 region (Fig. 1) the entire gene was not sequenced. The *T. cervi* Type F sequence, based on alignment of the V4 variable region, was found in all isolates tested (USWTD1, USWTD2, USWTD3, CNELK and OKELK1); *T. cervi* Type G was found in USWTD1 and CNELK (Table 1). Sub-

<i>T. parva</i>	1	15	16-180	181	195	196-210	211	225	226	240
<i>T. cervi</i>	AACCTGGTTGATCCT	ATTGCGCGCGTTTA	AACCGCTTG-CGTCG	TCCGGTGATTCATAA			
Type F	AACCTGGTTGATCCT	TTCGGGCGGCGTTTA	AACCGCTTG-CGTCG	TACGGTGATTCATAA			
Type G	AACCTGGTTGATCCT	TTCGGGCGGCGTTTA	AACCGCTTG-CGTCG	TACGGTGATTCATAA			
G1	AACCTGGTTGATCCT	GGTTGGCTGCGTTTA	ATTCTGAATGTCGAAA	ACCGGTGATTCATAA			
G2	AACCTGGTTGATCCT	TTTGGGCTGCGTTTA	ATTCTGAATGTCGAAAC	AC -GGTGATTCATAA			
G3			
<i>T. parva</i>	241	255	256	270	271-465	466	480	481-615	616	630
<i>T. cervi</i>	TAAATATGCGAATCG	T- ACTTAG	-- TGCG	ACGGGGCTTAAAGTC	ATTTCTGCTGTCATCG			
Type F	TAAACTTGC GAATCG	CGGCTTCGG- CTGCG	ACGGGGCTTAAAGTC	ATTTCTGCTGCTCCG				
Type G	TAAACTTGC GAATCG	CGGCTTCGG- CTGCG	ACGGGGCTTAAAGTC	ATTTCTGCTGCTCCG				
G1	TAAACTTGC GAATCG	CGGCTTACGGGCTGCG	ACGGGGCTTAAAGTC	ATTTCTGCTGCTCCG				
G2	TAAACTTGC GAATCG	TGGCTTAGG-CTGCG	ACGGGGCTTAAAGTC	ATTTCTGCTGCTCCG				
G3	ATTTCTGCTGCTCCG				
<i>T. parva</i>	631	645	646	660	661	675	676	690	691	705
<i>T. cervi</i>	C -TTGT - GTCCCTTC	--GGGGTCTCTGCA-	TGTGGCTTATTTTCGG	ACGGAGTTC-GCTTT	GTCTGGATGTTTACT					
Type F	CATT - - - TTCCCTTT	GAGGGGTTTTTGCGC	TGTGGCTTATTTTCGG	ACTGTGTTATGCACT	GTCCGGATGTTTACT					
Type G	CACATCTTCCCGTT	ATGGAGGTTTTGCGC	TGTGGCTTATTTTCGG	ACTGTGTTATGCACT	GTCCGGATGTTTACT					
G1	CACATCTTCCCGTT	ATGGAGGTTTTGCGC	TGTGGCTTATTTTCGG	ACTGTGTTATGCACT	GTCCGGATGTTTACT					
G2	CACATCTTCCCGTT	ATGGAGGTTTTGCGC	TGTGGCTTATTTTCGG	ACTGTGTTATGCACT	GTCCGGATGTTTACT					
G3	CACATCTTCCCGTT	ATGGAGGTTTTGCGC	TGTGGCTTATTTTCGG	ACTGTGTTATGCACT	GTCCGGATGTTTACT					
<i>T. parva</i>	706-35	736	750	751-810	811	825	826	840	841-915	916
<i>T. cervi</i>	TTTGCTTGAATAGT	AATGGTTAATAGGAA	CAGTTGGGGGCATTC	TTCATTAATCAAGAA			
Type F	TTTGCTTGAATAGT	AATGGTTAATAGGAA	CAGTTGGGGGCATTC	TTCATTAATCAAGAA			
Type G	TTTGCTTGAATAGT	AATGGTTAATAGGAA	CAGTTGGGGGCATTC	TTCATTAATCAAGAA			
G1	TTTGCTTGAATAGT	AATGGTTAATAGGAA	CAGTTGGGGGCATTC	TTCATTAATCAAGAA			
G2	TTTGCTTGAATAGT	AATGGTTAATAGGAA	CAGTTGGGGGCATTC	TTCATTAATCAAGAA			
G3	TTTGCTTGAATAGT	AATGGTTAATAGGAA	CAGTTGGGGGCATTC	TTCATTAATCAAGAA			
<i>T. parva</i>	931-75	976	990	991-1035	1036	1050	1051-70	1171	1185	1186
<i>T. cervi</i>	ACCATAAACTATGCC	TTGAGAGAAATCAAAA	TCCAGACAAAGGAAG	GATTGACAGATTGAT			
Type F	ACCATAAACTATGCC	TTGAGAGAAATCAAAA	TCCAGACAAAGGAAG	GATTGACAGATTGAT			
Type G	ACCATAAACTATGCC	TTGAGAGAAATCAAAA	TCCAGACAAAGGAAG	GATTGACAGATTGAT			
G1	ACCATAAACTATGCC	TTGAGAGAAATCAAAA	TCCAGACAAAGGAAG	GATTGACAGATTGAT			
G2	ACCATAAACTATGCC	TTGAGAGAAATCAAAA	TCCAGACAAAGGAAG	GATTGACAGATTGAT			
G3	ACCATAAACTATGCC	TTGAGAGAAATCAAAA	TCCAGACAAAGGAAG	GATTGACAGATTGAT			
<i>T. parva</i>	1201	1215	1216-1305	1306	1320	1321	1335	1336	1350	1351-1365
<i>T. cervi</i>	AGCTCTTCTTCTGATT	GGGTACGGGAATAAG	CTCTCGCTGTCCCGT	CATCGCTTCTTAGAG				
Type F	AGCTCTTCTTCTGATT	GGGTACGGGAATAAG	CTCTCGCTGTCCCGT	GATCGCTTCTTAGAG				
Type G	CGCCCTTCTTCTGATT	GCTCACGGGAATAAG	TTAAGACCGTCCCT	GGATGCTTCTTAGAG				
G1	CGCCCTTCTTCTGATT	GCTCACGGGAATAAG	TTAAGACCGTCCCT	GGATGCTTCTTAGAG				
G2	CGCCCTTCTTCTGATT	GCTCACGGGAATAAG	TTAAGACCGTCCCT	GGATGCTTCTTAGAG				
G3	CGCCCTTCTTCTGATT	GCTCACGGGAATAAG	TTAAGACCGTCCCT	GGATGCTTCTTAGAG				
<i>T. parva</i>	1366	1380	1381-1470	1471	1485	1486-1500	1501	1515	1516	1530
<i>T. cervi</i>	AAATCGCAAGGAAGT	GAGGCCCGGGTAATC	GATGGGGATCGAATA	TTGCAATTATTAATC				
Type F	AAATCGCAAGGAAGT	GAGGCCCGGGTAATC	GATGGGGATCGAATA	TTGCAATTATTAATC				
Type G	AAATCGCAAGGAAGT	GAGGCCCGGGTAATC	GATGGGGATCGAATA	TTGCAATTATTAATC				
G1	AAATCGCAAGGAAGT	GAGGT -GGGTAATC	GATGGGGATCGAATA	TTGCAATTATTAATC				
G2	AAATCGCAAGGAAGT	GAGGT -GGGTAATC	GATGGGGATCGAATA	TTGCAATTATTAATC				
G3	AAATCGCAAGGAAGT	GAGGT -GGGTAATC	GATGGGGATCGAATA	TTGCAATTATTAATC				
<i>T. parva</i>	1531-1545	1546	1560	1561-1753						
<i>T. cervi</i>	CATCAGCTTGTGCAG	1742						
Type F	CACCAGCTTGTGCAG	1748						
Type G	CACCAGCTTGTGCAG	1750						
G1	CACCAGCTTGTGCAG	1750						
G2	CACCAGCTTGTGCAG	1748						
G3	CACCAGCTTGTGCAG	1748						

FIGURE 1. Nucleotide sequence of *Theileria cervi* SSU rRNA gene types and subtypes aligned with the *Theileria parva* SSU rRNA gene nucleotide sequence. Gaps represented by dashes (-) were introduced into the aligned sequences to maximize homology. Nucleotide positions differing from those of *T. parva* are designated by bold type. The V4 variable region is delineated by arrows (nucleotide positions 621–661). Gaps where the sequence is identical in all genes are designated by dotted lines (.....). The unsequenced portion of Subtype G3 is shown by a broken line (---). The SSU rRNA gene sequence shown for *T. parva* was obtained from the GenBank data base, Accession Number M67476. *Theileria cervi* SSU rRNA gene GenBank Accession Numbers are as follows: Type F, U97054; Type G, U97055; Subtype G1, U97056; Subtype G2, AF86804; Subtype G3, AF86805.

type G1 was found in both deer and elk isolates, USWTD1 and OKELK1. Subtypes G2 and G3 were identified in USWTD3.

The total SSU rRNA gene lengths were: Type F, 1748 bp; Type G, 1750 bp; Subtype G1, 1750 bp; Subtype G2, 1748. G1, G2 and G3 are considered subtypes of

TABLE 2. Number of nucleotide position differences among the *Theileria cervi* SSU rRNA gene types. Top matrix shows differences found in the entire gene sequence; bottom matrix shows differences found only in the V4 variable region of the gene (nucleotide positions 621–661).

Nucleotide differences among <i>Theileria cervi</i> SSU rRNA gene types					
Type	G	G1	G2	G3	F
G	0	23	19	ND ^a	34
G1	2	0	11	ND	45
G2	1	3	0	ND	54
G3	2	4	1	0	ND
F	10	10	11	11	0

^a ND = not done. Entire *Theileria cervi* G3 SSU rRNA gene not sequenced.

Type G because they differ from G in only one or two nucleotide positions through the V4 region (nucleotide positions 623–663) (Fig. 1 and Table 2) and share sequence identity in nucleotide positions 1171–1174, 1199–1204, 1307–1309, 1319–1328, and 1334–1339 (Fig. 1). In contrast, Type F does not share sequence identity in these positions (Fig. 1) and also differs from Type G in 10 nucleotide positions through the V4 region (Fig. 1 and Table 2).

Differences in the V4 region sequences among the subtypes range from one nucleotide substitution between G and G2, and G2 and G3 to four substitutions between G1 and G3 (Table 2). Subtypes G2 and G3 share substitutions distinguishing them from G1: G2 and G3 have guanine (G) instead of thymidine (T) at position 657; G2 and G3 have cytosine (C) instead of thymidine (T) at 926 (Fig. 1).

Differences found in the entire gene sequences among the various *T. cervi* SSU rRNA genes are summarized in Table 2. Total Type G and Subtype G1 genes showed variation in 23 positions; Type G and Subtype G2 showed variation in 19 positions. Variation was only found in 11 positions between Subtypes G1 and G2. In contrast, differences at 33 positions were noted between Types F and G. No micro-

heterogeneity was found among the Type F sequences.

Identity values determined among the *Theileria* spp. and Types F and G and Subtype G1 showed Type G and Subtype G1 to share the highest identity among the obtained sequences, 98.8% (Table 3). The percent identity between *T. cervi* Type F and Type G was 97.8. When compared to SSU rRNA gene sequences reported for other *Theileria* spp., identity values for Type F ranged from 96.7 to 97.5%, and for Type G from 95 to 95.6%. These percentages reflect 43–57 position differences in Type F and 76–96 position differences in Type G when compared to other *Theileria* spp. (Table 3). *Theileria cervi* SSU rRNA sequence Types F and G (and subtypes) were closest in sequence homology to that of *T. parva* (GenBank Accession Nos. L02366 and L28999).

DISCUSSION

SSU rRNA data indicate that both white-tailed deer and elk harbor *T. cervi*, as supported by this study and previous work (Chae et al., 1998). Complete nucleotide sequences from SSU rRNA genes amplified from cervine blood infected with *Theileria* spp. confirmed that two sequence Types, F and G, were present in the same isolate population. Microheterogeneity in Type G was observed in isolates from two white-tailed deer and two elk, and subtypes G1, G2, and G3 were designated. Considering that Type G SSU rRNA gene heterogeneity was the norm rather than the exception among the small number of isolates in this study, it is likely that additional divergent Type G sequences will be found among other *T. cervi* isolates. Thus, the Type G gene and G subtypes likely comprise a polymorphic family of G SSU rRNA genes.

SSU rRNA Type G or a G subtype gene was not found in one isolate, USWTD2. Amplification of the SSU rRNA gene from this isolate was very difficult and the original ligation and transformation yielded only one plasmid clone with the SSU

TABLE 3. ALIGN program sequence homology of *Theileria* spp. SSU rRNA genes. SSU rRNA gene sequences from the bovine *Theileria* spp., *Theileria annulata*, *Theileria parva*, and *Theileria taurotragi*, and from the cervine *Theileria* spp., *Theileria* sp. from sable antelope and *Theileria cervi* SSU rRNA gene Types F and G and Subtype G1. Upper matrix shows percent identity between sequences. Lower matrix shows the number of nucleotide differences in bold print and the number of overlapping nucleotides in parentheses.

SSU rRNA ^a	<i>T. annulata</i>	<i>T. parva</i>	<i>T. taurotragi</i>	<i>Theileria</i> sp.	<i>T. buffeli</i>	<i>T. cervi</i> -F	<i>T. cervi</i> -G	<i>T. cervi</i> -G1
<i>T. annulata</i>	100 (1,744)	98.4	97.8	96.1	96.3	96.8	95.2	94.7
<i>T. parva</i>	28 (1,744)	100 (1,742)	98.5	96.6	96.6	97.5	95.6	95.2
<i>T. taurotragi</i>	39 (1,744)	27 (1,742)	100 (1,737)	96.6	96.4	97.0	95.4	95.0
<i>Theileria</i> sp.	68 (1,746)	59 (1,746)	59 (1,746)	100 (1,746)	96.8	96.7	95.2	94.8
<i>T. buffeli</i>	65 (1,744)	59 (1,742)	62 (1,740)	55 (1,746)	100 (1,740)	96.9	95.0	94.7
<i>T. cervi</i> -F	56 (1,748)	43 (1,748)	52 (1,748)	57 (1,748)	54 (1,748)	100 (1,748)	97.8	96.8
<i>T. cervi</i> -G	84 (1,750)	76 (1,750)	80 (1,750)	84 (1,750)	87 (1,750)	38 (1,750)	100 (1,750)	98.6
<i>T. cervi</i> -G1	92 (1,750)	83 (1,750)	88 (1,750)	91 (1,750)	93 (1,750)	56 (1,750)	24 (1,750)	100 (1,750)

^a GenBank Accession Numbers for SSU rRNA gene sequences: *Theileria annulata*, M64243 and M34845; *Theileria parva*, L02366 and L28999; *Theileria taurotragi*, L19082; *Theileria* sp., L19082; *Theileria buffeli*, Z15106; *Theileria cervi* Type F, U97054; *Theileria cervi* Type G, U97055; *Theileria cervi* Subtype G1, U97056.

rRNA gene insert (Chae et al., 1998). In this study, bacterial transformation by the original ligated plasmid preparation was repeated in an effort to obtain additional plasmid clones to sequence. Only three more clones containing the gene were obtained and all three contained inserts of the Type F sequence. Unfortunately, additional DNA was not available to repeat the entire protocol. Clearly, there is a possibility that this isolate possessed other SSU rRNA gene types that we did not find.

The existence of multiple *T. cervi* SSU rRNA gene sequence types may represent genes from mixed populations of parasites or multiple copy units within a parasite as shown in *T. parva* (Kibe et al., 1994). Inasmuch as both Types F and G (or G subtypes) were found in all isolates, with the exception of USWTD2, it appears likely that individual *T. cervi* parasites possess both gene types. It is not unlikely that different SSU rRNA genes may be expressed during different developmental stages of the *Theileria* life cycle. Transcript regula-

tion of distinct stage specific SSU rRNAs has been shown in *Plasmodium* spp. (Li et al., 1997; Waters et al., 1989; Corredor and Enea, 1994). Further studies are needed to determine if both gene types are represented in multiple gene clusters in *T. cervi*, whether both types produce functional rRNAs, and whether transcription of different SSU rRNA gene types is stage dependent in *T. cervi*. It is also possible that the microheterogeneity observed in the Type G sequence reflects the presence of pseudogenes.

SSU rRNA Type G microheterogeneity previously reported based on V4 region data (Chae et al., 1998) was confirmed by the present study. Subtype G1 was found in both white-tailed deer and elk isolates, USWTD1, CNELK, and OKELK1. Subtypes G2 and G3, which are nearly identical through the V4 region, were found in a white-tailed deer isolate, USWTD3. Alternatively, no microheterogeneity was found among the Type F SSU rRNA gene sequences regardless of host or geographic origin. Thus, the Type F gene appears

highly conserved among the *T. cervi*. Although a single base substitution in the V4 region has been reported (Chae et al., 1998), the Type F genes identified in this study did not show this substitution.

The conservation of the Type F SSU rRNA gene sequence among *T. cervi* isolates provides a base of reference when investigating new isolates and presents an opportunity to develop specific DNA primer sequences which could be used in amplification-based assays for detecting *T. cervi* infections in deer or vector ticks. Such assays could be used to screen ticks for the presence of the parasite to identify potential vectors aside from the known vector, *A. americanum*, although ultimately transmission experiments would be necessary to confirm the ability of a tick species to vector the parasite. Sensitive molecular methods could facilitate epidemiological studies in ticks, deer, and other suspected hosts of the parasite.

ACKNOWLEDGMENTS

The authors thank D. Cruz for excellent technical assistance. This research was supported in part by the Texas Agricultural Experiment Station (Project H-6261), Texas A&M University Faculty Minigrant (FMG 95-120), and the Korean Research Foundation (KRF 01-G-0162).

LITERATURE CITED

- ALLSOPP, B. A., M. CARRINGTON, H. BAYLIS, S. SOHAL, T. DOLAN, AND K. IAMS. 1989. Improved characterization of *Theileria parva* isolates using the polymerase chain reaction and oligonucleotide probes. *Molecular and Biochemical Parasitology* 35: 137–148.
- , H. A. BAYLIS, M. T. E. P. ALLSOPP, T. CAVALIER-SMITH, R. P. BISHOP, D. M. CARRINGTON, B. SOHANPAL, AND P. SPOONER. 1993. Discrimination between six species of *Theileria* using oligonucleotide probes which detect small subunit ribosomal RNA sequences. *Parasitology* 107: 157–165.
- ALLSOPP, M. T. E. P., T. CAVALIER-SMITH, D. T. DE WAAL, AND B. A. ALLSOPP. 1994. Phylogeny and evolution of the piroplasms. *Parasitology* 108: 147–152.
- CHAE, J. S., J. M. LEE, O. D. KWON, P. J. HOLMAN, S. D. WAGHELA, AND G. G. WAGNER. 1998. Nucleotide sequence heterogeneity in the small subunit ribosomal RNA gene variable (V4) region among and within geographic isolates of *Theileria* from cattle, elk and white-tailed deer. *Veterinary Parasitology* 75: 41–52.
- CORREDOR, V., AND V. ENEA. 1994. The small ribosomal subunit RNA isoforms in *Plasmodium cynomolgi*. *Genetics* 136: 857–865.
- ELWOOD, H. J., G. J. OLSEN, AND M. L. SOGIN. 1989. The small subunit ribosomal RNA gene sequences from the hypotrichous ciliates *Oxytricha nova* and *Stylonychia pustulata*. *Journal of Molecular Biology and Evolution* 5: 399–410.
- GAJADHAR, A. A., W. C. MARQUARDT, R. HALL, J. GUNDERSON, E. V. ARIZTIA-CARMONA, AND M. L. SOGIN. 1991. Ribosomal RNA sequences of *Sarcocystis muris*, *Theileria annulata* and *Cryptosporidium parvum* reveal evolutionary relationships among Apicomplexans, dinoflagellates and ciliates. *Molecular and Biochemical Parasitology* 45: 147–154.
- KIBE, M. K., O. K. OLE-MOIYOI, V. NENE, B. KHAN, B. A. ALLSOPP, N. E. COLLINS, S. P. MORZARIA, E. I. GOBRIGHT, AND R. P. BISHOP. 1994. Evidence for two single copy units in *Theileria parva* ribosomal RNA genes. *Molecular and Biochemical Parasitology* 66: 249–259.
- KINGSTON, N. 1981. Protozoan parasites. In *Diseases and parasites of white-tailed deer*, W. R. Davidson, F. A. Hayes, V. F. Nettles, and F. E. Kellogg (eds.). Tall Timbers Research Station Miscellaneous Publication No. 7, Tallahassee, Florida, pp. 193–236.
- KOCAN, A. A., AND K. M. KOCAN. 1991. Tick-transmitted protozoan diseases of wildlife in North America. *Bulletin of the Society of Vector Ecology* 16: 94–108.
- LAIRD, J. S., A. A. KOCAN, K. M. KOCAN, S. M. PRESLEY, AND J. A. HAIR. 1988. Susceptibility of *Amblyomma americanum* to natural and experimental infections with *Theileria cervi*. *Journal of Wildlife Diseases* 24: 679–683.
- LI, J., R. R. GUTELL, S. H. DAMBERGER, R. A. WIRTZ, J. C. KISSINGER, M. J. ROGERS, J. SATTABONGKOT, AND T. F. MCCUTCHAN. 1997. Regulation and trafficking of three distinct 18S ribosomal RNAs during development of the malaria parasite. *Journal of Molecular Biology* 269: 203–213.
- SCHAEFFLER, W. F. 1961. *Theileria* in white-tailed deer in the United States. *Journal of Protozoology* 8 (supplement): 10.
- . 1962. *Theileria cervi* infection in white-tailed deer (*Damas virginianus*) in the United States. Ph.D. Dissertation, University of Illinois, Urbana, Illinois, 104 pp.
- SCHULER, G. D., S. F. ALTCHUL, AND D. J. LIPMAN. 1991. A workbench for multiple alignment construction and analysis. *Proteins: Structure, Function, and Genetics* 9: 180–190.
- THOMPSON, J. D., D. G. HIGGINS, AND T. J. GIBSON.

1994. CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22: 4673–4680.
- WALDRUP, K. A., E. COLLISSE, S. E. BENTSEN, C. K. WINKLER, AND G. G. WAGNER. 1989. Prevalence of erythrocytic protozoa and serologic reactivity to selected pathogens in deer in Texas. *Preventive Veterinary Medicine* 7: 49–58.
- WATERS, A. P., C. SYIN, AND T. F. MCCUTCHAN. 1989. Developmental regulation of stage-specific ribosome populations in *Plasmodium*. *Nature* 342: 438–440.

Received for publication 31 August 1998.