

Survival of *Mycobacterium bovis* on Feedstuffs Commonly Used as Supplemental Feed for White-tailed Deer (*Odocoileus virginianus*)

Authors: Palmer, Mitchell V., and Whipple, Diana L.

Source: Journal of Wildlife Diseases, 42(4) : 853-858

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-42.4.853>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Survival of *Mycobacterium bovis* on Feedstuffs Commonly Used as Supplemental Feed for White-tailed Deer (*Odocoileus virginianus*)

Mitchell V. Palmer,^{1,2} and Diana L. Whipple¹ ¹ United States Department of Agriculture, Agricultural Research Service, National Animal Disease Center, Ames, Iowa, USA; ² Corresponding author (email: mpalmer@nadc.ars.usda.gov)

ABSTRACT: *Mycobacterium bovis*, the causative agent of bovine tuberculosis, has become established in free-ranging white-tailed deer *Odocoileus virginianus* in northeastern Michigan. The practice of supplemental feeding of white-tailed deer during the winter is believed to contribute to transmission of *M. bovis* between deer. The current study was conducted to determine the ability of *M. bovis* to survive on various feedstuffs commonly used as supplemental feed for deer in northeast Michigan (i.e., apples, corn, carrots, sugar beets, potatoes, and hay) and the effect of maintenance at -20°C , 8°C , and 23°C on survival. *Mycobacterium bovis* survived on all feedstuffs at all temperatures tested for at least 7 days. At 23°C , *M. bovis* could still be isolated from samples of apples, corn and potatoes at 112 days. This study suggests that contamination of feedstuffs by *M. bovis*-infected deer could act as a source of indirect transmission between deer because *M. bovis* is able to survive in temperatures similar to those recorded during winter months in northeastern Michigan. Current efforts to ban or control supplemental feeding of deer should have a positive effect on decreasing transmission of *M. bovis* among deer.

Key words: Feedstuffs, *Mycobacterium bovis*, *Odocoileus virginianus*, survivability, tuberculosis, white-tailed deer.

In 1994, a free-ranging white-tailed deer (*Odocoileus virginianus*) in Michigan was diagnosed with tuberculosis due to *Mycobacterium bovis*, the causative agent of bovine tuberculosis (Schmitt et al., 1997). Subsequent surveys identified a focus of *M. bovis* infection in free-ranging white-tailed deer in northeast Michigan (O'Brien et al., 2001, 2002). This represents the first known reservoir of *M. bovis* in free-ranging wildlife in the United States and a significant obstacle to the eradication of bovine tuberculosis in domestic livestock. Several factors are thought to have contributed to the estab-

lishment and persistence of *M. bovis* in this wildlife reservoir. It is postulated that *M. bovis* was transmitted from cattle to deer at some time during the early to mid 1900s when a large number of Michigan cattle were infected with *M. bovis* (Frye, 1995). During this same period, Michigan's deer population was steadily increasing beyond normal habitat carrying capacity to focal concentrations of 19 to 23 deer per km^2 (Schmitt et al., 1997). Transmission and maintenance of *M. bovis* among deer is thought to have been facilitated by the common practice in Michigan of long-term winter feeding of large volumes of sugar beets, carrots, corn, apples, pumpkins, and pelleted feed to deer by private citizens to prevent migration and decrease winter mortality in order to keep deer numbers high for hunting purposes (Schmitt et al., 1997). DNA analysis of *M. bovis* isolates from Michigan white-tailed deer showed that the majority of deer were infected with a common strain of *M. bovis* suggesting a single source of infection (Whipple et al., 1997).

Experimentally, transmission between deer and between deer and cattle has been demonstrated through the sharing of feed with no opportunity for direct contact (Palmer et al., 2004b). Research under a variety of conditions has demonstrated that *M. bovis* can persist for prolonged periods in the environment (Williams and Hoy, 1930; Maddock, 1933; Duffield and Young, 1985; Jackson et al., 1995; Tanner and Michel, 1999). *Mycobacterium bovis* has been shown to persist in cool, dark environments for 4–6 wk (Duffield and Young, 1985; Tanner and Michel, 1999), on growing grass for as long as 7 wk, and

in liquid manure for at least 16 wk (Williams and Hoy, 1930; Maddock, 1933). In contrast to such prolonged survival, other research has shown survival times of 7–28 days inside dens of brushtail possums in New Zealand (Jackson et al., 1995). Little information exists as to the ability of a particular strain, such as that isolated from white-tailed deer in northeast Michigan, to exist on specific substrates that are commonly used as feedstuffs. This study was conducted to determine the ability of *M. bovis* to survive on various feedstuffs commonly used as supplemental feed for deer in northeast Michigan and the effect of storage temperature on survival.

Mycobacterium bovis strain 1315 was used in the present study. Strain 1315 was originally isolated from a white-tailed deer in northeast Michigan and has the same RFLP pattern as the majority of isolates obtained from deer in Michigan. Bacterial suspensions consisted of mid-log-phase *M. bovis* grown in Middlebrook 7H9 liquid media supplemented with 10% oleic acid-albumin-dextrose complex (OADC) (Becton Dickinson Co., Sparks, Maryland) plus 0.05% Tween 80 (Sigma Chemical Co., St. Louis, Missouri) grown for 10 days at 37 C. To harvest bacilli from the culture media, cells were pelleted by centrifugation at $750 \times G$ and the pellet resuspended in phosphate-buffered saline solution (PBS, 0.01 M, pH 7.2) to the appropriate concentration. Bacterial suspensions were used immediately without freezing.

Samples of sugar beets, apples, carrots, and potatoes were diced into cubes of approximately $1\text{--}2\text{ cm}^3$. Samples of alfalfa hay were chopped to approximately 3–4 cm lengths. Shelled corn was divided into 10 g samples. Ten grams each of sugar beets, apples, carrots, potatoes, and shelled corn were placed into 50 ml centrifuge tubes. Ten grams of alfalfa hay was similarly placed in suitable plastic bags. To each tube or bag a 1 ml volume of a bacterial suspension containing 1.1×10^6 colony forming units (CFU) of *M.*

bovis was added. Tubes containing samples and bacterial suspension were rotated thoroughly to ensure mixing of the sample and suspension. Six different types of feedstuffs were examined in duplicate at 2 hr, and 1, 2, 3, 7, 14, 21, 28, 56, 84, and 112 days after inoculation. Each evaluation was done at three different environmental temperatures, -20 C (freezing), 8 C (refrigeration), and 23 C (room temperature), totaling 396 samples.

For isolation of *M. bovis* from carrots, apples, potatoes, and sugar beets, samples were homogenized with 50 ml of nutrient broth containing phenol red. Hay and corn samples were processed by mixing in a hand grinder with 40 ml and 60 ml, respectively, of phenol red nutrient broth. Tubes containing samples were thoroughly rinsed with nutrient broth as the sample was extracted for homogenization. Samples were homogenized for 30 sec and allowed to settle for 15 min. Twenty milliliters of suspension was removed, placed in a new 50-ml centrifuge tube, and incubated overnight (16–18 hr) at 37 C. After incubation, an equal volume of 1.0 N NaOH was added for decontamination. After 10 min, the sample was neutralized with 10 N HCl to pH 7 and centrifuged at $2,000 \times G$ for 20 min. The supernatant was decanted and the sediment suspended in 2 ml of phosphate buffered saline (PBS) containing amphotericin B (50 $\mu\text{g/ml}$). For each sample, two plates containing Middlebrook 7H11 media were inoculated with 0.2 ml of suspension. Plates were incubated at 37 C and growth evaluated at 7 days, 4 wk, and 8 wk. Two hundred microliters of each sample were also inoculated into vials of commercial liquid media (BACTEC 12B medium, Becton Dickinson) containing selected antimicrobials for isolation of mycobacteria (BACTEC PANTA Plus, Becton Dickinson). Growth on Middlebrook 7H11 agar plates was semiquantitatively scored using the following system; 0 (no growth), 1 (1–10 CFU), 2 (11–50 CFU), 3 (51–100 CFU), 4 (>100 CFU).

Mycobacterium bovis survived on all feedstuffs at all storage temperatures tested for at least 7 days (Table 1). At the termination of the study (112 days), *M. bovis* could be recovered from all feedstuffs at -20°C , from samples of apples, corn, hay, and potatoes at 8°C , and from samples of apples, corn and potatoes at 23°C . Survivability was least favorable on carrots and sugar beets; however, *M. bovis* could still be recovered from both vegetables at 112 days at -20°C and 14 and 84 days, respectively, for carrots and sugar beets at 8°C .

The prolonged survivability of *M. bovis* on feedstuffs commonly used for supplemental feeding of deer would increase the likelihood of transmission of *M. bovis* in areas where *M. bovis* is endemic in white-tailed deer, such as northeastern Michigan. In the current study, survivability of *M. bovis* was facilitated by decreased temperature, with recovery of viable *M. bovis* from all feedstuffs maintained at -20°C for 112 days, and recovery of *M. bovis* from apples, corn, hay, and potatoes after maintenance at 8°C for 112 days.

According to historical climate data (<http://www.crh.noaa.gov>), the average temperature in northeastern Michigan during the period 1971–2000, was 1.4°C , -4.0°C , -7.8°C , and -7.2°C for November, December, January and February, respectively. On average, temperatures during the winter in this region do not exceed 0°C for 55 out of 84 days. Environmental temperatures such as those recorded in winter in northeastern Michigan would increase the likelihood of *M. bovis* survival in the environment.

Supplemental feeding of deer in Michigan consists of placing large piles of feedstuffs, including those examined in the present study, and allowing wildlife free access over the entire winter (Schmitt et al., 1997). The volume of feed at some feeding sites can approach hundreds of thousands of kilograms per winter season (Miller et al., 2003). The results of the present study suggest that *M. bovis* can

remain viable on some feedstuff piles for most of the winter with possible exposure of large numbers of deer to viable *M. bovis*. Indeed, analysis has shown that supplemental feeding of deer is associated with the risk of tuberculosis in white-tailed deer (Miller et al., 2003). Specific risk factors include the presence of hardwood forests, other feeding sites in the area, the number of deer fed per year, and the quantity of fruits or vegetables provided (Miller et al., 2003). Interestingly, a commonly observed habit of white-tailed deer may also contribute to disease transmission. During winter, the outer surface of feed piles becomes frozen and deer break small holes, large enough to accommodate the muzzle, through the frozen surface to access nonfrozen feed (Miller et al., 2003). Later, other deer use the same holes to access feed. This habit may concentrate feed contamination to critical areas shared by numerous deer.

Previous studies have examined the ability of *M. bovis* to survive under various environmental conditions (Williams and Hoy, 1930; Maddock, 1933; Duffield and Young, 1985; Jackson et al., 1995; Tanner and Michel, 1999). Although results have varied, ranging from 4 days on pasture (Griffin et al., 2005) to 6 mo in dung (Maddock, 1933), reports have consistently demonstrated that survivability is enhanced in cool, moist conditions, especially when samples are protected from direct sunlight. Moreover, previous studies have taken place under environments commonly encountered in locales such as South Africa (Tanner and Michel, 1999), Australia (Duffield and Young, 1985), New Zealand (Jackson et al., 1995), and the United Kingdom (Williams and Hoy, 1930; Maddock, 1933) and have not considered environments that approach the cold climates of northern Michigan.

Mycobacterium bovis, like other mycobacteria has a unique cell wall, of which lipids account for approximately 60%. This extremely lipophilic cell wall allows *M. bovis* to resist adverse environmental

TABLE 1. Survivability of *M. bovis* on various feedstuffs used for supplemental feeding of white-tailed deer in Michigan at storage temperatures of 23 C, 8 C, and -20 C.

Feedstuff	Temp	Days of storage ^a										
		0 ^b	1	2	3	7	14.0	21	28	56	84.0	112
Apples	23	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	3±0.58	4.0±0.0	ND ^c	1.5±0.50	1±0.0	0	0.50±0.29
	8	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	1.5±0.29	3.33±0.33	3.50±0.50	4.0±0.0	4.0±0.0	3.0±0.58	2.0±0.58
Carrots	-20	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0
	23	4.0±0.0	4.0±0.0	2.0±0.58	1.0±0.41	0.50±0.29	0	ND	0	0	0	0
Corn	8	4.0±0.0	4.0±0.0	4.0±0.0	1.0±0.0	2.0±0.41	0.50±0.29	0	0	0	0	0
	-20	4.0±0.0	2.0±0.0	4.0±0.0	3.0±0.0	1.75±0.25	3.25±0.25	0.75±0.41	0.75±0.41	0.75±0.25	1.0±0.58	0.25±0.25
Hay	23	4.0±0.0	4.0±0.0	4.0±0.0	3.75±0.25	4.0±0.0	2.25±0.25	0.67±0.33	1.25±0.25	0.33±0.33	0	0.25±0.25
	8	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	2.75±0.41	4.0±0.0
Potatoes	-20	4.0±0.0	4.0±0.0	4.0±0.0	ND	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0
	23	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	ND	ND	ND	ND	ND	ND
Sugar Beets	8	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	3.5±0.5	4.0±0.0	4.0±0.0	3.5±0.5	4.0±0.0
	-20	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0
Sugar Beets	23	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	ND	2.75±0.63	4.0±0.0	4.0±0.0	4.0±0.0
	8	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	2.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0
Sugar Beets	-20	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0
	23	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	2.25±0.25	0	ND	2.25±0.25	ND	0	0
Sugar Beets	8	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	1.0±0.0	1.7±0.33	0.5±0.29	0.5±0.29	0
	-20	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0	4.0±0.0

^a Days represent mean of duplicate samples (±S.E.) and days after addition of *M. bovis* to feed samples that isolation of *M. bovis* was possible.
^b 0 = no growth, 1 = 1-10 CFU, 2 = 11-50 CFU, 3 = 51-100 CFU, 4 = >100 CFU.
^c Not done.

conditions such as drying (Willet, 1992). Cultures maintained at 37 C have been found viable after 12 yr of storage (Willet, 1992). The environment in which the bacilli are found is also critical to prolonged survival in the environment. Bacilli in sputum require 20 hr to 30 hr of exposure to direct sunlight to be killed, whereas bacilli from culture can be killed in 2 hr (Willet, 1992). Previous studies have demonstrated that *M. bovis*-infected white-tailed deer shed the organisms in nasal secretions and saliva (Palmer et al., 2004a, b). It is likely the environment found within these bodily fluids enhances survivability.

Current regulations in Michigan ban supplemental feeding of deer within the TB endemic area (<http://www.michigandnr.com/>) and restrict supplemental feeding in areas where TB has not been identified in deer. However, not all supplemental feeding of wildlife is intentional. Current agricultural practices, such as feeding cattle using hay in large round bales on pastures, do not preclude deer from sharing feed with other deer or livestock in regions where deer have access to livestock pastures.

Limitations of the current study include the lack of ability to evaluate the effect of factors such as moisture (humidity and precipitation); changes in temperature; freeze-thaw cycles; activity of other bacteria, protozoa, or fungi; and ultraviolet light on the survivability of *M. bovis*. It is likely that factors such as these would have a detrimental effect on *M. bovis* survivability. Although not conducted in regions with environmental conditions similar to northern Michigan, other studies conducted in natural settings where factors such as humidity, precipitation, and ultraviolet light were monitored have demonstrated prolonged survivability of *M. bovis* well within the span of time necessary to transmit *M. bovis* indirectly among deer through sharing of feed (Jackson et al., 1995; Tanner and Michel, 1999).

The lack of precise quantitative culture

results in the previous study also precludes conclusions concerning the likelihood of the presence of sufficient numbers of organisms at various time-points and their ability to infect white-tailed deer successfully. The dose of *M. bovis* that deer receive during natural infections is unknown and likely highly variable; however, previous experimental studies have demonstrated that low doses of *M. bovis* can produce tuberculous lesions in white-tailed deer (Palmer et al., 2002). Inoculation of 32 white-tailed deer with 300 CFU of the strain of *M. bovis* used in present study resulted in bacterial colonization and lesion formation in all deer (Palmer et al., 2002). Moreover, lesions have been successfully produced in white-tailed deer inoculated with as few as 40 CFU of *M. bovis* (M. Palmer, unpubl. obs.), in red deer with as few as 200 CFU of *M. bovis* (Griffin et al., 2006) and in cattle with a single CFU of *M. bovis* (Dean et al., 2005). This study suggests that contamination of feedstuffs by *M. bovis* infected deer could act as a source of indirect transmission to other deer because *M. bovis* is able to survive in temperatures similar to those recorded during the winter months in northeastern Michigan. Current efforts to ban or control supplemental feeding of deer should have a positive effect on decreasing transmission of *M. bovis* between deer.

LITERATURE CITED

- DEAN, G. S., S. G. RHODES, M. COAD, A. O. WHELAN, P. J. COCKLE, D. J. CLIFFORD, R. G. HEWINSON, AND H. M. VORDERMEIR. 2005. Minimum infective dose of *Mycobacterium bovis* in cattle. *Infection and Immunity* 73: 6467–6471.
- DUFFIELD, B. J., AND D. A. YOUNG. 1985. Survival of *Mycobacterium bovis* in defined environmental conditions. *Veterinary Microbiology* 10: 193–197.
- FRYE, G. H. 1995. Bovine tuberculosis eradication: The program in the United States. In *Mycobacterium bovis* infection in animals and humans, C. O. Thoen and J. H. Steele (eds.). Iowa State University Press, Ames, Iowa, pp. 119–129.
- GRIFFIN, J. F. T., C. G. MACKINTOSH, AND C. R. RODGERS. 2006. Factors influencing the protective efficacy of a BCG homologous prime-

- boost vaccination regime against tuberculosis. *Vaccine* 24: 835–845.
- JACKSON, R., G. W. DELISLE, AND R. S. MORRIS. 1995. A study of the environmental survival of *Mycobacterium bovis* on a farm in New Zealand. *New Zealand Veterinary Journal* 43: 346–352.
- MADDOCK, E. C. G. 1933. Studies on the survival time of the bovine tubercle bacillus in soil and dung, in dung and on grass, with experiments on the preliminary treatment of infected organic matter and the cultivation of the organism. *Journal of Hygiene* 33: 103–117.
- MILLER, R., J. B. KANEENE, S. D. FITZGERALD, AND S. M. SCHMITT. 2003. Evaluation of the influence of supplemental feeding of white-tailed deer (*Odocoileus virginianus*) on the prevalence of bovine tuberculosis in the Michigan wild deer population. *Journal of Wildlife Diseases* 39: 84–95.
- O'BRIEN, D. J., S. D. FITZGERALD, T. J. LYON, K. L. BUTLER, J. S. FIERKE, K. R. CLARKE, S. M. SCHMITT, T. M. COOLEY, AND D. E. BERRY. 2001. Tuberculous lesions in free-ranging white-tailed deer in Michigan. *Journal of Wildlife Diseases* 37: 608–613.
- , S. M. SCHMITT, J. S. FIERKE, S. A. HOGLE, S. R. WINTERSTEIN, T. M. COOLEY, W. E. MORITZ, K. L. DIEGEL, S. D. FITZGERALD, D. E. BERRY, AND J. B. KANEENE. 2002. Epidemiology of *Mycobacterium bovis* in free-ranging white-tailed deer, Michigan, USA, 1995–2000. *Preventive Veterinary Medicine* 54: 47–63.
- PALMER, M. V., W. R. WATERS, AND D. L. WHIPPLE. 2002. Lesion development in white-tailed deer (*Odocoileus virginianus*) experimentally infected with *Mycobacterium bovis*. *Veterinary Pathology* 39: 334–340.
- , ———, AND ———. 2004a. Investigation of the transmission of *Mycobacterium bovis* from deer to cattle through indirect contact. *American Journal of Veterinary Research* 11: 1483–1489.
- , ———, AND ———. 2004b. Shared feed as a means of deer to deer transmission of *Mycobacterium bovis*. *Journal of Wildlife Diseases* 40: 87–91.
- SCHMITT, S. M., S. D. FITZGERALD, T. M. COOLEY, C. S. BRUNING-FANN, L. SULLIVAN, D. BERRY, T. CARLSON, R. B. MINNIS, J. B. PAYEUR, AND J. SIKARSKIE. 1997. Bovine tuberculosis in free-ranging white-tailed deer from Michigan. *Journal of Wildlife Diseases* 33: 749–758.
- TANNER, M., AND A. L. MICHEL. 1999. Investigation of the viability of *M. bovis* under different environmental conditions in the Kruger National Park. *Onderstepoort Journal of Veterinary Research* 66: 185–190.
- WHIPPLE, D. L., R. M. MEYER, D. F. BERRY, J. L. JARNIGAN, AND J. B. PAYEUR. 1997. Molecular epidemiology of tuberculosis in wild white-tailed deer in Michigan and elephants. In *Report of the committee on tuberculosis. Proceedings of the United States Animal Health Association*. Pat Campbell and Associates and Spectrum Press, Richmond, Virginia, pp. 543–546.
- WILLET, H. P. 1992. *Mycobacteria*. In *Zinsser Microbiology*, W. K. Joklik, H. P. Willett, D. B. Amos and C. M. Wilfert (eds.). Appleton and Lange, Norwalk, Connecticut, pp. 497–525.
- WILLIAMS, R. S., AND W. A. HOY. 1930. The viability of *B. tuberculosis (bovinus)* on pasture land, in stored faeces and in liquid manure. *Journal of Hygiene* 30: 413–419.

Received for publication 25 January 2006.