

Assessing the risk of plant species invasion under different climate change scenarios in California

Authors: Renteria, Jorge L., Skurka Darin, Gina M., and Grosholz, Edwin D.

Source: Invasive Plant Science and Management, 14(3) : 172-182

Published By: Weed Science Society of America

URL: <https://doi.org/10.1017/inp.2021.23>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Research Article

Cite this article: Renteria JL, Darin GMS, and Grosholz ED (2021) Assessing the risk of plant species invasion under different climate change scenarios in California. *Invasive Plant Sci. Manag* **14**: 172–182. doi: [10.1017/inp.2021.23](https://doi.org/10.1017/inp.2021.23)

Received: 17 March 2021
Revised: 25 June 2021
Accepted: 23 July 2021
First published online: 30 July 2021

Associate Editor:

Catherine Jarnevich, U.S. Geological Survey

Keywords:

Exotic species; ornamental plants; climate change; Species Distribution Models; invasion risk

Author for correspondence:

Jorge L. Renteria, Department of Environmental Science and Policy, University of California, Davis, CA 95616.
(Email: jrenteria@ucdavis.edu)

Assessing the risk of plant species invasion under different climate change scenarios in California

Jorge L. Renteria¹ , Gina M. Skurka Darin² and Edwin D. Grosholz³

¹Assistant Project Scientist, Department of Environmental Science and Policy, University of California, Davis, CA, USA; ²Senior Environmental Scientist, Division of Environmental Services, California Department of Water Resources, Sacramento, CA, USA and ³Professor and Cooperative Extension Specialist, Department of Environmental Science and Policy, University of California, Davis, CA, USA

Abstract

Using species distribution models (SDMs), we predicted the distribution of 170 plant species under different climatic scenarios (current and future climatic conditions) and used this information to create invasion risk maps to identify potential invasion hot spots in California. The risk of invasion by individual species was also assessed using species' predicted area in combination with some biological traits associated with invasiveness (growth form, reproduction mechanisms, and age of maturity). A higher number of species would find suitable climatic conditions along the coast; the Central Western (CW) and South Western (SW) were ecoregions where a higher number of species were predicted. Overall, hot spots of species distribution were similar under current and future climatic conditions; however, individual species' predicted area (increase or decrease) was variable depending on the climate change scenario and the greenhouse gas emission. Out of the 170 species assessed, 22% ranked as high-risk species, with herbs, grasses, and vines accounting for 78% within this risk class, and a high proportion (67%) of Asteraceae species ranked as high risk. This study suggests that current climatic conditions of the central and south coastal regions of California would be considered as hot spots of new invasions, and for some species this risk might increase with hotter and drier future climatic conditions.

Introduction

Greater undesirable impacts are expected when regions become more susceptible to the establishment of exotic plant species. The interaction of California's complex topography with its climate and habitat diversity make the state home to a wide variety of native flora (3,400 species). The region is a biodiversity hot spot that contains 20% of all vascular plant species in the United States (Stein et al. 2000). However, these conditions are also favorable for the establishment of exotic plant species (Baldwin et al. 2012; Brusati et al. 2014). More than 1,500 exotic plant species have naturalized in agricultural systems and natural areas in California (DiTomaso and Healy 2007).

Studies have shown that the ornamental horticultural industry is the main pathway for plant invasions worldwide (van Kleunen et al. 2018). Ornamental plants are produced mainly for their aesthetic value and are grown for decorative purposes in gardens and landscape design projects. Ornamental plants constitute an important part of the global horticulture industry (Li et al. 2004; Mitrofanova et al. 2018). More than 50,000 exotic plant species have been introduced for ornamental purposes in the United States; although most of these species do not represent a problem, some do escape and become invasive (Reichard and White 2001). In California, exotic ornamental species represent 47% of the total number of invasive plants according to the California Invasive Plant Council's Invasive Plant Inventory (Bell et al. 2017; Conser et al. 2015). There is a significant risk of new plant invasions in California from subsequent introductions (Brusati et al. 2014; Conser et al. 2015; Reichard and White 2001).

The increase in the proportion of greenhouse gases in the Earth's atmosphere is likely to cause an average global warming of 1 to 3.5 C over the next century (IPCC 2014). This warming will vary spatially and is predicted to be most intense in the winter at high northern latitudes (Houghton 1997). Changes in global temperatures will also bring a change in precipitation regimes, but forecasting for the magnitude or direction of these changes is unclear. California has experienced an overall warming trend over the past century, and it is projected that the temperature will continue increasing. Statewide mean temperature could increase as much as 5.8 C by 2070 to 2099 relative to 1971 to 2000 under continued high greenhouse gas emissions (Cordero et al. 2011; Pierce et al. 2018; Rapacciuolo et al. 2014). Future change in precipitation for California is less certain, with both increases and decreases in annual

© The Author(s), 2021. Published by Cambridge University Press on behalf of the Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



Management Implications

This study assesses the risk of invasion by exotic ornamental plants in California under climate change conditions. This assessment uses a relatively simple methodology and provides a basis and rationale for prioritizing areas and species of potential concern. The evaluation identifies potential hot spots of plant invasion and ranks individual species according to the calculated risk. This is the first screening stage and provides to land managers and stakeholders an additional tool to identify feasible management strategies for potentially new invaders. These risk maps can be used to develop a regional surveying strategy to look for new potential invasive plant species in California. Resources and efforts should be focused on those areas where the conditions seem to be more suitable for the establishment of a high number of potential new invasive species. This assessment allowed us to classify species within three risk categories (high, moderate, and low). Depending on species naturalization status and risk of invasion, various interventions (trade ban, cessation of cultivation, monitoring, control, containment, or eradication) can be contemplated at different scales.

precipitation projected, depending on the general circulation model (GCM) considered (Berg and Hall 2015; Dettinger et al. 2015; Pierce et al. 2013).

On a local scale, the anticipated changes in climate will directly favor some species over others, and range shifts will consequently occur (Allen and Bradley 2016; Buckley and Csergő 2017; Dukes and Mooney 1999). Some studies have shown that cultivated species could expand and become problem species over larger areas if the limiting temperatures recede (Dullinger et al. 2017; Patterson 1995). Long-term observational studies suggest that an increase in annual precipitation in arid and semiarid regions of western North America could increase the dominance of invasive alien grasses (Boyte et al. 2016; Hobbs and Mooney 1991). In contrast, other studies have predicted decreases in the potential ranges and spatial shifts of some invasive plant species under future climatic scenarios (Beerling et al. 1995; Buckley and Csergő 2017; Manzoor et al. 2018). For example, a decrease in range size was predicted for five of South Africa's major plant invaders under future climatic scenarios (Richardson et al. 2000).

Increasing temperature and precipitation combined with more frequent and intense climatic events (very wet and very dry years) are likely to favor the establishment and spread of some invasive species (Bradley et al. 2010; Guan et al. 2020; Hellmann et al. 2008). Climate change may increase the probability of invasive species establishment by providing more favorable climatic conditions in areas where exotic species are currently unable to survive (e.g., ornamental exotic plants that currently depend on the artificial climate of a garden). Climate change might also facilitate exotic species establishment by increasing their competitive abilities or rate of spread; and finally, climate change might shift native species' geographic distributions, reducing their competitive resistance (Adhikari et al. 2019; Hellmann et al. 2008).

For new potentially invasive exotic species, the use of risk maps can guide management strategies by identifying areas where new invasive species are more likely to establish and cause negative impacts (Kriticos et al. 2013; Venette et al. 2010). Species distribution models (SDMs) are often used for a diverse range of ecological studies, including biological invasion studies (Guillera-Arroita et al. 2015); SDMs use mathematical algorithms to establish a

relationship between species' occurrence data and environmental variables. These models can then be projected across the environmental space to identify areas likely to have suitable conditions for a particular species; these outputs then can be used to support management decisions (Bradley et al. 2010; Guisan et al. 2013; Mammola and Leroy 2018).

Considering the large number of plant species that could become invasive in California (Brusati et al. 2014), deciding which species should be targeted for management can be challenging. Several frameworks have been developed to assess the risk of invasion by exotic plants, including trait-scoring, statistical, decision-tree, rapid screening, and mechanistic approaches, as well as other more detailed approaches that might include comprehensive information on the species, the region where it might be introduced, and the circumstances of its introduction (Keller and Kumschick 2017; Martin et al. 2020). These pre- or post-border assessments are usually based on a series of parameters related to the species' life history and ecology and its potential environmental and socioeconomic impacts (Cunningham et al. 2004; Darin et al. 2011; Hulme 2012; Kumschick et al. 2012; Roy et al. 2018). Depending on the number of parameters and information availability, completing these risk assessment frameworks for a large number of species can be challenging (Hulme 2012; Renteria et al. 2017; Verbrugge et al. 2019).

Managers often face the challenge of deciding which species should be targeted for intervention; however, given the amount of information required to make such a decision, it is likely that those interventions will be implemented under a certain level of uncertainty (Darin et al. 2011; Kumschick and Richardson 2013). Rapid prioritization assessments provide an initial screen, allowing managers to rank and identify potentially problematic species at different scales. Generally these assessments use few parameters and are less time-consuming compared with other more detailed schemes (Branquart et al. 2016; Martin et al. 2020).

Predicting future distributions of invasive species can aid development of effective management actions such as prevention of introductions and opportunities for eradication. Proactively identifying high-risk species and areas increases resource-use efficiency by preventing new invasions through targeted surveying in managed areas (Jiménez-Valverde et al. 2011; Venette et al. 2010). Assessing the risk of new invasion is beneficial for informing stakeholders and land managers, particularly in the face of climate change (Allen and Bradley 2016; Gallagher et al. 2013; O'Donnell et al. 2012). We used SDMs to predict the current and future potential distribution of 170 exotic ornamental plants listed as potentially new invasive species for California. Using GIS analysis, we identified geographic areas most at risk of invasion under different climate change scenarios. Using individual species' predicted area together with some biological traits, the risk of invasion of each species was assessed using a weighted-score approach. We discuss how these results can be used to help prioritize both high-risk areas and species for subsequent management intervention.

Methods

For this study, we considered the exotic ornamental plant species listed as potential new invaders for California generated by Brusati et al. (2014). Their assessment was based on species' invasiveness elsewhere with a similar Mediterranean climate or species listed as invasive in a neighboring state. Their risk analysis resulted in a total of 186 species arranged in two main groups:

naturalized (species that naturalized after 1940, 70 spp.) and non-naturalized (116 spp.). Naturalized species are often defined as exotic species that have been able to reproduce and maintain viable populations for substantial number of years in the recipient area, as opposed to non-naturalized exotic species, whose fate is either extinction or persistence through human aid (Pyšek and Prach 2003). Within these two groups, the assessment also categorized species according to their availability in Californian nurseries.

Areas at Risk of Invasion

Occurrence data from the 186 species were gathered from the Global Biodiversity Information Facility database (GBIF). Data from GBIF are derived from many sources ranging from museum specimens to citizen science data; every single species' occurrence record in GBIF goes through a series of data-quality steps until it becomes available for the users (www.gbif.org). Occurrences from species' native and introduced ranges (including records from California) were considered; this approach provides the best approximation of the range of niches a species can occupy and so can be used to understand its full potential (Jiménez-Valverde et al. 2011; Verbruggen et al. 2013).

Before analysis, all records were carefully checked to match species' taxonomy. We included occurrence records with geographic coordinates having at least two decimal places. Record occurrences having ≥ 1 km error or uncertainty (suspicious outliers) associated with the geographic coordinates were discarded. To avoid pseudo-replication, only one record per ~ 4.5 km² grid cell (based on the climatic variables resolution, 2.5 minutes) was used for model calibration. It is likely the data may exhibit spatial bias due to sampling effort, because the occurrences were not collected using a specific sampling methodology (Phillips et al. 2009). To reduce the geographic sampling biases, a geographic thinning (1 record per 4.5 km) was performed on all occurrences using the *spThin* R package (Aiello-Lammens et al. 2015). Species with fewer than 30 occurrences were not included; therefore, the analysis was carried out with 170 species (65 naturalized and 105 non-naturalized).

We used three SDMs to predict the species' potential distribution: a generalized linear model, a random forest model, and a support vector machine model. Six climatic variables identified as important in driving plant distributions in the western United States (Rehfeldt et al. 2006; Stephenson 1998) were considered as predictors of species occurrence: annual mean temperature, maximum temperature of warmest month, minimum temperature of coldest month, annual precipitation, precipitation of wettest month, and precipitation of driest month. These six bioclimatic variables (raster layers at 2.5 arc-minute resolution, historical climate data for 1970 to 2000) were acquired from the WorldClim database (Fick and Hijmans 2017). We examined collinearity among the six bioclimatic variables by running a Pearson correlation analysis. Annual mean temperature and annual precipitation were highly correlated (Pearson $r > 0.7$) to the other climatic variables and therefore were not considered to reduce the negative impact of multicollinearity in the modeling process.

Modeling was performed by randomly splitting the records into a calibration set (70% of the records) and a test set (30% of the records). The three SDM algorithms used in this study require absences or background data. Pseudo-absences were generated within the extent of the environmental rasters defined by

the maximum and minimum latitude and longitude values from all the species occurrence data (Vasquez et al. 2021). Using the selected method repeatedly under the SSDM package (default parameters) (Schmitt et al. 2017), a set of pseudo-absences was automatically generated for each SDM following recommendations from Barbet-Massin et al. (2012) (e.g., for GLM: 10 runs of 1,000 randomly selected pseudo-absences; for RF and SVM: same as number of presences, 100 or fewer presence points, a minimum of 10 runs with 100 pseudo-absences). Each model was run five times, and the average AUC (area under the receiver operating characteristic curve) was used to evaluate model performance. AUC values vary from 0 to 1; values below 0.7 represent poor model performance, whereas AUC values close to 1 indicate a high predictive model performance. We also assessed the variable relative importance generated by the SSDM package (Schmitt et al. 2017), which computes a simple Pearson's correlation r between predictions of the full model and the one without a variable and returns the score $1 - r$: the higher the value, the more influence the variable has on the model.

Models were projected onto the California near-current (years 1970 to 2000, hereafter referred to as "current") and future (year 2040) climatic conditions. Two GCMs (CNRM and MIROC) listed as good simulations of California's historic climate (Pierce et al. 2018) were used as future climatic scenarios. These two GCMs represent scenarios with the most extreme directional changes in precipitation (CNRM-wettest; MIROC-driest) and have shown an effect on species' habitat suitability prediction in California (Riordan et al. 2018). For each GCM, we considered two greenhouse gas scenarios (representative concentration pathways: RCP 4.5 and RCP 8.5) to create future projections. Each scenario defines a pathway in terms of the concentration of carbon in the atmosphere at any date: RCP 4.5 represents a target forcing of 4.5 W m^{-2} above the preindustrial baseline by 2100 and delivers a temperature increase of about 1.8°C ; RCP 8.5 corresponds to a high greenhouse gas emissions pathway of 8.5 W m^{-2} and delivers a temperature increase of about 3.7°C by 2100 (IPCC 2014). Future bioclimatic variables generated for the two GCMs (CNRM-CM6-1 and MIROC6) and the two greenhouse gas emissions (RCP 4.5 and 8.5) were acquired from the WorldClim database (raster layers at 2.5 arc-minute resolution, future climate data for year 2040) (Fick and Hijmans 2017).

Species' suitability maps generated by each of the three models were transformed to binary maps (presence/absence maps) using the maximum sum of sensitivity and the specificity as a threshold cutoff value. With this approach, a threshold value for each species was calculated to maximize the agreement between the observed and predicted distribution (Liu et al. 2013). Individual species' binary maps were combined, and the potential distribution for each individual species was calculated as the grid cells where at least two out of three binary maps predicted the species occurrence. Finally, the potential distributions of the 170 species were merged to produce the risk maps for California—resulting in five main risk maps: current climatic conditions, two GCMs, and two RCPs. To identify species-rich areas or hot spots of invasion, risk maps' cells were classified into four risk categories based on the number of species predicted to occur in a given cell (four equal intervals, range between 1 and the highest number of species predicted). All analyses were conducted using the SSDM package in the R environment and ArcGIS Desktop (ArcGIS Version 10.0, Environmental Systems Research Institute, Inc. [ESRI], Redlands, CA; R Core Team 2018; Schmitt et al. 2017).

Table 1. List of parameters, range values, and scores used to run the species' risk assessment.^a

Species' potential distribution			Species' biological traits		
Parameter	Range	Score	Parameter	Range	Score
Predicted area ^b	<1 %	0	Growth form	Tree	1
	1%–20 %	1		Shrub	2
	21%–40 %	2		Herb/grass/vine	3
	41%–60 %	3			
Number of ecoregions ^c	0	0	Reproduction	Seeds	1
	1–3	1		Seeds and vegetative	2
	4–6	2	Age maturity	>3 yr	1
	7–10	3		1–3 yr	2
Predicted area variation ^d	No increase	0		Within a year	3
	Increase under one GCM	1			
	Increase under both GCMs	2			

^aSpecies' parameters were scored from 0 to 3, the total score (sum of all parameter scores for a species) for a given species could vary from 3 to 16.

^bSpecies' predicted area in relation to California's total area.

^cNumber of ecoregions intersected by a species' predicted area (Hickman 1993).

^dDifference between predicted area under current climatic conditions and predicted area under future climatic scenarios. For each general circulation model (GCM), an increase in predicted area needed to occur under both emission scenarios (representative concentration pathways [RCPs] 4.5 and 8.5) to be categorized as an increase for the GCM.

Table 2. Relative variable importance (VI; average percentage \pm SE) across all species and percentage of species with the highest VI for each species distribution model (SDM): generalized linear model (GLM), random forest model (RF), and support vector machine model (SVM).

Climatic variable	Relative VI			Percentage of species with highest VI		
	GLM	RF	SVM	GLM	RF	SVM
Maximum temperature of warmest month	20.7 \pm 0.9	23.6 \pm 0.7	24.2 \pm 0.7	7.6	10	10.6
Minimum temperature of coldest month	45.9 \pm 1.6	45.0 \pm 1	43.5 \pm 1	72.4	79.4	74.1
Precipitation of wettest month	17 \pm 1.1	14.9 \pm 0.6	16.1 \pm 0.9	6.5	4.1	5.3
Precipitation of driest month	16.3 \pm 1.8	16.5 \pm 0.7	16.3 \pm 1	13.5	6.5	10

Species Risk Assessment

A rapid-invasion risk assessment was carried out using the species' potential distribution and some biological traits. The species' potential distribution (predicted area) under current climatic condition was evaluated in relation to: California total area, number of ecoregions overlapping with predicted area, and the variation of the predicted area relative to the prediction under two future climatic scenarios (GCMs: CNRM and MIROC; RCPs: 4.5 and 8.5). Higher risk of establishment was given to species with a broader predicted area within California, overlapping with various ecoregions, and an increase in predicted area under two future climatic scenarios (Supplementary Table S1).

Three species' biological traits were considered as factors that could facilitate the invasion process: growth form, reproduction mechanisms, and age of maturity. Studies suggest that herbs, grasses, and vines have higher invasion rates than other plant growth forms (Anning and Yeboah-Gyan 2007; Godoy et al. 2009; Johnson et al. 2020). Reproduction is a key factor in plant invasions. Effective reproduction mechanisms enable invasive plants to produce a large number of propagules to establish new populations and spread (Barrett 2011; Burns et al. 2013). Plant invasions have also correlated with high relative growth rate, small seed masses, and short juvenile period (Grotkopp et al. 2010; Rejmanek 1996). A higher risk was assumed for species classified as herbs/grasses/vines, with multiple reproduction mechanisms, and with a short time to reach maturity; information on species' traits was gathered from publications and Internet resources (Supplementary Table S2).

The assessment was carried out using a scored approach assigning a value from 0 to 3 to each of the parameters (species' potential distribution and biological traits). A total score was

calculated by adding the individual score from each parameter; the lower the species' total score, the lower the risk of it becoming invasive (Table 1).

Results and Discussion

Areas at Risk of Invasion

Species distribution models were fit for 170 species (65 naturalized and 105 non-naturalized) of the total 186 listed as potentially invasive plant species for California (Brusati et al. 2014). On average, the three models showed a very good AUC evaluation: GLM = 0.82 \pm 0.004 SE, RF = 0.93 \pm 0.002 SE, and SVM = 0.91 \pm 0.003 SE. GLM models for two species showed an AUC below 0.7; however, these values were reasonably high (AUC = 0.68) (Supplementary Table S3). Therefore, these models were still considered to predict species' distribution. Overall, the temperature of coldest month was the variable that consistently had the higher predictive power, whereas precipitation of wettest month had low importance across the three models (Table 2).

The analysis predicted that under current climatic conditions, 99% of California shows suitable climatic conditions for at least one species. Most of California (78%) shows suitable conditions for 1 to 30 species, whereas a greater number of species (91 to 125) are predicted to occur in only 3.3% of the total area (Table 3). A high number of species are predicted to occur along the coast, particularly in the Central Western (CW) and South Western (SW) ecoregions, where 46% and 27% of these regions could be suitable for more than 60 species. The CW could be considered as a potential hot spot for new invasions, while a small number of species (1 to 30) are predicted in inland regions (Figure 1).

Table 3. Variation in the percentage of predicted area by risk categories (no species predicted) relative to predicted area under current and future climatic conditions (GCMs: CNRM and MIROC; RCPs: 4.5 and 8.5).^a

Risk category	Current predicted area	Variation in area ^b			
		CNRM		MIROC	
		RCP 4.5	RCP 8.5	RCP 4.5	RCP 8.5
no. of species	%			%	
0	0.8	−0.6	−0.6	3.6	4.2
1–30	78.7	4.8	4.9	−4.6	−3.7
31–60	12.1	−2.6	−2.7	0.9	−0.4
61–90	5.1	−1.0	−1.0	0.1	0.1
91–125	3.3	−0.5	−0.7	0.1	−0.2

^aGCM, general circulation model; RCP, representative concentration pathway.
^bNegative values represent reductions in area with climate change.




Figure 1. California invasion risk map created by combining the 170 species' potential distribution under current and future climatic conditions; for future climatic conditions (year 2040; global circulation models [GCMs]: CNRM, and MIROC; greenhouse emissions: representative concentration pathways [RCP] 4.5, and 8.5). Codes correspond to Jepson ecoregions; CaR: Cascade Ranges; CW: Central Western CA; SNE: East of Sierra Nevada; GV: Great Valley; MP: Modoc Plateau; DMoj: Mojave Desert; NW: North Western CA; SN: Sierra Nevada; DSon: Sonoran Desert; SW: Southwestern CA (Hickman 1993).




Figure 2. Effect of future climatic conditions (year 2040; representative concentration pathways [RCP] 4.5 and 8.5) on species' predicted area in relation to species' predicted area under current climatic conditions: (A) under climatic conditions generated by GCM CNRM and (B) under climatic conditions generated by GCM MIROC.

Overall, species' distribution predicted under future climatic scenarios shows similar patterns to predictions under current climatic conditions (Figure 1). Climatic suitability for a greater number of species is predicted along the coast compared with the inland regions. Some level of variation on the predicted species richness by the two GCMs can be observed (Table 3). GCM CNRM predicted a decrease on the number of cells with high species richness under both greenhouse emissions (RCP 4.5 and 8.5). GCM MIROC predicted an increase on species richness under RCP 4.5 and a decrease under RCP 8.5. The effect of the two climatic scenarios (CNRM and MIROC) on individual species' predicted area was different (Figure 2). Under climatic scenario CNRM, 20% of the species showed an increase in predicted area versus 47% under climatic scenario MIROC.

Species Risk Assessment

The risk of invasion for 170 ornamental plants listed as future potential invasive species for California was investigated using species' potential distribution and some biological and ecological attributes. The species include 64 families, for which Fabaceae (21 spp.), Asteraceae (15 spp.), and Iridaceae (10 spp.) account for 27% of the total number of species. Regarding growth form, the species list includes 76 herbs/grasses/vines (45%), 54 woody/shrubs (32%), and 40 tree species (23%); as per habitat preference, 94% are terrestrial species (Supplementary Table S2).

The calculated species' total risk score varied from 3 to 15. Using a three equal score intervals (lowest score +4), species were grouped within three risk categories: low risk (total score: 3 to 7), moderate risk (total score: 8 to 11), and high risk (total score: 12 to 15) (Table 4; Supplementary Table S4). The evaluation ranked 38 species on top of the list as high-risk species, 112 species as moderate risk, and 20 species as low risk. Regarding families, 67% of the Asteraceae species and 30% of the Iridaceae species ranked as high risk, whereas most of the Fabaceae species ranked as moderate and

low risk (76% and 24%, respectively). Out of the 38 species categorized as high-risk species, herbs/grasses/vines accounted for 78%.

According to Brusati et al. (2014), out of the 170 species used on this assessment, 65 have been recorded as naturalized and 105 as non-naturalized in California. Because naturalized species are already established in California, the probability of these species becoming invasive is considerably higher than for non-naturalized species; results of the assessment are presented by naturalization status (Table 4). Within naturalized species, the risk assessment ranked 19 species as high risk, a second group of 41 species as moderate risk (score: 8 to 11), and 5 species as low risk. Species such as lantana (*Lantana camara* L.), french lavender (*Lavandula stoechas* L.), five-stamen tamarisk (*Tamarix chinensis* Lour.), and birdwood grass [*Pennisetum ciliare* (L.) Link] ranked on top, and these species are considered wildfire hazards (Supplementary Table S2). Within non-naturalized species, 19 species were ranked as high risk, 71 species as moderate risk, and 15 as low risk. Overall, the proportion of the number of species by risk categories follows the naturalization status trend (Figure 3A). Most of the naturalized and non-naturalized species were ranked as moderate risk; however, the number of species classified as high risk is greater for naturalized species. Within the growth forms, most of the species were classified as moderate risk (Figure 3B). However, the proportion of herbs, grasses, and vine species ranked as high risk was considerably greater than for shrubs and trees.

Using the potential distribution of 170 plant species, we created risk maps of invasion under "current" and future climatic conditions for California. The influence of climatic conditions on species distribution at regional scales is well known (Chapin and Díaz 2020). The use of temperature and precipitation as predictors of species distribution is very common and has provided a reasonably good approximation of species' environmental suitability (Bradie and Leung 2017; Bucklin et al. 2015). The contributions of the four climatic variables were consistent across models and species (Table 1). Results of the modeling indicate that extreme

Table 4. Species ranked as top 10 according to the assessment.^a

Species	Species' potential distribution						Species' biological traits						
	Predicted area		Predicted area increase		Ecoregions		Growth form		Reproductive mechanisms		Age of maturity		Total score
	Percentage of California	s	GCMs	s	No.	s		s		s			
Naturalized													
<i>Pennisetum ciliare</i> (L.) Link	53.0	3	CNRM	1	9	3	H,G,V	3	S&V	2	Within a year	3	15
<i>Lantana camara</i> L.	13.9	1	CNRM and MIROC	2	6	2	H,G,V	3	S&V	2	Within a year	3	13
<i>Lavandula stoechas</i> L.	18.8	1	MIROC	1	7	3	H,G,V	3	S&V	2	Within a year	3	13
<i>Osteospermum ecklonis</i> (DC) Norl.	48.1	3	Neither	0	10	3	Shrub	2	S&V	2	Within a year	3	13
<i>Osteospermum fruticosum</i> (L.) Norl.	23.3	2	MIROC	1	10	3	H,G,V	3	S&V	2	1–3 yr	2	13
<i>Tamarix chinensis</i> Lour.	62.5	3	Neither	0	10	3	Shrub	2	S&V	2	Within a year	3	13
<i>Cabomba caroliniana</i> A. Gray	17.7	1	Neither	0	8	3	H,G,V	3	S&V	2	Within a year	3	12
<i>Coreopsis lanceolata</i> L.	11.6	1	MIROC	1	9	3	H,G,V	3	S&V	2	1–3 yr	2	12
<i>Gazania linearis</i> (Thunb.) Druce	16.1	1	MIROC	1	7	3	H,G,V	3	Seeds	1	Within a year	3	12
<i>Helianthus tuberosus</i> L.	0.3	0	CNRM and MIROC	2	3	2	H,G,V	3	S&V	2	Within a year	3	12
Non-naturalized													
<i>Argemone ochroleuca</i> Sweet	36.0	2	CNRM	1	8	3	H,G,V	3	Seeds	1	Within a year	3	13
<i>Glandularia pulchella</i> (Sweet) Tronc.	36.3	2	MIROC	1	10	3	H,G,V	3	Seeds	1	Within a year	3	13
<i>Periploca graeca</i> L.	21.7	2	MIROC	1	9	3	H,G,V	3	S&V	2	1–3 yr	2	13
<i>Schkuhria pinnata</i> Lam.	40.8	3	Neither	0	9	3	H,G,V	3	Seeds	1	Within a year	3	13
<i>Ageratina riparia</i> (Regel) R.M.King & H.Rob.	3.4	1	MIROC	1	3	2	H,G,V	3	S&V	2	Within a year	3	12
<i>Alpinia zerumbet</i> (Pers.) B.L.Burt & R.M.Sm	3.0	1	CNRM and MIROC	2	4	2	H,G,V	3	S&V	2	1–3 yr	2	12
<i>Asparagus plumosus</i> Baker	14.1	1	CNRM and MIROC	2	7	3	H,G,V	3	Seeds	1	1–3 yr	2	12
<i>Canna indica</i> L.	6.8	1	MIROC	1	3	2	H,G,V	3	S&V	2	Within a year	3	12
<i>Coleostephus myconis</i> (L.) Rchb.fil.	16.9	1	MIROC	1	7	3	H,G,V	3	Seeds	1	Within a year	3	12
<i>Freesia leichtlinii</i> Klatt	21.3	2	Neither	0	8	3	H,G,V	3	S&V	2	1–3 yr	2	12

^aList of parameters and scores (s) used for the risk of invasion evaluation: growth form: H,G,V, herb, grass, or vine; reproductive mechanisms: S&V, seeds and vegetative.

^bGCM, general circulation model.

temperatures explained the distribution of most of the species used for this assessment.

Our analysis revealed that most of California would have suitable climatic conditions for a relatively low number of species. A higher number of species are predicted along the coastline, with the highest concentration in the CW and SW regions. Studies have shown that an invasive species is more likely to invade areas with conditions similar to those where it is indigenous (Thuiller et al. 2005). Most of the species used for this assessment originate from regions with a Mediterranean climate; therefore, suitable conditions for these species would be expected along the coastline, where the climatic conditions are likely to match species' climate requirements. These areas identified as having a high risk for invasion are also known to be rich in plant biodiversity and endemism (Kraft et al. 2010; Loarie et al. 2008). There is great potential for

undesirable impacts to these ecologically valuable and vulnerable ecoregions—particularly to evergreen and deciduous forest, woodland, chaparral, and open grassland vegetation types. As in other Mediterranean regions, the climate in California is defined by cool wet winters and hot dry summers (Harrison et al. 2020; Rundel et al. 2016). These climatic conditions, which vary considerably by ecoregion, have shaped native plant diversity, and vegetation communities might also be an important factor for future plant species establishment (Lenihan et al. 2003; Pyšek et al. 2017).

Future species' distribution varied widely depending on the climatic scenario considered. The wettest climatic scenario, CNRM (RCP 4.5 and 8.5), projected a reduction in invasive species richness, whereas the driest scenario, MIROC (RCP 4.5), projected an increase. Contrary to the negative effects that extreme climatic conditions such as those generated by climatic scenario MIROC




Figure 3. Number of species within the invasion risk categories: (A) by naturalized status in California and (B) by growth forms.

might have on California native vegetation (Riordan et al. 2018), a higher number of species (~50%) increased the predicted distribution under the MIROC climatic scenario. Extreme variations in climatic conditions are expected to have important impacts on distribution patterns of native and invasive plant species (Lenihan et al. 2003; Sandel and Dangremond 2012); however, the magnitude and direction of these impacts might depend on the climate change scenario and the species-specific responses to climatic conditions (Bellard et al. 2013; Finch et al. 2021; Guan et al. 2020; Petitpierre et al. 2016).

Informing government and society about areas at risk of invasion is necessary to guide management efforts and secure economic resources (del-Val et al. 2015). To our knowledge, this is one of the first assessments using a large number of species to produce an invasion risk map for California. These maps should be used as an early screening tool to identify potential areas suitable for invasion and spread (Montemayor et al. 2015; Pearson and Dawson 2003), allowing for a coarse identification of areas where effort should be focused to look for particular plants or areas at risk.

The list of ornamental plant species produced by Brusati et al. (2014) was created on the basis of species invasiveness in regions with similar climatic conditions or states neighboring California. Although all these species are at risk of becoming invasive in California, given the variety of climatic and topographic conditions, the magnitude of the risk would be expected to vary between species. Using the species' potential distribution and some biological attributes, we performed a rapid screening to rank and categorize species according to the calculated invasion risk. High-risk naturalized species should have the highest priority; these species have established and created self-sustainable populations, making them more likely to become invasive. For these species, monitoring should be considered to assess the spatial distribution and population dynamics with the aim of detecting species invasiveness behavior. As a precautionary principle for non-naturalized high-risk species, interventions should be focused on preventing species arrival; for those already introduced, state trade regulation might be required.

The scope of the assessment is the entire state of California; however, the list can be modified to generate a list of priority species at local scales. Although this rapid assessment does not replace other, more detailed risk assessment schemes, it can be used as an initial step in the prioritization process. Moreover, this assessment should be considered to be a dynamic process; the status of some species might change, and new invasive species are likely to arrive, so this ranking and categorization will need to be modified accordingly (Conser et al. 2015).

Predicting the distribution of an invasive species is not an easy task, and it becomes even more challenging when the exercise includes a large number of species. The invasion process is complex and involves the interaction of several biotic and abiotic factors that might influence the species' establishment and spread (Gantchoff et al. 2018; Lee and Lee 2006). Additionally, predicted distributions are sensitive to data and modeling processes (Sofaer et al. 2019; Zurell et al. 2020). As with any modeling effort, our approach is subject to constraints and limitations; for example, using records from species' native and introduced ranges to fit the models risks overestimating species' predicted distributions (Bradley 2013; Jiménez-Valverde et al. 2011). Further, it is likely the data may exhibit spatial bias due to sampling effort, because the occurrences were not collected using a specific sampling methodology (Phillips et al. 2009). Additionally, studies have shown variability among model predictions of species moving into new environments (Araújo and New 2007; Webber et al. 2011). We have tried to address limitations (e.g., selecting uncorrelated environmental variables, applying geographic thinning to reduce the geographic sampling biases, using three SDMs to reduce variability, generating pseudo-absences according to each model), aiming to reduce the source of error and increase the models' predictive ability. Species' predicted distributions and future shifts in range are approximations and do not represent an absolute measure of site suitability or change. Our risk analysis focuses on distribution patterns across the landscape rather than an accurate potential invasion area.

Despite assumptions and difficulties in evaluating predictions accurately (Araújo and Peterson 2012; Barbet-Massin et al. 2018), SDMs are a valuable tool to assess invasion risk and assist

in designing effective management strategies (Bradley et al. 2010; Barbet-Massin et al. 2018). This risk analysis is an important step toward the development of early warning systems to prevent the arrival or establishment of new potential invasive plant species in California.

Acknowledgments. We are grateful for the contributions made by Rachel Atkinson and Ryan Boynton to earlier versions of this article. This research received no specific grant from any funding agency or the commercial or not-for-profit sectors. No conflicts of interest have been declared.

Supplementary material. To view supplementary material for this article, please visit <https://doi.org/10.1017/inp.2021.23>

References

- Adhikari P, Jeon J-Y, Kim HW, Shin M-S, Adhikari P, Seo C (2019) Potential impact of climate change on plant invasion in the Republic of Korea. *J Ecol Environ* 43:1–12
- Aiello-Lammens ME, Boria RA, Radosavljevic A, Vilela B, Anderson RP (2015) spThin: an R package for spatial thinning of species occurrence records for use in ecological niche models. *Ecography* 38:541–545
- Allen JM, Bradley BA (2016) Out of the weeds? Reduced plant invasion risk with climate change in the continental United States. *Biol Conserv* 203:306–312
- Anning AK, Yeboah-Gyan K (2007) Diversity and distribution of invasive weeds in Ashanti Region, Ghana. *Afr J Ecol* 45:355–360
- Araújo MB, New M (2007) Ensemble forecasting of species distributions. *Trends Ecol Evol* 22:42–47
- Araújo MB, Peterson AT (2012) Uses and misuses of bioclimatic envelope modeling. *Ecology* 93:1527–1539
- Baldwin BG, Goldman D, Keil DJ, Patterson R, Rosatti TJ, Wilkin D (2012) The Digital Jepson Manual: Vascular Plants of California. 2nd ed. Oakland, CA: University of California Press
- Barbet-Massin M, Jiguet F, Albert CH, Thuiller W (2012) Selecting pseudo-absences for species distribution models: how, where and how many? *Methods Ecol Evol* 3:327–338
- Barbet-Massin M, Rome Q, Villemant C, Courchamp F (2018) Can species distribution models really predict the expansion of invasive species? *PLoS ONE* 13:1–14
- Barrett SCH (2011) Why reproductive systems matter for the invasion biology of plants. Pages 195–210 in Richardson DM, ed. *Fifty Years of Invasion Ecology: The Legacy of Charles Elton*. Chichester, UK: Blackwell
- Beerling DJ, Huntley B, Bailey JP (1995) Climate and the distribution of *Fallopia japonica*: use of an introduced species to test the predictive capacity of response surfaces. *J Veg Sci* 6:269–282
- Bell CE, DiTomaso JM, Wilen CA (2017) Invasive Plants. Integrated Pest Management for Home Gardeners and Landscape Professionals. Pest Notes 7. Oakland, CA: University of California Agriculture & Natural Resources Publication 74139. <http://ipm.ucanr.edu/PDF/PESTNOTES/pninvasiveplants.pdf>. Accessed: August 6, 2021
- Bellard C, Thuiller W, Leroy B, Genovesi P, Bakkenes M, Courchamp F (2013) Will climate change promote future invasions? *Global Change Biol* 19:3740–3748
- Berg N, Hall A (2015) Increased interannual precipitation extremes over California under climate change. *J Clim* 28:6324–6334
- Boyte SP, Wylie BK, Major DJ (2016) Cheatgrass percent cover change: comparing recent estimates to climate change–driven predictions in the northern Great Basin. *Rangeland Ecol Manag* 69:265–279
- Bradie J, Leung B (2017) A quantitative synthesis of the importance of variables used in MaxEnt species distribution models. *J Biogeogr* 44:1344–1361
- Bradley BA (2013) Distribution models of invasive plants over-estimate potential impact. *Biol Invasions* 15:1417–1429
- Bradley BA, Wilcove DS, Oppenheimer M (2010) Climate change increases risk of plant invasion in the Eastern United States. *Biol Invasions* 12:1855–1872
- Branquart E, Brundu G, Buholzer S, Chapman D, Ehret P, Fried G, Starfinger U, van Valkenburg J, Tanner R (2016) A prioritization process for invasive alien plant species incorporating the requirements of EU Regulation no. 1143/2014. *EPO Bull* 46:603–617
- Brusati E, Johnson D, DiTomaso J (2014) Predicting invasive plants in California. *Calif Agric* 68:89–95
- Buckley YM, Csörgő AM (2017) Predicting invasion winners and losers under climate change. *Proc Natl Acad Sci USA* 114:4040–4041.
- Bucklin DN, Basille M, Benscoter AM, Brandt LA, Mazzotti FJ, Romanach SS, Speroterra C, Watling JJ (2015) Comparing species distribution models constructed with different subsets of environmental predictors. *Divers Distrib* 21:23–35
- Burns JH, Pardini EA, Schutzenhofer MR, Chung YA, Seidler KJ, Knight TM (2013) Greater sexual reproduction contributes to differences in demography of invasive plants and their noninvasive relatives. *Ecology* 94:995–1004
- Chapin FS, Diaz S (2020) Interactions between changing climate and biodiversity: shaping humanity's future. *Proc Natl Acad Sci USA* 117:6295–6296
- Conser C, Seebacher L, Fujino DW, Reichard S, DiTomaso JM (2015) The development of a plant risk evaluation (PRE) tool for assessing the invasive potential of ornamental plants. *PLoS ONE* 10:e0121053
- Cordero EC, Kessomkiat W, Abatzoglou J, Maugé SA (2011) The identification of distinct patterns in California temperature trends. *Clim Change* 108:357–382
- Cunningham DC, Barry SC, Woldendorp G, Burgess MB (2004) A framework for prioritizing sleeper weeds for eradication. *Weed Technol* 18:1189–1193
- Darin GMS, Schoenig S, Barney JN, Panetta FD, DiTomaso JM (2011) WHIPPET: a novel tool for prioritizing invasive plant populations for regional eradication. *J Environ Manage* 92:131–139
- del-Val E, Balvanera P, Castellarini F, Espinosa-García FJ, Murguía M, Pacheco C (2015) Identifying areas of high invasion risk: a general model and an application to Mexico. *Rev Mex Biodivers* 86:208–216
- Dettinger M, Udall B, Georgakakos A (2015) Western water and climate change. *Ecol Appl* 25:2069–2093
- DiTomaso JM, Healy EA (2007) Weeds of California and Other Western States. Oakland, CA: University of California Agriculture & Natural Resources
- Dukes JS, Mooney HA (1999) Does global change increase the success of biological invaders? *Trends Ecol Evol* 14:135–139
- Dullinger I, Wessely J, Bossdorf O, Dawson W, Essl F, Gatteringer A, Klöner G, Kreft H, Kuttner M, Moser D, Pergl J, Pyšek P, Thuiller W, van Kleunen M, Weigelt P, et al. (2017) Climate change will increase the naturalization risk from garden plants in Europe. *Global Ecol Biogeogr* 26:43–53
- Fick SE, Hijmans RJ (2017) WorldClim 2: new 1km spatial resolution climate surfaces for global land areas. *Int J Climatol* 37:4302–4315
- Finch DM, Butler JL, Runyon JB, Fetting CJ, Kilkenny FF, Jose S, Frankel SJ, Cushman SA, Cobb RC, Dukes JS, Hicke JA, Amelon SK (2021) Effects of climate change on invasive species. Pages 57–83 in Poland TM, Patel-Weyand T, Finch DM, Miniati CF, Hayes DC, Lopez VM, eds. *Invasive Species in Forests and Rangelands of the United States: A Comprehensive Science Synthesis for the United States Forest Sector*. Cham, Switzerland: Springer International
- Gallagher RV, Duursma DE, O'Donnell J, Wilson PD, Downey PO, Hughes L, Leishman MR (2013) The grass may not always be greener: projected reductions in climatic suitability for exotic grasses under future climates in Australia. *Biol Invasions* 15:961–975
- Gantchoff MG, Wilton CM, Belant JL (2018) Factors influencing exotic species richness in Argentina's national parks. *PeerJ* 6:e5514
- Godoy O, Richardson DM, Valladares F, Castro-Díez P (2009) Flowering phenology of invasive alien plant species compared with native species in three Mediterranean-type ecosystems. *Ann Bot* 103:485–494
- Grotkopp E, Rejmánek M, Rost TL, Rejmanek M, Rost TL (2010) Toward a causal explanation of plant invasiveness: seedling growth and life-history strategies of 29 pine (*Pinus*) species. *Am Nat* 159:396–419
- Guan B, Guo H, Chen S, Li D, Liu X, Gong X, Ge G (2020) Shifting ranges of eleven invasive alien plants in China in the face of climate change. *Ecol Inform* 55:101024
- Guillera-Arroita G, Lahoz-Monfort JJ, Elith J, Gordon A, Kujala H, Lentini PE, McCarthy MA, Tingley R, Wintle BA (2015) Is my species distribution model fit for purpose? Matching data and models to applications. *Global Ecol Biogeogr* 24:276–292

- Guisan A, Tingley R, Baumgartner JB, Naujokaitis-Lewis I, Sutcliffe PR, Tulloch AIT, Regan TJ, Brotons L, McDonald-Madden E, Mantyka-Pringle C (2013) Predicting species distributions for conservation decisions. *Ecol Lett* 16:1424–1435
- Harrison S, Spasojevic MJ, Li D (2020) Climate and plant community diversity in space and time. *Proc Natl Acad Sci USA* 117:4464–4470
- Hellmann JJ, Byers JE, Bierwagen BG, Dukes JS (2008) Five potential consequences of climate change for invasive species. *Conserv Biol* 22:534–543
- Hickman JC (1993) *The Jepson Manual: Higher Plants of California*. Berkeley: University of California Press. 1424 p
- Hobbs RJ, Mooney HA (1991) Effects of rainfall variability and gopher disturbance on serpentine annual grassland dynamics. *Ecology* 72:59–68
- Houghton J (1997) *Global Warming: The Complete Briefing*. 2nd ed. Cambridge University Press, Cambridge. 437 p
- Hulme PE (2012) Weed risk assessment: a way forward or a waste of time? *J Appl Ecol* 49:10–19
- [IPCC] Intergovernmental Panel on Climate Change (2014) *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate*. Geneva, Switzerland: IPCC. 169 p
- Jiménez-Valverde A, Peterson AT, Soberón J, Overton JM, Aragón P, Lobo JM (2011) Use of niche models in invasive species risk assessments. *Biol Invasions* 13:2785–2797
- Johnson LR, Trammell TLE, Bishop TJ, Barth J, Drzyzga S, Jantz C (2020) Squeezed from all sides: urbanization, invasive species, and climate change threaten riparian forest buffers. *Sustainability* 12:1448
- Keller RP, Kumschick S (2017) Promise and challenges of risk assessment as an approach for preventing the arrival of harmful alien species. *Bothalia-African Biodivers Conserv* 47:1–8
- Kraft NJB, Baldwin BG, Ackerly DD (2010) Range size, taxon age and hotspots of neoenidism in the California flora. *Divers Distrib* 16:403–413
- Kriticos DJ, Venette RC, Baker RHA, Brunel S, Koch FH, Rafoss T, der Werf W, Worner SP (2013) Invasive alien species in the food chain: advancing risk assessment models to address climate change, economics and uncertainty. *NeoBiota* 18:1–7
- Kumschick S, Bacher S, Dawson W, Heikkilä J, Sendek A, Pluess T, Robinson TB, Ingolf K (2012) A conceptual framework for prioritization of invasive alien species for management according to their impact. *NeoBiota* 15:69–100
- Kumschick S, Richardson DM (2013) Species-based risk assessments for biological invasions: advances and challenges. *Divers Distrib* 19:1095–1105
- Lee H, Lee C (2006) Environmental factors affecting establishment and expansion of the invasive alien species of tree of heaven (*Ailanthus altissima*) in Seoripool Park, Seoul. *Integr Biosci* 10:27–40
- Lenihan JM, Drapek R, Bachelet D, Neilson RP (2003) Climate change effects on vegetation distribution, carbon, and fire in California. *Ecol Appl* 13:1667–1681
- Li Y, Cheng Z, Smith WA, Ellis DR, Chen Y, Zheng X, Pei Y, Luo K, Zhao D, Yao Q, Duan H, Li Q (2004) Invasive ornamental plants: problems, challenges, and molecular tools to neutralize their invasiveness. *CRC Crit Rev Plant Sci* 23:381–389
- Liu C, White M, Newell G (2013) Selecting thresholds for the prediction of species occurrence with presence-only data. *J Biogeogr* 40:778–789
- Loarie SR, Carter BE, Hayhoe K, McMahon S, Moe R, Knight CA, Ackerly DD (2008) Climate change and the future of California's endemic flora. *PLoS ONE* 3:e2502
- Mammola S, Leroy B (2018) Applying species distribution models to caves and other subterranean habitats. *Ecography* 41:1194–1208
- Manzoor SA, Griffiths G, Iizuka K, Lukac M (2018) Land cover and climate change may limit invasiveness of *Rhododendron ponticum* in Wales. *Front Plant Sci* 9:664
- Martin CD, Jewell SD, Hoff MH, Givens CE, Marcot BG (2020) Comparing invasive species risk screening tools FISRAM, ERSS, and FISK/AS-ISK as a response to Hill et al. (2020). *Manag Biol Invasions* 11:342
- Mitrofanova IV, Zakubanskiy AV, Mitrofanova OV (2018) Viruses infecting main ornamental plants: an overview. *Ornam Hort* 24:95–102
- Montemayor SI, Dellapé PM, Melo MC (2015) Predicting the potential invasion suitability of regions to cassava lacebug pests (Heteroptera: Tingidae: *Vatiga* spp.). *Bull Entomol Res* 105:173–181
- O'Donnell J, Gallagher R V, Wilson PD, Downey PO, Hughes L, Leishman MR (2012) Invasion hotspots for non-native plants in Australia under current and future climates. *Global Change Biol* 18:617–629
- Patterson D (1995) Weeds in a changing environment. *Weed Sci* 43:685–701
- Pearson RG, Dawson TP (2003) Predicting the impacts of climate change on the distribution of species: are bioclimate envelope models useful? *Global Ecol Biogeogr* 12:361–371
- Petitpierre B, McDougall K, Seipel T, Broennimann O, Guisan A, Kueffer C (2016) Will climate change increase the risk of plant invasions into mountains? *Ecol Appl* 26:530–544
- Phillips SJ, Dudik M, Elith J, Graham CH, Lehmann A, Leathwick J, Ferrier S (2009) Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. *Ecol Appl* 19:181–197
- Pierce DW, Das T, Cayan DR, Maurer EP, Miller NL, Bao Y, Kanamitsu M, Yoshimura K, Snyder MA, Sloan LC (2013) Probabilistic estimates of future changes in California temperature and precipitation using statistical and dynamical downscaling. *Clim Dyn* 40:839–856
- Pierce DW, Kalansky JF, Cayan DR (2018) Climate, Drought, and Sea Level Rise Scenarios for the Fourth California Climate Assessment. California's Fourth Climate Change Assessment, California Energy Commission. Publication number CNRA-CEC-2018-006. 78 p
- Pyšek P, Pergl J, Essl F, Lenzner B, Dawson W, Kreft H, Weigelt P, Winter M, Kartesz J, Nishino M (2017) Naturalized alien flora of the world. *Preslia* 89:203–274
- Pyšek P, Prach K (2003) Research into plant invasions in a crossroads region: history and focus. *Biol Invasions* 5:337–348
- Rapacciuolo G, Maher SP, Schneider AC, Hammond TT, Jabis MD, Walsh RE, Iknayan KJ, Walden GK, Oldfather MF, Ackerly DD (2014) Beyond a warming fingerprint: individualistic biogeographic responses to heterogeneous climate change in California. *Global Change Biol* 20:2841–2855
- R Core Team (2018) R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing
- Reichard S, White P (2001) Horticulture as a pathway of invasive plant introductions in the United States. *BioScience* 51:103–113
- Rehfeldt GE, Crookston NL, Warwell MV, Evans JS (2006) Empirical analyses of plant-climate relationships for the western United States. *Int J Plant Sci* 167:1123–1150e
- Rejmanek M (1996) A theory of seed plant invasiveness: the first sketch. *Biol Conserv* 78:171–181
- Renteria JL, Rouget M, Visser V (2017) Rapid prioritization of alien plants for eradication based on climatic suitability and eradication feasibility. *Austral Ecol* 42:995–1005
- Richardson DM, Bond WJ, Dean WRJ, Higgins SI, Midgley GF, Milton SJ, Powrie LW, Rutherford MC, Samways MJ, Schulze RE (2000) Invasive alien species and global change: a South African perspective. Pages 303–349 in Mooney HA, Hobbs RJ, eds. *Invasive Species in a Changing World*. Washington, DC: Island Press
- Riordan EC, Montalvo AM, Beyers JL (2018) Using Species Distribution Models with Climate Change Scenarios to Aid Ecological Restoration Decisionmaking for Southern California Shrublands. *Res. Rep. PSW-RP-270*. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 130 p
- Roy HE, Rabitsch W, Scalera R, Stewart A, Gallardo B, Genovesi P, Essl F, Adriaens T, Bacher S, Booy O (2018) Developing a framework of minimum standards for the risk assessment of alien species. *J Appl Ecol* 55:526–538
- Rundel PW, Arroyo MTK, Cowling RM, Keeley JE, Lamont BB, Vargas P (2016) Mediterranean biomes: evolution of their vegetation, floras, and climate. *Annu Rev Ecol Evol Syst* 47:383–407
- Sandel B, Dangremond EM (2012) Climate change and the invasion of California by grasses. *Global Change Biol* 18:277–289
- Schmitt S, Pouteau R, Justeau D, de Boissieu F, Birnbaum P (2017) ssdm: an R package to predict distribution of species richness and composition based on stacked species distribution models. *Methods Ecol Evol* 8:1795–1803
- Sofaer HR, Jarnevich CS, Pearse IS, Smyth RL, Auer S, Cook GL, Edwards TC Jr, Guala GF, Howard TG, Morissette JT, Hamilton H (2019) Development and Delivery of Species Distribution Models to Inform Decision-Making. *BioScience* 69:544–557

- Stein BA, Kutner LS, Adams JS (2000) *Precious Heritage: The Status of Biodiversity in the United States*. Oxford: Oxford University Press on Demand
- Stephenson N (1998) Actual evapotranspiration and deficit: Biologically meaningful correlates of vegetation distribution across spatial scales. *J Biogeogr* 25:855–870
- Thuiller W, Richardson DM, Pyšek P, Midgley G, Hughes GO, Rouget M (2005) Niche-based modelling as a tool for predicting the risk of alien plant invasions at a global scale. *Global Change Biol* 11:2234–2250
- van Kleunen M, Essl F, Pergl J, Brundu G, Carboni M, Dullinger S, Early R, González-Moreno P, Groom QJ, Hulme PE, Kueffer C, Kühn I, Máguas C, Maurel N, Novoa A, *et al.* (2018) The changing role of ornamental horticulture in alien plant invasions. *Biol Rev* 93:1421–1437
- Vasquez VL, de Lima AA, dos Santos AP, Pinto MP (2021) Influence of spatial extent on habitat suitability models for primate species of Atlantic For *Ecol Inform* 61:101179
- Venette RC, Kriticos DJ, Magarey RD, Koch FH, Baker RHA, Worner SP, Gómez Raboteaux NN, McKenney DW, Dobesberger EJ, Yemshanov D, De Barro PJ (2010) Pest risk maps for invasive alien species: a roadmap for improvement. *BioScience* 60:349–362
- Verbrugge LNH, de Hoop L, Aukema R, Beringen R, Creemers RCM, van Duinen GA, Hollander H, de Hullu E, Scherpenisse M, Spikmans F, van Turnhout CAM, Wijnhoven S, Leuven RSEW (2019) Lessons learned from rapid environmental risk assessments for prioritization of alien species using expert panels. *J Environ Manage* 249:109405
- Verbruggen H, Tyberghein L, Belton GS, Mineur F, Jueterbock A, Hoarau G, Gurgel CFD, De Clerck O (2013) Improving transferability of introduced species' distribution models: new tools to forecast the spread of a highly invasive seaweed. *PLoS ONE* 8:e68337
- Webber BL, Yates CJ, Le Maitre DC, Scott JK, Kriticos DJ, Ota N, McNeill A, Le Roux JJ, Midgley GF (2011) Modelling horses for novel climate courses: insights from projecting potential distributions of native and alien Australian acacias with correlative and mechanistic models. *Divers Distrib* 17:978–1000
- Zurell D, Franklin J, König C, Bouchet PJ, Dormann CF, Elith J, Fandos G, Feng X, Guillera-Arroita G, Guisan A (2020) A standard protocol for reporting species distribution models. *Ecography* 43:1261–1277