

Susceptibility of native and invasive submersed plants in New Zealand to florpypauxifen-benzyl in growth chamber exposure studies

Authors: Howell, Andrew W., Hofstra, Deborah E., Heilman, Mark A., and Richardson, Robert J.

Source: Invasive Plant Science and Management, 15(3) : 133-140

Published By: Weed Science Society of America

URL: <https://doi.org/10.1017/inp.2022.22>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Research Article

Cite this article: Howell AW, Hofstra DE, Heilman MA, and Richardson RJ (2022) Susceptibility of native and invasive submersed plants in New Zealand to florypyrauxifen-benzyl in growth chamber exposure studies. *Invasive Plant Sci. Manag* **15**: 133–140. doi: [10.1017/inp.2022.22](https://doi.org/10.1017/inp.2022.22)

Received: 9 June 2022

Revised: 20 August 2022

Accepted: 23 August 2022

First published online: 1 September 2022

Associate Editor:

Ryan M. Wersal, Minnesota State University

Keywords:

Arylpicolinates; herbicide concentration; invasive species; oxygen weeds; synthetic auxin

Author for correspondence:

Robert J. Richardson, Department of Crop and Soil Sciences, North Carolina State University, Campus Box 7620, Raleigh, NC 27695. (Email: rob_richardson@ncsu.edu)

Susceptibility of native and invasive submersed plants in New Zealand to florypyrauxifen-benzyl in growth chamber exposure studies

Andrew W. Howell¹ , Deborah E. Hofstra² , Mark A. Heilman³  and Robert J. Richardson⁴ 

¹Graduate Research Assistant, Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC, USA; ²Programme Leader, National Institute of Water and Atmospheric Research, Hillcrest, Hamilton, New Zealand; ³Director of Aquatic Technology, SePRO Corporation, Carmel, IN, USA and ⁴Professor, Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC, USA

Abstract

Invasive aquatic plants constantly threaten freshwaters and associated environs globally. Water resource managers frequently seek new control tactics to combat invasive macrophytes, especially when the availability of herbicides registered for submersed plant control is limited. The synthetic auxin herbicide, florypyrauxifen-benzyl, recently registered (2018) for aquatic site applications in the United States, has shown success in controlling several invasive aquatic weeds. Studies were conducted to evaluate responses of native and invasive submersed plants to florypyrauxifen-benzyl under growth chamber conditions to provide insight on the selectivity of varying herbicide concentrations in New Zealand. Florypyrauxifen-benzyl concentrations evaluated ranged from 0.01 to 107.86 $\mu\text{g ai L}^{-1}$, encompassing the maximum use concentration (48 $\mu\text{g L}^{-1}$) for submersed plant applications. Dose–response metrics indicated the New Zealand native species watermilfoil [*Myriophyllum triphyllum* Orchard] was highly sensitive to florypyrauxifen-benzyl following a 21-d static exposure, having a dry weight 50% effective concentration (EC_{50}) value of 1.2 $\mu\text{g L}^{-1}$. The invasive species oxygen-weed [*Lagarosiphon major* (Ridley) Moss] and Canadian waterweed [*Elodea canadensis* Michx.] were less sensitive, with dry weight EC_{50} values of 35.4 and $>107.86 \mu\text{g L}^{-1}$, respectively. Brazilian waterweed (*Egeria densa* Planch.) was most tolerant to the tested concentrations, as EC_{50} values were not achieved. Overall, results indicate florypyrauxifen-benzyl demonstrates potential for controlling *L. major*, with further large-scale screening required to confirm control among field site applications. As the native species (*M. triphyllum*) was most sensitive to florypyrauxifen-benzyl compared with the invasive plant evaluated (I/N ratio indicated >31.3 times more sensitive), any targeted concentration used for invasive plant control for field applications would likely injure the native *M. triphyllum* plants. Future studies should investigate additional native and invasive species for management guidance and consider how exposure times influence plant response using similar florypyrauxifen-benzyl concentrations tested in the present study.

Introduction

Preservation of native submersed aquatic plants is vital for conserving biodiversity within waterways (Hofstra et al. 2021; Kovalenko et al. 2010), as macrophytes are essential for numerous ecosystem services (Carpenter and Lodge 1986; Cyr and Downing 1988; Madsen et al. 2001; Petr 2000; Valley et al. 2004). Conversely, the intrusion of invasive submersed plants into aquatic biomes can displace endemic flora and fauna through structural and resource competition resulting from the formation of aggressive monotypic stands that limit light, carbon, and nutrient availability (Hofstra et al. 2018; True-Meadows et al. 2016; Wells et al. 1997). Similarly, invasive submersed macrophytes that produce high biomass yielding canopies (e.g., Hydrocharitaceae) commonly obstruct navigation, clog water intakes for irrigation and hydroelectric generation, and impede recreation and economic opportunities (Carpenter and Lodge 1986; Clayton and Champion 2006; Langeland 1996). In New Zealand, some of the most problematic invasive submersed plants are known as the “oxygen weeds” (Clayton 1996), which includes the species oxygen-weed [*Lagarosiphon major* (Ridley) Moss], Canadian waterweed (*Elodea canadensis* Michx.), and Brazilian waterweed (*Egeria densa* Planch.). Water resource managers regularly seek effective control methods to eradicate or manage invasive aquatic plants to enable the recovery of desirable native habitats.

While biological, physical, and mechanical control options do exist, herbicides are largely the most economic, effective, and selective management tool utilized to control invasive aquatic weeds (Hussner et al. 2017; Muller et al. 2021). It is important to recognize that applying herbicides for aquatic weed management requires several factors to be considered, some of which

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



Management Implications

Since their introduction in the early 1900s, invasive submersed plants have adversely affected lakes and waterways on both islands of New Zealand. Some of the worst plant invaders in the country are from the “oxygen weed” (Hydrocharitaceae) family and yield high biomass plant canopies that displace native flora and fauna and obstruct the recreational use and economic activities in affected lakes. Water resource managers regularly utilize herbicides to control aquatic weeds to restore invaded lakes and waterways. However, there are only two aquatic herbicides currently registered in New Zealand for submersed weed control, which limits the scope of management opportunity. Registration of the synthetic auxin herbicide, florypyrauxifen-benzyl, in the United States has provided another chemical option for aquatic weed control. However, limited data are available for florypyrauxifen-benzyl concentrations required to effectively control frequently managed Hydrocharitaceae in New Zealand. A dose–response study was conducted to examine the sensitivity of three New Zealand invasive species [oxygen-weed, *Lagarosiphon major* (Ridley) Moss; Brazilian waterweed, *Egeria densa* Planch.; and Canadian waterweed, *Elodea canadensis* Michx.] and one native submersed plant (watermilfoil, *Myriophyllum triphyllum* Orchard) to the herbicide. Among early invasion scenarios, native plants are frequently found cohabitating with invasive weed species. Therefore, herbicides that provide selective control with minimal impact to native plant species are desired. Following the 21-d growth chamber evaluations, we found the native plant *M. triphyllum* to be the species most sensitive to herbicide. The invasive plant *L. major* was also sensitive to florypyrauxifen-benzyl. The invasive plants *E. canadensis* and *E. densa* displayed a sublethal response from herbicide, and control was not achieved at any florypyrauxifen-benzyl concentration labeled for submersed plant applications ($48 \mu\text{g ai L}^{-1}$). Therefore, targeted concentrations deployed for invasive plant control within mixed communities would likely injure the native *Myriophyllum* spp. However, native species do recover from seedbanks following invasive plant removal. Future research should evaluate additional native and introduced invasive species for best management guidance in New Zealand and investigate approaches, including concentration and exposure time relationships, to provide effective control of submersed aquatic weeds.

include regulatory and economic constraints, herbicide efficacy and selectivity, and public support for the treatment (Champion et al. 2002; Clayton 1996; Madsen 2000; Stallings et al. 2015). Combinations of these factors influence initiatives used to broaden application strategies for restoring native plant populations within invaded waterways (Getsinger et al. 2008; Weber et al. 2020). When considering herbicide-based management strategies, it is critical to understand target and non-target plant responses. Further, these data will help to provide appropriate recommendations for management action in mixed assemblages of native and invasive species, especially because most available herbicides do not provide uniform control of aquatic weeds.

Currently, only two herbicides are labeled for submersed aquatic plant control in New Zealand, diquat-dibromide (WSSA Group 22; photosystem I inhibitor) and endothall dipotassium salt (WSSA Group 31; protein phosphatase inhibitor). For perspective, 16 herbicides are presently labeled for aquatic weed management in the United States (Gettys et al. 2020). A limited herbicide

portfolio restricts management options and prompts selection pressures, which can select for herbicide-resistant plant populations that further complicate future invasive plant control (Richardson 2008). While endothall dipotassium salt and diquat-dibromide effectively control *L. major* (Wells and Champion 2010; Wells et al. 2014), only diquat-dibromide is efficacious on *E. canadensis* and *E. densa* (Glomski et al. 2005; Hofstra and Clayton 2001; Sesin et al. 2018; Skogerboe et al. 2006). Previous studies have examined additional herbicide sites of action (SOAs) including triclopyr (WSSA group 4; synthetic auxin), dichlobenil (WSSA Group 29; cellulose synthase inhibitor), and fluridone (WSSA Group 12; phytoene desaturase inhibitor) to control invasive submersed plants; however, these herbicides are not efficacious on *L. major*, *E. canadensis*, and *E. densa* in mesocosm or field studies (Hofstra and Clayton 2001; Wells et al. 1986), nor are these herbicides registered for aquatic weed control applications in New Zealand (Champion et al. 2019). While endothall dipotassium salt has shown some selectivity on desirable native plants in the United States and New Zealand (Hofstra and Clayton 2001; Skogerboe and Getsinger 2002), diquat-dibromide is a nonselective contact herbicide, and applications are prone to off-target injury to native plants. Wells and Champion (2010) have suggested diquat had transient injury on native charophytes in New Zealand following invasive plant eradication efforts, though recovery was not immediate. Nevertheless, there remains a need to evaluate new herbicides as they become available to enhance current invasive aquatic plant management programs, while protecting native plant species like watermilfoil (*Myriophyllum triphyllum* Orchard), which may be frequently intermixed or adjacent to invasives (Rattray et al. 1994).

Florypyrauxifen-benzyl (WSSA Group 4; synthetic auxin) is a relatively new herbicide initially introduced for rice (*Oryza sativa* L.) production (Epp et al. 2016) and registered for aquatic use in the United States in 2018. Synthetic auxins have been utilized for crop and non-crop weed management since development in the 1940s (Peterson et al. 2016). This class of herbicides is frequently cited for their favorable management characteristics compared with other herbicide SOAs (Glomski and Netherland 2010; Grossmann 2010; Heap 2022; Sprecher et al. 1998). Synthetic auxin herbicides like florypyrauxifen-benzyl are unique in both selectiveness and phloem mobility within susceptible plants, as they mimic the natural plant growth hormone indole-3-acetic acid (Epp et al. 2016). Endogenous auxin compounds are essential for plant cell elongation and division, phototropism, apical dominance, and additional developmental processes (Gaines 2020; Grossmann 2010). Susceptible plants treated with synthetic auxins undergo rapid growth complexes when transcription factor proteins responsible for plant regulation become overwhelmed, triggering uncontrolled gene expression (Grossmann 2010; McCauley et al. 2020; Parry et al. 2009). Eventually, the process of uncontrolled gene expression initiates abscisic acid, hydrogen peroxide, and ethylene accumulation, leading to leaf senescence, cell death, and loss of turgor through multifaceted processes that are still undergoing investigation (Grossmann 2010; McCauley et al. 2020). Synthetic auxin overload in susceptible plants characteristically results in apical epinasty, twisting, and curling of leaf tissues. In the United States, florypyrauxifen-benzyl has provided an additional herbicide for selective invasive aquatic plant management among several common invasive weeds including hydrilla [*Hydrilla verticillata* (L.f.) Royle], watermilfoils (*Myriophyllum* spp.), and crested floatingheart [*Nymphoides cristata* (Roxb.) Kuntze.] while having limited activity on native *Potamogeton*

spp. (Beets and Netherland 2018; Mudge et al. 2021; Netherland and Richardson 2016; Richardson et al. 2016; Sperry et al. 2021). Additionally, florypyrauxifen-benzyl was classified by the U.S. Environmental Protection Agency as a reduced-risk herbicide (USEPA 2017) with favorable toxicological profiles (Buczek et al. 2020).

Given the limited number of registered herbicides in New Zealand, there remains a need to evaluate additional selective herbicides that provide different SOAs than those currently registered. Registration of such a herbicide would promote herbicide stewardship and increase treatment options for controlling invasive submersed plants. The objective of this study was to implement a small-scale screening method for evaluating relative sensitivities to florypyrauxifen-benzyl of native and invasive plant species found in New Zealand using dose–response protocols. Based on previous screening studies (e.g., Beets et al. 2019; Howell et al. 2021; Netherland and Richardson 2016; Richardson et al. 2016), we hypothesize the native species, *M. triphyllum*, will be the most sensitive to the herbicide; however, we anticipate the invasive species tested to display comparable sensitivity to florypyrauxifen-benzyl.

Materials and Methods

A growth chamber experiment was conducted at the National Institute of Water and Atmospheric Research (NIWA) Ruakura Campus, Hamilton, New Zealand in fall (April to May) 2018. The experimental design closely followed a modified version of the Organisation for Economic Co-operation and Development (OECD) protocol described by Netherland and Richardson (2016) to evaluate the sensitivity of *M. triphyllum*, *L. major*, *E. canadensis*, and *E. densa* to florypyrauxifen-benzyl. Tested species were collected from on-site stock tanks or field harvested locally within the Waikato basin in March and April 2018. Plant species were then propagated in aerated outdoor tanks under ambient environmental conditions ($\mu = 18.5$ to 20.0 °C) with 50% shade fabric and monitored twice weekly to ensure adequate population vigor before testing.

At experiment initiation, apical shoot tips (6 cm) of each species were removed from the outdoor tanks. The basal portions of the apical shoots were secured with lead weights to ensure submersion during propagation. Weighted plant shoots were then placed in aerated 6-L bins containing dechlorinated tap water. Bins were subsequently placed in the growth chamber and monitored for 7 to 10 d for confirmation of plant root generation and shoot elongation. Growth chamber conditions were set to a constant 16-h light:8-h dark photoperiod, 21.5 °C temperature, and light intensity of 130 to 160 $\mu\text{E m}^{-2} \text{s}^{-1}$ at bench level. Following rooting confirmation, one shoot of each species was planted in a 20-ml vial filled with 16 cm of washed sand (i.e., a single shoot per vial). At minimum, 3 cm of the shoot was buried in the sand. Seven days before treatment, 1-L jars were filled with 750 ml of Smart and Barko solution, and each jar was provided with supplemental aeration (Smart and Barko 1985). Vials containing plants were then placed in the respective treatment jars. Prior studies noted *Myriophyllum* spp. to be highly sensitive to florypyrauxifen-benzyl (Netherland and Richardson 2016; Richardson et al. 2016). As such, *M. triphyllum* plants and the Hydrocharitaceae evaluated in this study were isolated in separate treatment jars to evaluate herbicide concentration response endpoints based on the recognized sensitivity. Stock solutions of florypyrauxifen-benzyl suspension concentrate (ProcellaCOR SC, SePRO, Carmel, IN, USA) were produced for treatment and injected into the water column to achieve desired

Table 1. Herbicide concentrations used to evaluate species sensitivity to florypyrauxifen-benzyl in the growth chamber study.

Species	Family (division)	New Zealand plant status	Evaluated concentrations $\mu\text{g ai L}^{-1}$
<i>Myriophyllum triphyllum</i>	Haloragaceae (dicotyledon)	Native	0, 0.01, 0.04, 0.13, 0.4, 1.33, 4, 11.98, 35.96, 107.86
<i>Lagarosiphon major</i>	Hydrocharitaceae (monocotyledon)	Invasive	0, 0.13, 0.4, 1.33, 4, 11.98, 35.96, 66.58, 107.86
<i>Elodea canadensis</i>		Invasive	
<i>Egeria densa</i>		Invasive	

serial dose–response concentrations. Treatments consisted of static exposure to a geometric series of rates ranging from 0 to 107.86 $\mu\text{g ai L}^{-1}$ (Table 1). Pretreatment water pH was 8.2 (SD ± 0.2) and temperature was 22.2 °C (SD ± 1.4). All treatments were replicated five times (one jar was considered as one replication) following a randomized complete block design, and experiments were repeated in time (two consecutive runs). At treatment, five nontreated jars were removed to determine pretreatment weight and shoot length of each species. Trials lasted 21 d, and dechlorinated water was added to the jars as water loss occurred. Visual observations of auxin herbicide symptoms (e.g., chlorosis, epinasty, leaf-shattering) were documented throughout experimentation. At 21 d after treatment (DAT), above-sediment green plant tissue was harvested and blotted dry with paper towels, and then fresh weight (g) and shoot length (cm) were immediately recorded. Plants were then oven-dried at 60 °C for 72 h to obtain a constant dry weight. Plant dry weights were measured using an analytical-grade balance with 0.001 g accuracy.

Data Analysis

There was no significant run effect according to ANOVA ($P > 0.05$), so treatment data were pooled over experiments to account for inherent response variability in the growth chamber studies. Plant shoot length, fresh biomass, and dry biomass metrics were transformed to percent inhibition (%In) of the nontreated control to standardize plant response to tested florypyrauxifen-benzyl concentrations tested using Equation 1:

$$\%In = [(\mu_c - \mu_t) / \mu_c] \times 100 \quad [1]$$

Where (μ_c) is the mean of the nontreated group and (μ_t) is the mean value of the treatment group. Percent inhibition (%In) was limited to the logical extremes (0% to 100%) to achieve appropriate parameters for modeling plant response to herbicide concentrations tested, as plant inhibition cannot physically exceed a 100% threshold. Following similar statistical procedures as Netherland and Richardson (2016), nonlinear regression analyses were performed in SigmaPlot v. 14.0 (SigmaPlot v. 14.0.3.192, Systat Software, Point Richmond, CA, USA) to estimate 50% effective dose concentrations (EC_{50}). Equation 2 is the four-parameter log-logistic regression curve used to estimate EC_{50} shoot length and dry weight inhibition metrics at tested florypyrauxifen-benzyl concentrations as described in detail by Ritz et al. (2015) and Seefeldt et al. (1995).

Table 2. Metrics of 50% effective concentrations (EC₅₀) derived from log-logistic four-parameter or Weibull four-parameter dose–response models following plant exposure to florypyrauxifen-benzyl at 0 to 107.86 µg ai L^{–1}, the lowest observed effect concentration (LOEC) calculated from Dunnett’s test at the 0.05 significance level, and invasive-to-native plant tolerance index.

Species	EC ₅₀ florypyrauxifen-benzyl metrics ^a					I/N ratio ^b
	Shoot length inhibition	Fresh weight inhibition	Dry weight inhibition	Dry weight LOEC		
		µg ai L ^{–1}				
<i>Myriophyllum triphyllum</i>	4.7	5.0	1.2	0.4		—
<i>Lagarosiphon major</i>	na	49.5	35.4	35.96		31.3
<i>Elodea canadensis</i>	na	> 107.86	> 107.86	> 107.87		93.8
<i>Egeria densa</i>	na	na	na	na		na

^aEC₅₀ values were not achieved (na) if species exhibited low inhibition (µ < 50%) for the metric tested.
^bEstimated invasive species (Hydrocharitaceae; I) dry weight EC₅₀ value divided by the native species (*M. triphyllum*; N) dry weight EC₅₀ value.

$$Y = y_o + \{a/[1 + (x/x_{EC50})^b]\}$$
 [2]

For Equation 2, the parameters y_o and a represent the limit extreme and difference values, b is the slope of the inflection point, x is the herbicide concentration, and x_{EC50} is the herbicide concentration providing 50% inhibition of the maximum (i.e., 100% In). Equation 3 is the Weibull four-parameter model used to estimate EC₅₀ fresh weight inhibition metrics at tested florypyrauxifen-benzyl concentrations as described in detail by Price et al. (2012) and Brown and Mayer (1988).

$$Y = a \times \{1 - \exp[-(x - x_{EC50} + b + \ln(2)^{1/c})/b]^c\}$$
 [3]

For Equation 3, the parameter a is the upper asymptote, b is the slope of the inflection point, c is the shape of the curve, x is the herbicide concentration, and x_{EC50} is the herbicide concentration providing 50% inhibition of the maximum (i.e., 100% In). The Weibull model is suitable when asymmetric data, like plant fresh weight, define a response variable at a different rate than could be described using a log-logistic curve (Price et al. 2012). Selected models were chosen when they converged across the applicable dataset(s) and when the Shapiro-Wilks normality assumptions were met ($\alpha = 0.05$). For each species by metric modeled, lack-of-fit tests were performed to ensure the selected model was appropriate. Graphical outputs from the models used log-transformed values of tested florypyrauxifen-benzyl concentrations for each species response.

A Dunnett’s test ($P < 0.05$) was performed in RStudio (R v. 4.0.3, R Foundation for Statistical Computing, Vienna, Austria) to establish the lowest observed effect concentration (LOEC) between the nontreated and treated plant dry biomass at the select florypyrauxifen-benzyl concentrations using the BASE and MULTCOMP packages (Horthorn et al. 2008; R Core Team 2022). An index comparing the estimated herbicide tolerance of invasive plants to the estimated herbicide susceptibility of the native plant (I/N ratio) was also calculated. The I/N ratio was defined as the estimated dry weight EC₅₀ value of invasive plant species (Hydrocharitaceae; I) divided by the corresponding native species (*M. triphyllum*; N) dry weight EC₅₀ value (EC₅₀ invasive species/EC₅₀ native species).

Results and Discussion

Nontreated reference plants exhibited shoot elongation and axillary branching during the 21-d static exposure. Nontreated plant biomass increased by 2.5 to 9.5 times that of the pretreatment biomass, which conforms to experiment validation standards of the OECD protocol (OECD 2014). The tested florypyrauxifen-benzyl

concentrations evaluated in this study ranged 0.02% to 225% of the commonly recommended formulated product maximum use rate (48 µg L^{–1}) for submersed plant applications; therefore, plant control was compared directly using predicted EC₅₀ values of shoot length, fresh weight, and dry weight metrics.

The native species, *M. triphyllum*, was the most sensitive plant evaluated in this study, with a dry weight EC₅₀ value of 1.2 µg L^{–1} and LOEC of 0.4 µg L^{–1} (Table 2; Figure 1). Within 1 to 2 DAT, *M. triphyllum* exhibited auxin herbicide exposure symptoms, with epinasty appearing as the first sign of plant injury (data not shown). At the 2 wk after treatment (WAT) evaluation, plant symptoms had progressed to necrotic shoots with black nodes at concentrations >1.33 µg L^{–1}. Following plant harvest, *M. triphyllum* treated with >11.98 µg L^{–1} had <20% biomass remaining, and shoot lengths were reduced by 65% relative to the nontreated plants (Figure 1). These results confirm our hypothesis that *M. triphyllum* is highly sensitive to florypyrauxifen-benzyl, as the EC₅₀ values in this study align well with previous small-scale herbicide-screening observations among other *Myriophyllum* spp. (Netherland and Richardson 2016; Richardson et al. 2016).

The most sensitive invasive species tested was *L. major*, which had an estimated dry weight EC₅₀ value of 35.4 µg L^{–1} and LOEC of 35.96 µg L^{–1} (Table 2; Figure 1). Nevertheless, an EC₅₀ for shoot length was not achieved (shoot inhibition µ < 50%). *Lagarosiphon major* injury from florypyrauxifen-benzyl was also rapid, with proximal bending and minor chlorosis observed within 1 to 3 DAT (data not shown). Leaf abscission from the apical shoots ensued ca. 7 DAT, when plants were gently agitated using forceps, with apical shoots fragmenting at concentrations >11.98 µg L^{–1} 2 WAT. Howell et al. (2021) noted a similar response to herbicide in an outdoor mesocosm study in which *L. major* exposed to florypyrauxifen-benzyl at 30 and 50 µg ai L^{–1} displayed proximal leaf abscission and stem fragmentation at 5 to 7 DAT. At harvest, biomass of *L. major* shoots treated with concentrations >66.58 µg L^{–1} was reduced more than 62% compared with the nontreated plants. Though sensitive, the calculated I/N ratio suggests *L. major* would require 31.3 times more herbicide to produce EC₅₀ values similar to that of the native species, *M. triphyllum* (Table 2).

Elodea canadensis was not as sensitive to florypyrauxifen-benzyl as *L. major*, with fresh and dry weight EC₅₀ value estimates greater than the highest concentration tested of 107.86 µg L^{–1}. Consequently, the LOEC for *E. canadensis* was also >107.86 µg L^{–1}. Shoot length inhibition did not meet the criteria for estimating an EC₅₀ at any tested concentration (shoot inhibition µ < 50%) (Table 2; Figure 1). While response metrics suggest low sensitivity in this study, plant injury and growth abnormalities were present. Shoots became brittle, with necrotic tissue forming at the nodes




Figure 1. Native (*Myriophyllum triphyllum*) and invasive (*Lagarosiphon major*, *Elodea canadensis*, and *Egeria densa*) plant responses following 21-d static exposure to tested florypyrauxifen-benzyl serial concentrations expressed as percent inhibition of the nontreated control: (A) shoot inhibition, (B) fresh weight inhibition, and (C) dry weight inhibition. Data points with standard error bars represent mean response of the metric evaluated. Herbicide concentration is provided on a log₁₀ scale. Regression analyses implemented for plant shoot and dry weight inhibition correspond to the log-logistic four-parameter model equation: $Y = y_0 + \{a/[1 + (x/x_{EC50})^b]\}$, while fresh weight inhibition was modeled using the Weibull four-parameter equation: $Y = a \times \{1 - \exp[-(x - x_{EC50} + b + \ln(2)^{1/c})/b]^c]\}$.

with exposures to concentrations $>11.98 \mu\text{g L}^{-1}$ at 2 to 3 WAT (data not shown). Conversely, a slight increase in axillary branching was noted at harvest among several plants treated at $\leq 11.98 \mu\text{g}$

L^{-1} . Based on the I/N ratio, *E. canadensis* would require >90 times more herbicide to produce an EC_{50} value comparable to that of *M. triphyllum* (Table 2). Similarly, Beets et al. (2019) noticed *E.*

canadensis biomass was not affected when testing florypyrauxifen-benzyl at 3 to 27 $\mu\text{g ai L}^{-1}$ in a 60-d concentration and exposure (CET) study.

Of the Hydrocharitaceae species evaluated, *E. densa* was the least sensitive, with EC_{50} and LOEC values not achieved with any test concentrations (Table 2). The trend in the data appeared linear with a shallow slope, indicating limited plant response to increased florypyrauxifen-benzyl concentrations during the 21-d exposure (Figure 1). Much like *E. canadensis*, apical portions of treated plants displayed auxin herbicide exposure characteristics, with slight epinasty, shoot twisting, and internode lengthening observed at 2 WAT at concentrations $\geq 35.96 \mu\text{g L}^{-1}$ (data not shown). However, these abnormal growth patterns did not appreciably reduce biomass or shoot lengths collected at harvest (Table 2). Howell et al. (2021) noted 80% *E. densa* visual control was not achieved until 7 WAT with florypyrauxifen-benzyl at 30 $\mu\text{g L}^{-1}$ in outdoor mesocosm studies. Likewise, Haug et al. (2021) indicated *E. densa* had less shoot absorption and translocation than the Hydrocharitaceae species *H. verticillata* in a ^{14}C experiment applying 10 $\mu\text{g ai L}^{-1}$ florypyrauxifen-benzyl during a 192-h exposure period. These previous studies suggest longer static exposure periods (e.g., >4 wk) may improve control of *E. densa*, as indicated by the low sensitivity shown in the present study, especially at lower concentrations (e.g., <36.96 $\mu\text{g L}^{-1}$). Consistent with this hypothesis, Madsen et al. (2021) showed *E. densa* dry biomass was reduced by approximately 50% compared with the nontreated plants following a 10-wk static exposure to 50 $\mu\text{g ai L}^{-1}$ florypyrauxifen-benzyl.

Elodea canadensis and *E. densa* proliferation in axillary branching and shoot development at the lower treatment concentrations ($\leq 11.98 \mu\text{g L}^{-1}$) further conveys the typical synthetic auxin properties of florypyrauxifen-benzyl, despite limited overall efficacy (personal observation). Hormesis, or augmented growth following sublethal herbicide concentrations, is characteristic of low-dose auxin injury (Belz and Duke 2014; Cedergreen et al. 2007; Jalal et al. 2021). Hormesis was noted in a previous study that documented a stimulated increase in yield for *E. densa* treated with the auxin herbicide, 2,4-D, applied at 1 to 11 mg ai L^{-1} (Peres et al. 2017). Similarly, Mudge et al. (2021) suggested potential hormesis occurred for *E. canadensis* in a 6-wk CET study when exposed to florypyrauxifen-benzyl at 3, 6, and 9 $\mu\text{g ai L}^{-1}$. While macrophyte hormesis literature is limited for florypyrauxifen-benzyl, findings from these previous auxin herbicide screenings closely align with the observations of *E. canadensis* and *E. densa* response to treatment in the present study. Further, these data denote the perceptive effective dose thresholding, which can occur among auxin herbicides, and the varying sensitivity found even within the same plant family (e.g., *L. major* dry weight EC_{50} was ~3-fold less than *E. canadensis* in the present study). Further research is required to specifically evaluate the lower florypyrauxifen-benzyl threshold concentrations and exposures that deter possible hormesis in common field application scenarios; notably in high water-exchange situations (e.g., flowing systems).

Though complexities of induced hormesis do exist with auxin herbicides (Belgers et al. 2007; Peres et al. 2017), submersed plant tolerance and sensitivity to synthetic auxins is well documented (Getsinger et al. 2003; Haug and Bellaud 2013; Hofstra and Clayton 2001; Parsons et al. 2001; Richardson et al. 2016; Sperry et al. 2021; Wersal et al. 2010). Unlike nonselective herbicides like diquat-dibromide, synthetic auxin herbicides, like 2,4-D and triclopyr largely act as selective compounds, which typically do not adversely affect monocotyledons compared with dicotyledon

(broadleaf species) counterparts (Gettys et al. 2020; Madsen 2000). However, as part of the arylpicolinate class of auxins, florypyrauxifen-benzyl associates with a binding-site receptor atypical of common predecessor auxin classes (e.g., 2,4-D belongs within the phenoxy-carboxylate class) (Epp et al. 2016; Hoyerova et al. 2018; Lee et al. 2014). The mobility of florypyrauxifen-benzyl acid metabolites (Haug et al. 2021) and subsequent auxin derivatives within susceptible aquatic plants denotes the unique activity levels of this herbicide, as several selectivity phenomena were evident in the present study. For example, Hofstra and Clayton (2001) noted that *M. triphyllum* was not controlled in greenhouse trials using triclopyr at 0.25 to 2.5 mg L^{-1} . Yet rapid sensitivity and plant death was observed for *M. triphyllum* at very low florypyrauxifen-benzyl concentrations ($\leq 1.2 \mu\text{g ai L}^{-1}$). In the same trial, triclopyr did not provide adequate control of *L. major* in New Zealand (Hofstra and Clayton 2001), while *L. major* showed rapid sensitivity with no signs of recovery at florypyrauxifen-benzyl concentrations >35 $\mu\text{g ai L}^{-1}$ in the present study.

While literature documenting submersed plant control with florypyrauxifen-benzyl is still developing at the international scale, results from the present study corroborate the findings of *Myriophyllum* spp. sensitivity among previous studies (Beets et al. 2019; Haug et al. 2021; Richardson et al. 2016). However, our findings do contradict those originally found by Netherland and Richardson (2016), which indicated greater sensitivity for *E. canadensis* with EC_{50} values of 6.9 and 13.1 $\mu\text{g ai L}^{-1}$, as *E. canadensis* EC_{50} required more than twice the maximum labeled concentration of formulated herbicide in this 21-d study. Florypyrauxifen-benzyl degrades primarily through photolysis (1- to 2-d half-life), with secondary degradation occurring through hydrolysis with increasing pH (pH 7 to 9; 111- to 2-d half-life, respectively) (Heilman and Getsinger 2018; MDA 2018). Though unlikely, treatment pH in this study ($\mu = 8.2$) could have nominally influenced herbicide activity on *E. canadensis*. A rapid conversion of florypyrauxifen-benzyl to the less-active parent acid under growth chamber conditions could have also reduced the observed herbicide activity on the more tolerant invasive plants, although this was not specifically tested for (Netherland and Richardson 2016; Richardson et al. 2016; Sperry et al. 2021). A more likely explanation for greater *E. canadensis* sensitivity in prior research is the potential for genotypic differences between naturalized plant populations in New Zealand versus the native range cohorts in North America. In New Zealand, *E. canadensis* accessions occur solely via clonal propagation, whereas viable seed production can occur within various regions in North America (Swearingen and Barger 2016). Past genetic comparisons were performed within established Hydrocharitaceae populations in New Zealand (Lambertini et al. 2010); however, this type of study typically focuses on species plasticity and invasion potential within invaded waterways rather than genetic parallels to the endemic plant populations. Further genetic screening comparing test species in the United States and New Zealand would allow for more relevant plant response comparisons for confirmation of this hypothesis.

In conclusion, this study indicates florypyrauxifen-benzyl would be a prospective candidate for *L. major* management, with further evaluation required to develop tactics that produce adequate control levels for *E. canadensis* and *E. densa*. Large-scale mesocosm trials would be beneficial in elucidating *E. canadensis* whole-plant response to validate plant tolerance metrics shown in this study, as previous outdoor mesocosm experiments showed more favorable results for *E. densa* control (Howell et al. 2021). Given the

sensitivity of *M. triphyllum* compared with the Hydrocharitaceae tested, any targeted concentration used for invasive plant control in field scenarios would likely seriously injure native *Myriophyllum* spp. However, resource managers should note that native species, such as *M. triphyllum* and red pondweed (*Potamogeton cheesemaniae* A. Benn.), generally produce large seedbanks that could support reestablishment following invasive plant eradication programs (Hofstra et al. 2018; de Winton and Clayton 1996; de Winton et al. 2000). As documented in previous studies evaluating florypyrauxifen-benzyl for invasive submersed plant control (Beets et al 2019; Richardson et al. 2016; Sperry et al. 2021), future research is needed to test additional native species' sensitivity for best management guidance. Similarly, future investigations should assess native and invasive species in mixed communities to show side-by-side confirmation of results in field-based plant management scenarios. While small-scale screenings can overestimate herbicide activity on plants (Richardson et al. 2016), this study exemplifies the benefits of alternative small-scale trials for quickly gauging submersed plant response to new herbicide chemistries and provides a foundation for future screening activity that could prove expedient for herbicide registration purposes.

Acknowledgments. The authors would like to thank Denise Rendle and Susie Elcock for their assistance with plant collection, propagation, and harvesting. Funding for NIWA trials provided through the Strategic Science Investment Fund. SePRO Corporation provided florypyrauxifen-benzyl material for testing. The authors recognize that the coauthor Mark Heilman is employed by SePRO Corporation. The remaining authors declare no conflicts of interest.

References

- Beets J, Heilman M, Netherland MD (2019) Large-scale mesocosm evaluation of florypyrauxifen-benzyl, a novel arylpicolinate herbicide, on Eurasian and hybrid watermilfoil and seven native submersed plants. *J Aquat Plant Manag* 57:49–55
- Beets J, Netherland MD (2018) Mesocosm response of crested floating heart, hydrilla, and two native emergent plants to florypyrauxifen-benzyl. *J Aquat Plant Manag* 56:57–62
- Belgers JDM, Van Lieverloo RJ, Van der Pas LJT, Van de Brink PJ (2007) Effects of the herbicide 2,4-D on the growth of nine aquatic macrophytes. *Aquat Bot* 86:260–268
- Belz RG, Duke SO (2014) Herbicides and plant hormesis. *Pest Manag Sci* 70:698–707
- Brown RF, Mayer DG (1988) Representing cumulative germination. 2. The use of the Weibull function and other empirically derived curves. *Ann Bot* 61:127–138
- Buczek SB, Archambault JM, Cope WG, Heilman MA (2020) Evaluation of juvenile freshwater mussel sensitivity to multiple forms of florypyrauxifen-benzyl. *Bull Environ Contam Toxicol* 105:588–594
- Carpenter SR, Lodge DM (1986) Effects of submersed macrophytes on ecosystem processes. *Aquat Bot* 26:341–370
- Cedergreen N, Streibig JC, Kudsk P, Mathiassen SK, Duke SO (2007) The occurrence of hormesis in plants and algae. *Dose Response* 5:150–162
- Champion P, Clayton J, Rowe D (2002) *Lake Manager's Handbook: Alien Invaders*. Wellington, New Zealand: Ministry for the Environment. 49 p
- Champion P, Hofstra D, de Winton M (2019) *Best Management Practice for Aquatic Weed Control, Part One: The Framework*. Hamilton, New Zealand: National Institute of Water & Atmospheric Research Ltd. Client Report. 82 p
- Clayton J, Champion P (2006) Risk assessment method for submerged weeds in New Zealand hydroelectric lakes. *Hydrobiologia* 570:183–188
- Clayton JS (1996) Aquatic weeds and their control in New Zealand lakes. *Lake Reserv Manag* 12:477–486
- Cyr H, Downing JA (1988) The abundance of phytophilous invertebrates on different species of submerged macrophytes. *Freshw Biol* 20:365–374
- Epp JB, Alexander AL, Balko TW, Buysse AM, Brewster WK, Bryan K, Daeuble JF, Fields SC, Gast RE, Green RA, Irvine NM, Lo WC, Lowe CT, Renga JM, Richburg JS, et al. (2016) The discovery of Arylex™ active and Rinskor™ active: two novel auxin herbicides. *Bioorg Med Chem* 24:362–371
- Gaines T (2020) The quick and the dead: a new model for the essential role of ABA accumulation in synthetic auxin herbicide mode of action. *J Exp Bot* 71:3383–3385
- Getsinger K, Sprecher S, Smagula A (2003) Effects of triclopyr on variable-leaf watermilfoil. *J Aquat Plant Manag* 41:124–126
- Getsinger KD, Netherland MD, Grue CE, Koschnick TJ (2008) Improvements in the use of aquatic herbicides and establishment of future research directions. *J Aquat Plant Manag* 46:32–41
- Gettys LA, Haller WT, Petty D, eds (2020) *Biology and Control of Aquatic Plants*. 4th ed. Marietta, GA: Aquatic Ecosystem Restoration Foundation. 237 p
- Glomski L, Skogerboe J, Getsinger K (2005) Comparative efficacy of diquat to two members of the Hydrocharitaceae family: elodea and hydrilla. *J Aquat Plant Manag* 43:103–105
- Glomski LM, Netherland MD (2010) Response of Eurasian and hybrid watermilfoil to low use rates and extended exposures of 2,4-D and triclopyr. *J Aquat Plant Manag* 48:12–16
- Grossmann K (2010) Auxin herbicides: current status of mechanism and mode of action. *Pest Manag Sci* 66:113–120
- Haug E, Ahmed K, Gannon T, Richardson R (2021) Absorption and translocation of florypyrauxifen-benzyl in ten aquatic plant species. *Weed Sci* 69: 624–630
- Haug EJ, Bellaud MD (2013) Efficacy of 2,4-D ester on variable-leaf milfoil control for partial lake treatments in New Hampshire waterbodies. *J Aquat Plant Manag* 51:49–52
- Heap I (2022) The International Herbicide-Resistant Weed Database. <http://www.weedscience.org>. Accessed: February 23, 2022
- Heilman MA, Getsinger K (2018) Early Operational Demonstration of Selective Invasive Watermilfoil Control with ProcellaCOR. https://bugwoodcloud.org/mura/mipn/assets/File/UMISC-2018/Wednesday/PM/Heilman_Early_Operational_Demo_ProcellaCOR.pdf. Accessed: April 18, 2022
- Hofstra D, Clayton J, Champion P, de Winton M (2018) Control of invasive aquatic plants. Pages 267–298 in Hamilton D, Collier K, Quinn J, Howard-Williams C, eds. *Lake Restoration Handbook*. Cham, Switzerland: Springer
- Hofstra D, Clements D, Rendle D, Champion P (2021) Evaluation of flumioxazin on seven submersed macrophytes in New Zealand. *J Aquat Plant Manag* 5:27–34
- Hofstra DE, Clayton JS (2001) Evaluation of selected herbicides for the control of exotic submerged weeds in New Zealand: I. The use of endothall, triclopyr and dichlobenil. *J Aquat Plant Manag* 39:20–24
- Horthorn T, Bretz F, Westfall P (2008) Simultaneous inference in general parametric models. *Biom J* 50:346–363
- Howell AW, Hofstra DE, Heilman MA, Richardson RJ (2021) Evaluation of florypyrauxifen-benzyl to control three problematic submersed macrophytes in New Zealand. *J Aquat Plant Manag* 59:66–71
- Hoyerova K, Hosek P, Quareshy M, Li J, Klima P, Kubes M, Yemm AA, Neve P, Tripathi A, Bennett MJ, Napier RM (2018) Auxin molecular field maps define AUX1 selectivity: many auxin herbicides are not substrates. *New Phytol* 217:1625–1639
- Hussner A, Stiers I, Verhofstad MJJM, Bakker ES, Grutters BMC, Haury J, van Valkenburg JLCH, Brundu G, Newman J, Clayton JS, Anderson LWJ, Hofstra D (2017) Management and control methods of invasive alien freshwater aquatic plants: a review. *Aquat Bot* 136:112–137
- Jalal A, de Oliveira J Jr, Ribeiro J, Fernandes G, Mariano G, Trindade V, dos Reis A (2021) Hormesis in plants: physiological and biochemical responses. *Ecotoxicol Environ Saf* 207:1–12
- Kovalenko KE, Dibble ED, Slade J (2010) Community effects if invasive macrophyte control: role of invasive plant abundance and habitat complexity. *J Appl Ecol* 47:318–328
- Lambertini C, Riis T, Olesen B, Clayton JS, Sorrell BK, Brix H (2010) Genetic diversity in three invasive clonal aquatic species in New Zealand. *BMC Genet* 11:1–18

- Langeland KA (1996) *Hydrilla verticillata* (LF) Royle (Hydrocharitaceae), “the perfect aquatic weed.” *Castanea* 61:293–304
- Lee S, Sundaram S, Armitage L, Evans JP, Hawkes T, Kepinski S, Ferro N, Napier RM (2014) Defining binding efficiency and specificity of auxins for SCF(TIR1/AFB)-Aux/IAA co-receptor complex formation. *ACS Chem Biol* 9:673–682
- Madsen JD (2000) Advantages and Disadvantages of Aquatic Plant Management Techniques. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 31 p
- Madsen JD, Chambers PA, James WF, Koch EW, Westlake DF (2001) The interaction between water movement, sediment dynamics and submersed macrophytes. *Hydrobiologia* 444:71–84
- Madsen JD, Morgan C, Miskella J, Kyser G, Gilbert P, O’Brien J, Getsinger KD (2021) Brazilian egeria herbicide mesocosm and field trials for managing the Sacramento–San Joaquin River Delta. *J Aquat Plant Manag* 59s:90–97
- McCauley CL, McAdam SAM, Bhide K, Thimmapuram J, Banks JA, Young BG (2020) Transcriptomics in *Erigeron canadensis* reveals rapid photosynthetic and hormonal responses to auxin herbicide application. *J Exp Bot* 71:3701–3709
- [MDA] Minnesota Department of Agriculture (2018) New Active Ingredient Review: Florypyrauxifen-Benzyl. <https://www.mda.state.mn.us/sites/default/files/inline-files/Florypyrauxifen-benzyl.pdf> Accessed: April 18, 2022
- Mudge CR, Sartain BT, Sperry BP, Getsinger KD (2021) Efficacy of Florypyrauxifen-Benzyl for Eurasian Watermilfoil Control and Nontarget Illinois Pondweed, Elodea, and Coontail Response. ERDC/TN APCRP CC-24. Vicksburg, MS: U.S. Army Engineer Research and Development Center. 7 p
- Muller C, Hofstra D, Champion P (2021) Eradication economics for invasive alien aquatic plants. *Manag Biol Invasions* 12:253–271
- Netherland MD, Richardson RJ (2016) Evaluating sensitivity of five aquatic plants to a novel arylpicolinate herbicide utilizing an organization for economic cooperation and development protocol. *Weed Sci* 64:181–190
- [OECD] Organisation for Economic Co-operation and Development (2014) Test No. 239: Water-Sediment *Myriophyllum spicatum* Toxicity Test, OECD Guidelines for the Testing of Chemicals. Section 2. Paris: OECD Publishing. 10.1787/9789264224155-en
- Parry G, Calderon-Villalobos LI, Prigge M, Peret B, Dharmasiri S, Itoh H, Lechner E, Gray WM, Bennett M, Estelle M (2009) Complex regulation of the TIR1/AFB family of auxin receptors. *PNAS USA* 106:22540–22545
- Parsons JK, Hamel KS, Madsen JD, Getsinger KD (2001) The use of 2,4-D for selective control of an early infestation of Eurasian watermilfoil in Loon Lake, Washington. *J Aquat Plant Manag* 39:117–125
- Peres LRS, Della Vechia JF, Cruz (2017) Hormesis effect of herbicides subdoses on submerged macrophytes in microassay conditions. *Planta Daninha* 35, 10.1590/s0100-83582017350100076
- Peterson MA, McMaster SA, Riechers DE, Skelton J, Stahlman PW (2016) 2,4-D past, present, and future: a review. *Weed Technol* 30:303–345
- Petr T (2000) Interactions between Fish and Aquatic Macrophytes in Inland Waters: A Review. FAO Fisheries Technical Paper 396. Rome, Italy: FAO. 185 p
- Price WJ, Shafii B, Seefeldt SS (2012) Estimation of dose–response models for discrete and continuous data in weed science. *Weed Technol* 26:587–601
- Ratray MR, Howard-Williams C, Brown JMA (1994) Rates of early growth of propagules of *Lagarosiphon major* and *Myriophyllum triphyllum* in lakes of differing trophic status. *NZ J Mar Freshw Res* 28:235–241
- R Core Team (2022) R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org>.
- Richardson RJ (2008) Aquatic plant management and the impact of emerging herbicide resistance issues. *Weed Technol* 22:8–15
- Richardson RJ, Haug EJ, Netherland MD (2016) Response of seven aquatic plants to a new arylpicolinate herbicide. *J Aquat Plant Manag* 54:26–31
- Ritz C, Kniss AR, Streibig JC (2015) Research methods in weed science: statistics. *Weed Sci* 63:166–187
- Seefeldt SS, Jensen JE, Fuerst EP (1995) Log-logistic analysis of herbicide dose–response relationships. *Weed Technol* 9:218–227
- Sesin V, Dalton RL, Boutin C, Robinson SA, Bartlett AJ, Pick FR (2018) Macrophytes are highly sensitive to the herbicide diquat dibromide in test systems of varying complexity. *Ecotoxicol Environ Saf* 165:325–333
- Skogerboe JG, Getsinger KD (2002) Endothall species selectivity evaluation: northern latitude aquatic plant community. *J Aquat Plant Manag* 40:1–5
- Skogerboe JG, Getsinger KD, Glomski LAM (2006) Efficacy of diquat on submersed plants treated under simulated flowing water conditions. *J Aquat Plant Manag* 44:122–125
- Smart RM, Barko JW (1985) Laboratory culture of submersed freshwater macrophytes on natural sediments. *Aquat Bot* 21:251–263
- Sperry BP, Leary JK, Jones KD, Ferrell JA (2021) Observations of a submersed field application of florypyrauxifen-benzyl suppressing hydrilla in a small lake in central Florida. *J Aquat Plant Manag* 29:20–26
- Sprecher SL, Getsinger KD, Stewart AB (1998) Selective effects of aquatic herbicides on sago pondweed. *J Aquat Plant Manag* 36:64–68
- Stallings KD, Seth-Carley D, Richardson RJ (2015) Management of aquatic vegetation in the southeastern United States. *J Integr Pest Manag* 6:1–5
- Swearingen J, Barger C (2016) Invasive Plant Atlas of the United States. Tifton: University of Georgia Center for Invasive Species and Ecosystem Health. <http://www.invasiveplantatlas.org>. Accessed: February 21, 2022
- True-Meadows S, Haug EJ, Richardson RJ (2016) Monoecious hydrilla—a review of the literature. *J Aquat Plant Manag* 54:1–11
- [USEPA] U.S. Environmental Protection Agency (2017) Final Registration Decision on the New Active Ingredient Florypyrauxifen-Benzyl. Washington, DC: U.S. Environmental Protection Agency. 21 p
- Valley RD, Cross TK, Radomski P (2004) The Role of Submersed Aquatic Vegetation as Habitat for Fish in Minnesota Lakes, Including the Implications of Non-native Plant Invasions and Their Management. Special Publication 160. Saint Paul, MN: Minnesota Department of Natural Resources. 25 p
- Weber MA, Wainger LA, Harms NE, Nesslage GM (2020) The economic value of research in managing invasive hydrilla in Florida public lakes. *Lake Reserv Manag* 37:63–76
- Wells RDS, Champion PD (2010) Endothall for aquatic weed control in New Zealand. Pages 307–310 in *Proceedings of the 17th Australian Weeds Conference*, Christchurch, New Zealand, September 26–30, 2010
- Wells RDS, Champion PD, Clayton JS (2014) Potential for lake restoration using the aquatic herbicide endothall. Science, community and food security: the weed challenge. Pages 1–4 in *Proceedings of the 19th Australasian Weeds Conference*, Hobart, Australia, September 1–4, 2014
- Wells RDS, Coffey BT, Lauren DR (1986) Evaluation of fluridone for weed control in New Zealand. *J Aquat Plant Manag* 24:39–42
- Wells RDS, de Winton M, Clayton JS (1997) Successive macrophyte invasions within the submerged flora of Lake Tarawera, Central North Island, New Zealand. *NZ J Mar Freshw Res* 31:449–459
- Wersal RM, Madsen JD, Woolf TE, Eckberg N (2010) Assessment of herbicide efficacy on Eurasian watermilfoil and impacts to the native submersed plant community in Hayden Lake, Idaho, USA. *J Aquat Plant Manag* 48:5–11
- Winton MD de, Clayton JS (1996) The impact of invasive submerged weed species on seed banks in lake sediments. *Aquat Bot* 53:31–45
- Winton M de, Clayton J, Champion P (2000) Seedling emergence from seed banks of 15 New Zealand lakes with contrasting vegetation histories. *Aquat Bot* 66:181–194