

Evidence of large sturgeons in the Paleocene of North America

Author: Brownstein, Chase Doran

Source: Journal of Paleontology, 97(1) : 218-222

Published By: The Paleontological Society

URL: <https://doi.org/10.1017/jpa.2022.87>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Evidence of large sturgeons in the Paleocene of North America

Chase Doran Brownstein* 

Department of Ecology and Evolutionary Biology, Yale University, New Haven, Connecticut 06520, and Stamford Museum and Nature Center, Stamford, Connecticut 06903, USA <chase.brownstein@yale.edu>

Abstract.—Ray-finned fishes comprise nearly half of extant vertebrate species and include several ancient lineages with fossil records that stretch over 200 Myr in time. One of these old clades, the sturgeons and paddlefishes, is distributed across the Northern Hemisphere and includes some of the largest known freshwater fishes. Yet, the fossil record of this lineage (Acipenseriformes) is poor compared to similarly ancient ray-finned fish clades. Here, I describe sturgeon fossils from two geological units in North America < 10 Myr younger than the Cretaceous–Paleogene boundary. Both come from individuals of ~1.5 m in length. These Paleogene forms establish the long history of large body size in Acipenseriformes and reveal sturgeons were some of the largest inhabitants of freshwater ecosystems that were still recovering from the Cretaceous–Paleogene extinction.

Introduction

North America is a hotspot of ancient aquatic vertebrate diversity. All but one of the classic ancient lineages of ray-finned fishes (Polypteridae; e.g., Near et al., 2014) are found on the continent, and at least three of these are presently entirely confined to North America (gars, Bowfin, *Hiodon* Lesueur, 1818; Grande and Bemis, 1998; Hilton, 2003; Hilton and Grande, 2008; Grande, 2010). Acipenseriformes, which currently comprises 27 species of sturgeons and the Paddlefish and includes some of the largest living freshwater fishes (*Huso* Linnaeus, 1758; see Berg, 1948) is the most diverse and geographically widespread of the major ancient ray-finned fish clades. Despite the status of this clade, the evolutionary history of Acipenseriformes remains poorly understood. The anatomy, fossil record, and interrelationships of this clade have been the subject of numerous detailed morphological studies (Grande and Bemis, 1991; Bemis et al., 1997; Grande and Hilton, 2006; Hilton and Grande, 2006; Hilton et al., 2011; Sato et al., 2018; Murray et al., 2020; During et al., 2022).

Despite the large number of specialized features unique to Paddlefish and sturgeons, the available fossil record for these fishes indicates the phenotype in Acipenseriformes has remained remarkably unchanged over perhaps 100 Myr (Hilton et al., 2011; Sato et al., 2018; Murray et al., 2020). Unfortunately, gaps in the fossil record of Acipenseriformes for key intervals like the aftermath of the Cretaceous–Paleogene mass extinction 66 Ma have obscured whether this pattern of phenotypic conservatism is a biological reality or an artifact of sampling (Grande and Hilton, 2006; Hilton and Grande, 2006; Hilton et al., 2011; Sato et al., 2018; Hilton and Grande, 2022).

Here, I describe two records of sturgeons from the 7 Myr following the Cretaceous–Paleogene mass extinction in western North America (Fig. 1). These lateral osteoderms demonstrate the presence of large (>1.5 m) sturgeons in early-middle Paleocene freshwater ecosystems. In turn, they extend the record of large (>1 m) body size in Acipenseriformes back by 60 Myr and imply that the high degree of body size disparity observed in this clade relative to other vertebrates (Rabosky et al., 2013) is a consistent pattern in their evolutionary history.

Materials

Repository and institutional abbreviation.—YPM, Yale Peabody Museum, New Haven, Connecticut, USA.

Systematic paleontology

Actinopterygii Cope, 1887
Acipenseriformes Berg, 1940
Acipenseridae Bonaparte, 1831
Acipenseridae morphotype A
Figures 2, 3

Occurrence.—Layer #3, Eagle Mine near Bear Creek, Carbon County, Montana (Bear Creek Local Fauna, Fort Union Formation, Tiffanian–Clarkfordian, 61.2–56.6 Ma; Barnosky et al., 2014, fig. 1). Precise coordinates are available to qualified researchers at YPM.

Description.—YPM VPPU 17066 consists of an exquisitely preserved, isolated dermal scute (Fig. 2.1–2.3) assignable to a large acipenserid sturgeon (Hilton and Grande, 2006; Hilton et al., 2011; Thieren et al., 2015; Sato et al., 2018; Murray

*Corresponding author.




Figure 1. Locality and horizon of the new sturgeon material: (1) map of the United States showing localities where sturgeon records reported in this paper were recovered; (2) simplified stratigraphic column of the Cretaceous-Paleogene in Montana showing horizons from which sturgeon material reported in this paper was recovered.

et al., 2020). The size and shape of the scute allow assignment of YPM VPPU 17066 to the sturgeon crown group Acipenseridae because the closest fossil relatives of this lineage lack large lateral scutes (e.g., Hilton and Grande, 2006). The scute, which represents the best record of an acipenseriform from the Bear Creek Local Fauna, comes from the middle portion of the lateral row based on its dorsoventral asymmetry and

median ridge size (Hilton et al., 2011), which produce a slight posterior slant for the scute in lateral and medial views (Fig. 2.1, 2.2).

The external surface is ornamented with unorganized deep pitting and several large ridges that radiate from the defined central ridge and produce a rugose outline. These ridges are much larger than those seen in extant sturgeons, e.g., *Acipenser* spp. (e.g., Hilton et al., 2011; Thieren et al., 2015). Pitting is not confined to the center of the external surface of the scute and variously reaches the borders of this bone (Fig. 2.1, 2.2). Size estimation using the lateral scute maximum equation of Thieren and Van Neer (2016) using the dorsoventral maximum height of YPM 17066 (70 mm) provides a total length of 1.650 m for YPM VPPU 17066 (Fig. 3).

Material.—YPM VPPU 17066, a complete lateral scute.

Remarks.—YPM VPPU 17066 is identifiable as the lateral dermal scute of an acipenserid.

Acipenseridae morphotype B

Figures 2, 3

Occurrence.—Highway Blowout Site, Fallon County, Montana (Tongue River Member, Fort Union Formation, Tiffanian-




Figure 2. Anatomy of the new sturgeon specimens of Acipenseridae morphotype A: (1–3) YPM VPPU 17066, Bear Creek Site, Montana: (1) lateral view; (2) medial view; (3) anterior view; (4–6) YPM VPPU 16646, Highway Blowout Site, Montana: (4) lateral view; (5) medial view; (6) dorsal view.

Clarkfordian, ~62–60 Ma; Peppe et al., 2011). Precise coordinates are available to qualified researchers at YPM.

Description.—YPM VPPU 16646 is another complete dermal scute from a large acipenserid fish (Fig. 2.4–2.6). This specimen is slightly larger than YPM VPPU 17066 and shows a very different set of features. Although the dorsoventral asymmetry of YPM VPPU 16646 demonstrates that it comes from the lateral scute row, the specimen is strongly angled along its dorsoventral midline (Fig. 2.4). This produces distinctive dorsal and ventral faces. The form of ornamentation on YPM VPPU 16646 is also markedly different from YPM VPPU 17066 and more closely resembles the ornamentation of extant sturgeons like *Acipenser* spp., *Huso* spp., *Scaphirhynchus* spp., and *Pseudoscaphirhynchus* spp. (Hilton et al., 2011; Thieren et al., 2015) and extinct crown-group acipenserids, e.g., †*Anchiacipenser acanthaspis* Sato et al., 2018. Although the size of individual pits on YPM VPPU 16646 is larger than most crown group sturgeons (Hilton et al., 2011; Thieren et al., 2015; Sato et al., 2018) and stem-group taxa from North America (Hilton et al., 2011; Sato et al., 2018), YPM VPPU 16646 shares with these forms

reduced radiating ridges and pitting ornamentation that is widely separated from the scute border by a relatively smooth surface bearing small radiating ridges (Fig. 2.4–2.6).

YPM VPPU 16646 is unusual in that it bears a large flange directed toward the posterior end of the body. Although some extant sturgeon lateral scutes show small posterior spines (Thieren et al., 2015), the posterior flange on YPM VPPU 16646 is two-thirds of the anteroposterior length of the main body of the osteoderm. Another odd feature of YPM VPPU 16646 is the rounded outline of the main body of the scute. In most crown sturgeons, the lateral scutes have main bodies that terminate dorsally and ventrally at distinct apices, forming a rhomboid shape in lateral and medial views. Size estimation using the lateral scute maximum equation of Thieren and Van Neer (2016) using the dorsoventral maximum height of YPM VPPU 16646 (= 75 mm) gives a total length of 1.653 m for the Highway Blowout acipenserid (Fig. 3).

Material.—YPM VPPU 16646, a complete lateral scute.

Remarks.—YPM VPPU 16646 is identifiable as the lateral dermal scute of an acipenserid.




Figure 3. Size comparison of the two new sturgeon records of Acipenseridae morphotype A: Bear Creek sturgeon (left); Highway Blowout sturgeon (right). Silhouette is a 1.85 m tall man.

Discussion

Eight extant sturgeon species are currently found in North America, but the fossil record of the crown group on the continent stretches back to the Cretaceous (Hilton and Grande, 2006; Hilton et al., 2011; Sato et al., 2018; Murray et al., 2020). Although living sturgeons in the genera *Acipenser* Linnaeus, 1758 and *Huso* are the largest known freshwater fishes (e.g., Berg, 1948; Hilton et al., 2011), most of the published fossil record of the crown clade is comprised by relatively small (~1.0 m or less) species (Hilton and Grande, 2006; Sato et al., 2018; Murray et al., 2020). The ancient ages of the two lineages in the crown group are known to include giant species. For example, various time-calibrated molecular phylogenies suggest that giant species, e.g., *Acipenser transmontanus* Richardson, 1836 and *Huso* spp., last share common ancestry with other sturgeons varying between 8 and >50 Ma (Peng et al., 2007; Luo et al., 2019; Shen et al., 2020). These estimated ages imply long ghost lineages leading to extant giant sturgeons.

The two new sturgeon specimens, although not complete enough to justify naming new species, represent giant acipenserids from two different geological units in the Paleocene of the American West. Both specimens are within the upper size bracket for Acipenseridae (Thieren et al., 2015; Thieren and Van Neer, 2016), stretching the record of large size in this clade back by at least 60 Myr to the recovery period following the Cretaceous-Paleogene mass extinction. Although their assignment to subclades in Acipenseridae is limited by the wide variability of osteoderm shape in sturgeons (Hilton and Grande, 2006; Hilton et al., 2011; Sato et al., 2018; Murray et al., 2020), both the Bear Creek and Highway Blowout scutes belong to pan-acipenserids, which are characterized by the presence of distinctive dermal osteoderms (Hilton and Grande, 2006; Hilton et al., 2011; Sato et al., 2018; Murray et al., 2020).

The high degree of body size disparity in extant sturgeons (Rabosky et al., 2013) contrasts with their status as a living fossil lineage, i.e., a long-lived, species-poor clade that shows little phenotypic change from ancient fossil relatives (Gardiner, 1984; Grande and Bemis, 1991; Bemis et al., 1997; Hilton et al., 2011). A similar pattern of proportionally high variation in body size relative to species diversity has also been documented in latimeroid coelacanths, which include the freshwater-brackish clade †Mawsoniidae (Cavin et al., 2021).

Finally, the new large-bodied sturgeon records add to the growing body of evidence that ancient freshwater fish assemblages from the Paleocene and Eocene of North America contained numerous large-bodied clades with different body plans (e.g., Grande and Bemis, 1991, 1998; Grande, 2010; Hilton et al., 2011; Brownstein, 2022). The suction-feeding style of sturgeons (e.g., Carroll and Wainwright, 2003) represents a unique component to these assemblages, which are already known to include durophagous and macropredatory gars with shortened to longirostrine skulls (Grande and Bemis, 1998; Grande, 2010; Brownstein, 2022), filter-feeding paddlefishes (Grande and Bemis, 1998), and large, predatory osteoglossiform fishes with the characteristic rasping device that gives the clade its name (Forey and Hilton, 2010; Hilton and Lavoué, 2018). In turn, the specimens described in this contribution indicate a

legacy of giant acipenserids in Paleocene North American ecosystems in need of further discoveries to be fully revealed.

Acknowledgments

I thank D. Brinkman, G. Watkins-Colwell, and V. Rhue for access to the ichthyology and palaeoichthyology collections of YPM. This research was supported by the Yale Undergraduate Richter Summer Fellowship through Pierson College. I thank the editor A. Murray, D. Brinkman, and an anonymous reviewer for their comments, which greatly improved this manuscript. Sturgeon silhouettes used in Figure 3 are in the public domain (<http://phylopic.org/>). This paper is dedicated to my parents D.I. Brownstein and L.R. Tannebaum, who first introduced me to sturgeons as the tastiest smoked fish.

References

- Barnosky, A.D., Holmes, M., Kirchholti, R., Lindsey, E., Maguire, K.C., Poust, A.W., Stegner, M.A., Sunseri, J., Swartz, B., Swift, J., and Villavicencio, N.A., 2014, Prelude to the Anthropocene: Two new North American land mammal ages (NALMAs): The Anthropocene Review, v. 1, no. 3, p. 225–242, <https://doi.org/10.1177/2053019614547433>.
- Bemis, W.E., Findeis, E.K., and Grande, L., 1997, An overview of Acipenseriformes: Environmental Biology of Fishes, v. 48, no. 1, p. 25–71.
- Berg, L.S., 1940, Classification of Fishes Both Recent and Fossil, edition with English translation: Bangkok, Thai National Document Center, 517 p.
- Berg, L.S., 1948, Freshwater Fishes of the USSR and Adjacent Countries, Volume 2: Jerusalem, Israel Program for Scientific Translations, 496 p.
- Bonaparte, C.L., 1831, Saggio di una distribuzione metodico degli animali vertebrati: Giornale Arcadico di Scienze, Lettere ad Arti, v. 49, p. 1–77.
- Brownstein, C.D., 2022, Unappreciated Cenozoic ecomorphological diversification of stem gars revealed by a new large species: Acta Palaeontologica Polonica, v. 67, no. 3, p. 559–568, <https://doi.org/10.4202/app.00957.2021>.
- Carroll, A.M., and Wainwright, P.C., 2003, Functional morphology of prey capture in the sturgeon, *Scaphirhynchus albus*: Journal of Morphology, v. 256, no. 3, p. 270–284, <https://doi.org/10.1002/jmor.10095>.
- Cavin, L., Piuze, A., Ferrante, C., and Guinot, G., 2021, Giant Mesozoic coelacanths (Osteichthyes, Actinistia) reveal high body size disparity decoupled from taxic diversity: Scientific Reports, v. 11, no. 1, p. 1–13, <https://doi.org/10.1038/s41598-021-90962-5>.
- Cope, E.D., 1887, Zittel's Manual of Palaeontology: The American Naturalist, v. 21, p. 1014–1019.
- During, M.A.D., Smit, J., Voeten, D.F.A.E., Berruyer, C., Tafforeau, P., Sanchez, S., Sten, K.H.W., Verdegaaal-Warmerdam, S.J.A., and van der Lubbe, J.H.J.L., 2022, The Mesozoic terminated in boreal spring: Nature, v. 603, p. 91–94, <https://doi.org/10.1038/s41586-022-04446-1>.
- Forey, P.L., and Hilton, E.J., 2010, Two new Tertiary osteoglossid fishes (Teleostei: Osteoglossomorpha) with notes on the history of the family, in Elliot, D., Yu, X., Maisey, J., and Miao, D., eds. Fossil Fishes and Related Biota: Morphology, Phylogeny and Paleobiogeography—In Honor of Meemann Chang: Munich, Germany, Friedrich Pfeil, p. 215–246.
- Gardiner, B.G., 1984, Sturgeons as living fossils, in Eldredge, N., and Stanley, S.M., eds., Living Fossils: New York, Springer, p. 148–152.
- Grande, L., 2010, An empirical synthetic pattern study of gars (Lepisosteiformes) and closely related species, based mostly on skeletal anatomy: The resurrection of Holostei: Copeia, v. 10, no. 2A, p. 1–871.
- Grande, L., and Bemis, W.E., 1991, Osteology and phylogenetic relationships of fossil and recent paddlefishes (Polyodontidae) with comments on the interrelationships of Acipenseriformes: Journal of Vertebrate Paleontology, v. 11, no. S1, p. 1–121.
- Grande, L., and Bemis, W.E., 1998, A comprehensive phylogenetic study of amiid fishes (Amiidae) based on comparative skeletal anatomy: An empirical search for interconnected patterns of natural history: Journal of Vertebrate Paleontology, v. 18, suppl. 1, p. 1–696.
- Grande, L., and Hilton, E.J., 2006, An exquisitely preserved skeleton representing a primitive sturgeon from the Upper Cretaceous Judith River Formation of Montana (Acipenseriformes: Acipenseridae: n. gen. and sp.): Journal of Paleontology, v. 80, no. S65, p. 1–39, <https://doi.org/10.1666/05032.1>.
- Hilton, E.J., 2003, Comparative osteology and phylogenetic systematics of fossil and living bony-tongue fishes (Actinopterygii, Teleostei, Osteoglossomorpha): Zoological Journal of the Linnean Society, v. 137, no. 1, p. 1–100, <https://doi.org/10.1046/j.1096-3642.2003.00032.x>.

- Hilton, E.J., and Grande, L., 2006, Review of the fossil record of sturgeons, family Acipenseridae (Actinopterygii: Acipenseriformes), from North America: *Journal of Paleontology*, v. 80, no. 4, p. 672–683, [https://doi.org/10.1666/0022-3360\(2006\)80\[672:ROTFROJ\]2.0.CO;2](https://doi.org/10.1666/0022-3360(2006)80[672:ROTFROJ]2.0.CO;2).
- Hilton, E.J., and Grande, L., 2008, Fossil moonies (Teleostei: Hiodontiformes, Hiodontidae) from the Eocene of western North America, with a reassessment of their taxonomy: *Geological Society, London, Special Publications*, v. 295, no. 1, p. 221–251, <https://doi.org/10.1144/SP295.13>.
- Hilton, E.J., and Lavoué, S., 2018, A review of the systematic biology of fossil and living bony-tongue fishes, Osteoglossomorpha (Actinopterygii: Teleostei): *Neotropical Ichthyology*, v. 16, e180031, <https://doi.org/10.1590/1982-0224-20180031>.
- Hilton, E.J., Grande, L., and Bemis, W.E., 2011, Skeletal anatomy of the short-nose sturgeon, *Acipenser brevirostrum* Lesueur, 1818, and the systematics of sturgeons (Acipenseriformes, Acipenseridae): *Fieldiana Life and Earth Sciences*, no. 3, 168 p.
- Hilton, E., and Grande, L., 2022, Late Cretaceous sturgeons (Acipenseridae) from North America, with two new species from the Tanis site in the Hell Creek Formation of North Dakota: *Journal of Paleontology*, 1–29, <https://doi.org/10.1017/jpa.2022.81>.
- Lesueur, C.A., 1818, Description of several new species of North American fishes: *Journal of the Academy of Natural Sciences of Philadelphia*, v. 1, ser. 1, p. 222–235.
- Linnaeus, C., 1758, *Systema Naturae per Regna Tria Naturae* (tenth edition), Volume 1, *Regnum Animale*: Stockholm, Laurentii Salvii, 824 p.
- Luo, D., Li, Y., Zhao, Q., Zhao, L., Ludwig, A., and Peng, Z., 2019, Highly resolved phylogenetic relationships within order Acipenseriformes according to novel nuclear markers: *Genes*, v. 10, no. 1, 38, <https://doi.org/10.3390/genes10010038>.
- Murray, A.M., Brinkman, D.B., DeMar, D.G. Jr., and Wilson, G.P., 2020, Paddlefish and sturgeon (Chondrostei: Acipenseriformes: Polyodontidae and Acipenseridae) from lower Paleocene deposits of Montana, USA: *Journal of Vertebrate Paleontology*, v. 40, no. 2, e1775091, <https://doi.org/10.1080/02724634.2020.1775091>.
- Near, T.J., Dornburg, A., Tokita, M., Suzuki, D., Brandley, M.C., and Friedman, M., 2014, Boom and bust: Ancient and recent diversification in bichirs (Polypteridae: Actinopterygii), a relictual lineage of ray-finned fishes: *Evolution*, v. 68, no. 4, p. 1014–1026, <https://doi.org/10.1111/evo1.2323>.
- Peng, Z., Ludwig, A., Wang, D., Diogo, R., Wei, Q., and He, S., 2007, Age and biogeography of major clades in sturgeons and paddlefishes (Pisces: Acipenseriformes): *Molecular Phylogenetics and Evolution*, v. 42, no. 3, p. 854–862, <https://doi.org/10.1016/j.ympev.2006.09.008>.
- Peppe, D.J., Johnson, K.R., and Evans, D.A., 2011, Magnetostratigraphy of the Lebo and Tongue River Members of the Fort Union Formation (Paleocene) in the northeastern Powder River Basin, Montana: *American Journal of Science*, v. 311, no. 10, p. 813–850, <https://doi.org/10.2475/10.2011.01>.
- Rabosky, D.L., Santini, F., Eastman, J., Smith, S.A., Sidlauskas, B., Chang, J., and Alfaro, M.E., 2013, Rates of speciation and morphological evolution are correlated across the largest vertebrate radiation: *Nature Communications*, v. 4, no. 1, 1958, <https://doi.org/10.1038/ncomms2958>.
- Richardson, J., 1836, *Fauna Boreali-Americana, or the Zoology of the Northern Parts of British America, containing descriptions of the objects of natural history collected on the late northern land expeditions under command of Captain Sir John Franklin, R.N., Part 3, The Fish*: London, Richard Betley, 327 p., pls. 74–97.
- Sato, H., Murray, A.M., Vernygora, O., and Currie, P.J., 2018, A rare, articulated sturgeon (Chondrostei: Acipenseriformes) from the Upper Cretaceous of Dinosaur Provincial Park, Alberta, Canada: *Journal of Vertebrate Paleontology*, v. 38, no. 4, p. 1–15, <https://doi.org/10.1080/02724634.2018.1488137>.
- Shen, Y., Yang, N., Liu, Z., Chen, Q., and Li, Y., 2020, Phylogenetic perspective on the relationships and evolutionary history of the Acipenseriformes: *Genomics*, v. 112, no. 5, p. 3511–3517, <https://doi.org/10.1016/j.ygeno.2020.02.017>.
- Thiery, E., and Van Neer, W., 2016, New equations for the size reconstruction of sturgeon from isolated cranial and pectoral girdle bones: *International Journal of Osteoarchaeology*, v. 26, no. 2, p. 203–210, <https://doi.org/10.1002/oa.2407>.
- Thiery, E., Wouters, W., and Van Neer, W., 2015, Guide for the identification of archaeological sea sturgeon (*Acipenser sturio* and *A. oxyrinchus*) remains: *Cybium*, v. 39, no. 3, p. 175–192.

Accepted: 12 September 2022