

The assemblage of birds struck by aircraft differs among nearby airports in the same bioregion

Authors: Steele, William K., and Weston, Michael A.

Source: Wildlife Research, 48(5) : 422-425

Published By: CSIRO Publishing

URL: <https://doi.org/10.1071/WR20127>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

The assemblage of birds struck by aircraft differs among nearby airports in the same bioregion

William K. Steele^A and Michael A. Weston^{B,C} 

^AEcology Balance, PO Box 2513, Ringwood North, Vic. 3134, Australia.

^BDeakin University, Geelong, Australia, School of Life and Environmental Sciences, Faculty of Science, Engineering and the Built Environment, Melbourne Campus, 221 Burwood Highway, Burwood, Vic., 3125, Australia.

^CCorresponding author. Email: mweston@deakin.edu.au

Abstract

Context. Bird–aircraft collisions impose an economic cost and safety risk, yet ecological studies that inform bird hazard management are few, and to date no study has formally compared species’ strike profiles across airports. In response to strike risks, airports have implemented customised management on an airport-by-airport basis, based on the assumption that strike risk stems from prevailing local circumstances. We tested this assumption by comparing a decade of wildlife–aircraft strikes at three airports situated in the same bioregion (likely to have similar fauna) of Victoria, Australia.

Aim. To compare the assemblage of wildlife struck by aircraft at three major airports in the same bioregion.

Method. Standardised wildlife strike data were analysed from three airports (Avalon, Melbourne and Essendon Airports), in the Victorian Volcanic Plains bioregion, central Victoria, Australia. Ten discrete 1-year sampling periods from each airport were compared, spanning the period 2009–19. Bird data were comparable, and data on mammals were considered less reliable, so emphasis was placed on birds in the present study.

Results. In total, 580 bird strikes were analysed, with the most commonly struck species being Australian magpie (*Cracticus tibicen*; 16.7%), Eurasian skylark (*Alauda arvensis*; 12.2%), Australian pipit (*Anthus australis*; 12.1%), masked lapwing (*Vanellus miles*; 5.9%), nankeen kestrel (*Falco cenchroides*; 5.0%), house sparrow (*Passer domesticus*; 4.8%), welcome swallow (*Hirundo neoxena*; 4.3%) and tree martin (*Petrochelidon nigricans*; 4.0%). The assemblage of birds struck by aircraft over the decade of study differed between airports. The most commonly struck species drove the assemblage differences between airports.

Conclusions and implications. In the present study system, airports experienced discrete strike risk profiles, even though they are in the same bioregion. The airports examined differed in terms of air traffic movement rates, aircraft types, landscape context and bird hazard management effort. Given that strike risks profiles differ among airports, customised management at each airport, as is currently the case, is supported.

Keywords: aerodrome, aircraft, airfield, collision, community, mammals.

Received 27 July 2020, accepted 13 January 2021, published online 18 March 2021

Introduction

Although aircraft strikes on wildlife are not common events, they present a real hazard to aircraft and exact a significant cost in terms of both human lives and money (Cleary and Dolbeer 2005; El-Sayed 2019). The problem of mammal– and bird–aircraft collisions is likely to become greater in the future as the volume of air traffic increases (DeVault *et al.* 2013). Continued exacerbation of the problem seems likely – it has been suggested that modern commercial aircraft, which are quieter and have larger engine air intakes than older models, are involved in proportionately more bird strikes than older aircraft because birds are less able to detect them in time to avoid collisions (Chilvers *et al.* 1997; Ministerie van Verkeer en Waterstaat 1999). Furthermore, there are concerns, particularly in North America,

that populations of high strike-risk species of wetland bird are increasing (North American Bird Conservation Initiative, US Committee 2014).

Although commercial aircraft generally fly too high to be at risk of colliding with birds, many military and light aircraft utilise the same air space as birds, and all aircraft are exposed to the risk of a bird strike when landing or taking off at airports. The threat of bird strikes in the vicinity of an airport is increased because some bird species congregate at these sites (Burger 1983; E&SS 1994; Ministerie van Verkeer en Waterstaat 1999; Transport Canada 2004; DeVault *et al.* 2013; El-Sayed 2019). Thus, the management of wildlife at airports – particularly bird populations – to reduce the risk of wildlife strikes is becoming increasingly important to airport operators, including those in

Australia (ATSB 2012, 2019a, 2019b). Such efforts generally occur on an aerodrome-by-aerodrome basis, with management plans and management implementation occurring usually by aerodrome-specific operational staff, or by consultants contracted by specific aerodromes (Bunker and Jones 2008). An underlying assumption of this is that aerodromes differ in their circumstances and risks, and therefore require customised management responses. The Australian Transport Safety Bureau (and its predecessor, the Bureau of Air Safety Investigation) has maintained records of bird–aircraft collisions for Australia aerodromes and seeks to share best-practice wildlife management practices (BASI 1996; ATSB 2003, 2008, 2010, 2012, 2014, 2017, 2019a, 2019b).

Australia has 10 aerodromes classified as ‘major class C aerodromes’ and 12 classified as ‘regional towered’ (ATSB 2019a, 2019b). Three of these 22 larger aerodromes are located to the north and west of Melbourne, Victoria: (1) Melbourne International Airport is a major aerodrome; and (2) Essendon Fields Airport and (3) Avalon Airport are both regional towered aerodromes. All three aerodromes lie within the Victorian Volcanic Plain Bioregion of Victoria (DELWP 2021), but support different numbers and types of aircraft. The local pool of species can influence wildlife species occurring near airports (Alquezar *et al.* 2020). Assuming the wildlife communities at these three aerodromes were similar by virtue of being in the same bioregion, we investigated whether the bird species struck by aircraft at each aerodrome were also similar. We therefore explore how generalisable bird–aircraft strikes are in a biogeographically matched set of airports, and examine the assumption that bird hazard mitigation is best implemented at the scale of the aerodrome.

Methods

Three large aerodromes lying within the Victorian Volcanic Plains Bioregion of Australia were included in this study. Melbourne Airport (IATA code: MEL; 37°40′24″S, 144°50′36″E) is the second busiest in Australia; most aircraft movements are regular passenger transport (RPT) flights, or large passenger jet aircraft. Essendon Fields Airport (IATA code: MEB; 37°43′41″S, 144°54′07″E) lies less than 10 km from the Melbourne Airport runway intersection. This aerodrome supports emergency services’ flights, private and corporate aircraft, and flying schools, so has fewer Air Traffic Movements (ATMs; take-offs, landings and touch-and-go) than Melbourne Airport, and does not see the large jets. Avalon Airport (IATA code: AVV; 38°02′26″S, 144°28′15″E) is situated near Little River, and operates some RPT flights, with maintenance and training facilities for large jets. Every second year (odd-numbered years), this aerodrome hosts the Avalon International Airshow, which sees a huge increase in air traffic for one week.

Data on bird strikes were obtained from the Australian Transport Safety Bureau’s National Aviation Occurrence Database (ATSB 2019a, 2019b). All records for the 10-year period between 1 July 2009 and 30 June 2019 for occurrence types ‘Bird strike’ and ‘Wildlife – other’ within 25 km of the three airports were obtained and attributed to the appropriate aerodrome. WKS was the wildlife hazards management consultant at two of the three aerodromes and vetted all reports. Any bias in

detectability of species struck (a hypothetical effect, which has not been demonstrated), would likely exist among aerodromes and so not confound our comparisons.

Monthly ATM figures for the three aerodromes, Melbourne, Essendon Field and Avalon, were obtained from the Air Services Australia website (Airservices Australia 2019) and summed to show the total number of ATMs per financial year: 2009–10 to 2018–19. All three airports had substantive aircraft movements, although the number of movements differed among airports (repeated-measures generalised linear model on logged data, $F_{1,181, 10,630} = 1934.295$, $P < 0.001$; means \pm s.e. number of movements annually: MEB 225 643 \pm 5523; MEL 53 692 \pm 351; AVV, 7504 \pm 523).

Statistical analyses

We analysed the assemblages of birds struck by aircraft using multivariate analyses implemented in PRIMER (v. 7). Airport was specified as a factor, and 10 replicate 12-month sampling periods, from July to June inclusive, were available for each airport; these periods were the same across all airports and spanned 2009–19. We present analyses of the frequency of strikes of each species (number of strikes within the given period), but analyses of the presence or absence of a species being struck in a given period revealed qualitatively identical results and are therefore not presented. Analyses of strike rate (species strike rate per ATM) also revealed differences between airports, but rates were extremely low and interpretation proved difficult. Resemblance matrices were based on zero-inflated Bray–Curtis similarity measures, and these were visualised using non-metric Multidimensional Scaling (MDS). Differences were tested using Analysis of Similarities (ANOSIM), and where differences existed these were explored using the test of homogeneity in Permutational Dispersions (PERMDISP) and Similarity Percentages Analysis (SIMPER; contributions to dissimilarities of $\geq 10\%$ are presented).

Results

Unidentified birds were removed from the dataset, leaving 580 bird strikes (54 avian taxa). The most commonly struck species were Australian magpie (*Cracticus tibicen*; 16.7%), Eurasian skylark (*Alauda arvensis*; 12.2%), Australian pipit (*Anthus australis*; 12.1%), masked lapwing (*Vanellus miles*; 5.9%), nankeen kestrel (*Falco cenchroides*; 5.0%), house sparrow (*Passer domesticus*; 4.8%), welcome swallow (*Hirundo neoxena*; 4.3%) and tree martin (*Petrochelidon nigricans*; 4.0%). The assemblage of birds struck by aircraft differed among airports (ANOSIM, $r = 0.801$, $P = 0.001$), with pairwise tests revealing each airport had a different assemblage being struck (MEL v. AVV, $r = 0.865$, $P = 0.001$; MEL v. MEB, $r = 0.988$, $P = 0.001$; AVV v. MEB, $r = 0.530$, $P = 0.001$) (Fig. 1). Differences in multivariate dispersions were not evident (PERMDISP, $F_{1,27} = 1.539$, $P = 0.307$), indicating that differences lay only in the location of assemblage compositions in multivariate space. We used SIMPER to explore these differences, and identified which species drove pairwise differences among airports, i.e. contributed $\geq 10\%$ of the observed dissimilarity (also, see the vectors plotted on Fig. 1). More Eurasian skylark, Australian magpie and Australian pipit were struck at




Fig. 1. Non-metric multidimensional scaling plot of birds struck at each airport (two-dimensional stress, 0.13). Vectors indicate correlations ≥ 0.7 . MEL, Melbourne Airport; AVV, Avalon Airport; MEB, Essendon Fields Airport.

MEL than at AVV or MEB. Australian magpie, masked lapwing and house sparrow were struck more frequently at MEB than at AVV. Nankeen kestrel, tree martin and welcome swallow (but no other species) all contributed 5–10% of the observed dissimilarity among airports.

Our data on mammal strikes included three bats (grey-headed flying fox, *Pteropus poliocephalus*; Gould's wattled bat, *Chalinolobus gouldii*; and white-striped freetail bat, *Tadarida australis*) and four terrestrial species (short-beaked echidna, *Tachyglossus aculeatus*; European rabbit, *Oryctolagus cuniculus*; European hare, *Lepus capensis*; and red fox, *Vulpes vulpes*) – 85 strikes of identified species in total. Mammal strikes were less comparable than bird strikes among airports (airport operations staff focus on bird hazards) but we note the assemblage differed among airports (excluding echidna, which were infrequently struck; ANOSIM, $r = 0.548$, $P = 0.001$), although pairwise tests revealed that the assemblage struck did not differ between MEB and AVV ($r = 0.067$, $P = 0.230$). All of the airports are fenced but the areas enclosed vary markedly, with Melbourne Airport covering a substantially greater area than the other two and able to support intractable populations of some mammal species.

Discussion

With one exception (Pfeiffer *et al.* 2018), formal multi-aerodrome comparisons of species struck by aircraft have not previously been performed, although Soldatini *et al.* (2011) consider different animal guilds struck by aircraft at eight Italian airports in deriving their risk assessment model. In the USA, positive correlations in strike rates of species among distant airports were evident (Fernández-Juricic *et al.* 2018), and an analysis of 98 civil airports revealed that higher surrounding landscape heterogeneity (especially the proximity of farmland and water) was associated with higher strike rates with aircraft (Pfeiffer *et al.* 2018). Here we showed that three Australian aerodromes in the same bioregion experience different assemblages of birds struck. This reinforces the current approaches to

bird hazard management for aircraft, which are implemented on an aerodrome-by-aerodrome basis (Dolbeer *et al.* 1993). However, some similarities among airports within the studied bioregion are evident. The main species struck were insectivorous or generalist, aerial or ground foragers; this contrasts with airports in northern and Western Australia, where raptors and waterfowl are commonly struck (ATSB 2019a, 2019b).

The underlying causes of the variation in the assemblage of birds struck between airports are unknown, but could involve factors associated with species and/or those associated with airports. Species distributions and associations with habitat have led to bioregions being the basis of large-scale study of bird occurrence (Barrett *et al.* 2003), but finer-scale variation in species and habitat distributions also occur. Thus, within bioregions, birds may respond to landscape- and local-scale factors (Callaghan *et al.* 2018) such that some are present near or transit through particular aerodromes, whereas others are not. For coastal biomes, the influence of adjacent waters also likely influences the occurrence of some species (Alquezar *et al.* 2020; but see Callaghan *et al.* 2018). Although Avalon airport is near-coastal, differences persisted between the other two landlocked aerodromes we examined. Airports themselves, with their associated light and noise, may effectively 'filter' the local pool of species that venture onto the aerodromes, with some species being 'airport adapters' and some being 'airport avoiders' (Alquezar *et al.* 2020). For example, Australian magpies on airfields appear to habituate to aircraft (Linley *et al.* 2018). Other airport factors involve aircraft type and movement rates, local management of bird hazards (including local habitat management), size and seasonal and temporal patterns in air traffic (Cleary and Dolbeer 2005).

Conclusion

The biogeography of bird hazards to aircraft warrants attention, and would usefully distinguish species, local, landscape and airport factors from any large-scale biogeographical influences (Fernández-Juricic *et al.* 2018; Alquezar *et al.* 2020). Broader-scale comparisons may provide useful patterns at higher spatial

scales (Alquezar *et al.* 2020), but given the degree of within-bioregion variation we describe here, such comparisons may be unlikely to usefully inform anything other than generalised management directions. Thus, currently the airport remains the ideal scale at which to plan and deliver bird hazard management. However, we suggest that an understanding of how spatial and biogeographical factors influence strikes would usefully inform planning for new aerodromes, or modifications to existing ones or their surrounding landscape context.

Conflicts of interest

WKS was contracted to provide wildlife hazards management advice at all three study sites, although only briefly at two of these. MAW declares no conflict of interest; he was not involved in design or data collection but in analysis and drafting.

Acknowledgements

This work analysed publicly available data; WKS collected data and assisted writing up. MAW did not collect data or design the project. MAW was supported during write up by the Beach Ecology and Conservation Hub (Venus Bay).

References

- Airservices Australia (2019). Movements at Australian Airports. Airservices Australia, Canberra. Available at <https://www.airservicesaustralia.com/publications/reports-and-statistics/movements-at-australian-airports/> [verified 14 August 2019].
- Alquezar, R. D., Tolesano-Pascoli, G., Gil, D., and Macedo, R. H. (2020). Avian biotic homogenization driven by airport-affected environments. *Urban Ecosystems* **23**, 507–517. doi:10.1007/s11252-020-00936-0
- Australian Transport Safety Bureau (ATSB) (2003). The hazard posed to aircraft by birds. Australian Transport Safety Bureau, Canberra, ACT, Australia.
- Australian Transport Safety Bureau (ATSB) (2008). An analysis of Australian birdstrike occurrences 2002 to 2006. ATSB Transport Safety Investigation Report, Aviation Research and Analysis Report AR-2008-027. Australian Transport Safety Bureau, Canberra, ACT, Australia.
- Australian Transport Safety Bureau (ATSB) (2010). Australian aviation wildlife strike statistics: bird and animal strikes 2002 to 2009. ATSB Transport Safety Report, Aviation Research and Analysis Report AR-2009-064. Australian Transport Safety Bureau, Canberra, ACT, Australia.
- Australian Transport Safety Bureau (ATSB) (2012). Australian aviation wildlife strike statistics: bird and animal strikes 2002 to 2011. ATSB Transport Safety Investigation Report, Aviation Research and Analysis Report AR-2012-031. Australian Transport Safety Bureau, Canberra, ACT, Australia.
- Australian Transport Safety Bureau (ATSB) (2014). Australian aviation wildlife strike statistics: 2004 to 2013. ATSB Transport Safety Report, Aviation Research Statistics AR-2014-075. Australian Transport Safety Bureau, Canberra, ACT, Australia.
- Australian Transport Safety Bureau (ATSB) (2017). Australian aviation wildlife strike statistics: 2006 to 2015. ATSB Transport Safety Report, Aviation Research Statistics AR-2016-063. Australian Transport Safety Bureau, Canberra, ACT, Australia.
- Australian Transport Safety Bureau (ATSB) (2019a). Australian aviation wildlife strike statistics: 2008 to 2017. ATSB Transport Safety Report, Aviation Research Statistics AR-2018-035. Australian Transport Safety Bureau, Canberra, ACT, Australia.
- Australian Transport Safety Bureau (ATSB) (2019b). ATSB National Aviation Occurrence Database. Australian Transport Safety Bureau. Available at <https://www.atsb.gov.au/avdata/> [verified 14 August 2019].
- Barrett, G., Silcocks, A., Barry, S., Cunningham, R., and Poulter, R. (2003). 'The New Atlas of Australian Birds.' (Birds Australia: Melbourne.)
- BASI (1996). Reported bird strikes in Australia. Information Bulletin, Bureau of Air Safety Investigation, Department of Transport and Regional Development, Canberra.
- Bunker, J. A., and Jones, D. N. (2008). A decade of bird monitoring at Brisbane Airport: bird abundance and aircraft bird-strikes 1996–2006. Report for Brisbane Airport Corporation by the Centre of Innovative Conservation Strategies, Griffith University, Brisbane.
- Burger, J. (1983). Bird control at airports. *Environmental Conservation* **10**, 115–124. doi:10.1017/S0376892900012200
- Callaghan, C. T., Major, R. E., Lyons, M. B., Martin, J. M., and Kingsford, R. T. (2018). The effects of local and landscape habitat attributes on bird diversity in urban greenspaces. *Ecosphere* **9**, e02347. doi:10.1002/ecs2.2347
- Chilvers, B. L., Ryan, C. J., and Hickling, G. J. (1997). Factors affecting pilot-reported bird strike rates at Christchurch International Airport, New Zealand. *New Zealand Journal of Zoology* **24**, 1–7. doi:10.1080/03014223.1997.9518100
- Cleary, E. C., and Dolbeer, R. A. (2005). 'Wildlife Hazard Management at Airports: A Manual for Airport Personnel,' 2nd edn. (Federal Aviation Administration in co-operation with US Department of Agriculture, Animal and Plant Health Inspection Service, Wildlife Services.)
- DeVault, T. L., Blackwell, B. F., and Belant, J. L. (2013). 'Wildlife in Airport Environment: Preventing Animal–Aircraft Collisions through Science-based Management.' (John Hopkins University Press: Baltimore, USA.)
- DELWP (2021). Bioregions and EVC benchmarks. Available at <https://www.environment.vic.gov.au/biodiversity/bioregions-and-evc-benchmarks> [verified 24 February 2021].
- Dolbeer, R. A., Belant, J. L., and Sillings, J. L. (1993). Shooting gulls reduces strikes with aircraft at John F. Kennedy International Airport. *Wildlife Society Bulletin* **21**, 442–450.
- El-Sayed, A. F. (2019). 'Bird Strike in Aviation: Statistics, Analysis and Management.' (John Wiley & Sons Ltd: Chichester, UK.)
- E&SS (1994). 'Wildlife Control Manual.' (Environment and Support Services, Airports Group, Transport Canada, Ontario, Canada.)
- Fernández-Juricic, E., Brand, J., Blackwell, B. F., Seamans, T. W., and DeVault, T. L. (2018). Species with greater aerial maneuverability have higher frequency of collisions with aircraft: a comparative study. *Frontiers in Ecology and Evolution* **6**, 17. doi:10.3389/fevo.2018.00017
- Linley, G., Kostoglou, K., Jit, R., and Weston, M. A. (2018). Australian magpies exhibit increased tolerance of aircraft noise on an airport, and are more responsive to take-off than to landing noises. *Wildlife Research* **45**, 282–286. doi:10.1071/WR18039
- Ministerie van Verkeer en Waterstaat (1999). 'Bird Control at Airports: An Overview of Bird Control Methods and Case Descriptions.' (Ministerie van Verkeer en Waterstaat: The Netherlands.)
- North American Bird Conservation Initiative, US Committee (2014). The state of the birds 2014 report. US Department of Interior, Washington, DC, USA.
- Pfeiffer, M. B., Kougher, J. D., and DeVault, T. L. (2018). Civil airports from a landscape perspective: a multi-scale approach with implications for reducing bird strikes. *Landscape and Urban Planning* **179**, 38–45. doi:10.1016/j.landurbplan.2018.07.004
- Soldatini, C., Albores-Barajas, Y. V., Lovato, T., Andreon, A., Torricelli, P., Montemaggiore, A., Corsa, C., and Georgalas, V. (2011). Wildlife strike risk assessment in several Italian airports: lessons from BRI and a new methodology implementation. *PLoS One* **6**(12), e28920. doi:10.1371/journal.pone.0028920
- Transport Canada (2004). 'Sharing the Skies: An Aviation Industry Guide to the Management of Wildlife Hazards,' 2nd edn. (Aviation Publishing Division, Transport Canada: Ottawa.) Available at <https://tc.canada.ca/en/aviation/publications/tp-13549-sharing-skies> [verified 1 October 2020].

Handling Editor: Stephanie Shwiff