

Identity and distribution of *Sonchus* leaf-curling mite (Acari: Eriophyidae) in New Zealand

Authors: Martin, Nicholas, and Zhang, Zhi-Qiang

Source: Systematic and Applied Acarology, 15(3) : 263-264

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.15.3.12>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Identity and distribution of *Sonchus* leaf-curling mite (Acari: Eriophyidae) in New Zealand

NICHOLAS MARTIN¹ & ZHI-QIANG ZHANG²

New Zealand Arthropod Collection, Landcare Research, Private Bag 92170, Auckland, New Zealand

¹ martinn@landcareresearch.co.nz ² zhangz@landcareresearch.co.nz (for correspondence)

Lamb (1960) in his summary of galls known in New Zealand lists a leaf edge roll on *Sonchus oleraceus* L. (Asteraceae) that was found in February 1951 at Mangere and Mount Albert in Auckland. The mite species associated with those damage symptoms was not identified, nor described in subsequent papers on New Zealand Eriophyoidea (Manson 1984a,b; Xue & Zhang 2008), though a European species, *Aceria sonchi* (Nalepa, 1902) (Acari: Eriophyidae), that causes leaf galls, was known (Knihinicki *et al.* 2009). This recent paper (Knihinicki *et al.* 2009) describes a new species of eriophyoid mite that induces leaf edge curling and rolling in *Sonchus* species in Australia. Knihinicki *et al.* (2009) also postulated that the New Zealand mite on *Sonchus* might be the same species as the newly described species from Australia based on the damage symptoms recorded by Lamb (1960).

After becoming aware of the symptoms of leaf edge rolling and curling on *Sonchus* species as reported in Knihinicki *et al.* (2009), *Sonchus* plants were examined and samples showing such symptoms were collected for mite identification. Plants of *Sonchus oleraceus* with leaf edge curling and rolling (Fig. 1) were found in November, 2008 and, March and July, 2009, and January 2010 at



FIGURE 1. *Sonchus oleraceus* L. (Asteracea) showing severe leaf curling and rolling induced by *Aceria thalgi* Knihinicki, 2009 (Acari: Eriophyidae). Collected on 1 November 2008 from Panmure Basin, Auckland, New Zealand (photograph by Nicholas A. Martin, copyright Plant & Food Research).

six locations, five in the Auckland Region and one in North Waikato. Mites from Mount Albert Research Centre, Auckland—one of the first sites recorded by Lamb (1960)—were identified as *Aceria thalgi* Knihinicki in Knihinicki *et al.* (2009) (See Fig. 2).

Plants showing symptoms of leaf curling and rolling were found on or close to beaches on the East and West coasts in Auckland and Waikato regions. At Raglan, Waikato, the plant was in the hollow of a sand dune, while two west coast beaches were stony above the high tide. Plants not growing on the beach were at the base of walls, mostly concrete. One stony beach had many infested plants, but at the other sites, damage symptoms were usually found on one or two plants near each other.

The distribution of plants in New Zealand suggests that they prefer sites that get hot in sunny weather during spring and summer. The isolated distribution of plants indicates that the mites have a high level of dispersal that enables them to colonize isolated plants.

Voucher specimens. Collected by N.A. Martin on 20 Jan. 2010 from the rolled leaf edges of *S. oleraceus* in Mt Albert Research Centre, Auckland; mites mounted in Hoyers medium on 4 slides, with serial number 10-858 Z; deposited in the New Zealand Arthropod Collection, Auckland.



FIGURE 2. Photomicrograph (differential interference contrast microscopy at 1000x oil immersion, Nikon E-90 microscope with the imaging software NIS-Elements BR 3.10) of *Aceria thalgi* Knihinicki, 2009 collected in Auckland, New Zealand. Female adult in dorsal view showing the pro-dorsal shield pattern and arrangement of microtubercles on the dorsal annuli. Scale: the distance between the bases of two *sc* setae is 28 μ m. Photograph by Zhi-Qiang Zhang.

References

- Knihinicki, D.K., McCarren, K.L. & Scott, J.K. (2009) A new species of *Aceria* (Acari: Eriophyidae) damaging sowthistles, *Sonchus* spp. (Asteraceae), in Australia with notes on *Aceria sonchi* (Nalepa, 1902). *Zootaxa*, 2119, 23–38.
- Lamb, K.P. (1960) A check list of New Zealand plant galls (Zooecidia). *Transactions of the Royal Society of New Zealand*, 88 (1), 121–139.
- Manson, D.C. (1984a) *Fauna of New Zealand No.4 Eriophyoidea except Eriophyinae* (Arachnida: Acari). Science Information Publishing Centre, DSIR, Wellington, New Zealand, 142 pp.
- Manson, D.C. (1984b) *Fauna of New Zealand No.5 Eriophyinae* (Arachnida: Acari: Eriophyoidea). Science Information Publishing Centre, DSIR, Wellington, New Zealand, 123 pp.
- Nalepa, A. (1902) Neue Gallmilben (21. Fortsetzung). *Anzeiger der kaiserlichen Akademie Wissenschaften. Mathematische-naturwissenschaftliche Klasse, Wien*, 39, 221–223.
- Xue, X.-F. & Zhang, Z.-Q. (2008) New Zealand Eriophyoidea (Acari: Prostigmata): an update with descriptions of one new genus and six new species. *Zootaxa*, 1962, 1–32.

Accepted by O. Seeman 26 Nov. 2010; published 10 Dec. 2010