

Parasitism of the carpenter bee, *Xylocopa virginica* (L.) (Hymenoptera: Apidae), by larval *Dermacentor variabilis* (Say) (Acari: Ixodidae)

Authors: Goddard, Jerome, and Bircham, Lawrence

Source: Systematic and Applied Acarology, 15(3) : 195-196

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.15.3.4>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Parasitism of the carpenter bee, *Xylocopa virginica* (L.) (Hymenoptera: Apidae), by larval *Dermacentor variabilis* (Say) (Acari: Ixodidae)

JEROME GODDARD¹ & LAWRENCE BIRCHAM²

¹Department of Entomology and Plant Pathology, 100 Twelve Lane, Clay Lyle Entomology, Mississippi State University, Mississippi State, MS 39762, U.S.A. E-mail: jgoddard@entomology.msstate.edu

²220 Martin Drive South, Brandon, MS 39042 U.S.A.

Abstract

Seven larval *Dermacentor variabilis* were found during April 2010 on a male carpenter bee, *Xylocopa virginica*, at Rayne, Louisiana, USA, while looking for chaetodactylid mites. One tick appeared to be partially engorged. Subsequently, 10 more carpenter bees (7 males, 3 females) were collected from the same location and examined for ticks. None were infested. The significance of this finding is discussed.

Ticks are obligate blood-sucking parasites of all terrestrial vertebrates: mammals, birds, reptiles and amphibians. Only very rarely have they been found attached to invertebrates. Hooker reported a beetle as the host of an *Amblyomma maculatum* (no stage specified) (Hooker *et al.* 1912), and an adult male *Rhipicephalus* (*Boophilus*) *annulatus* was reported from a horse fly (Leprince *et al.* 1988).

On April 4, 2010, seven larval *Dermacentor variabilis* were found on a male carpenter bee, *Xylocopa virginica*, at Rayne, Louisiana, by the second author while looking for chaetodactylid mites (specifically genus *Sennertia*) on two male bees. Deutonymphs of *Sennertia* are commonly phoretic on *Xylocopa* and *Ceratina* carpenter bees (Gerling *et al.* 1989). No attempt was made to initially scan the bee for attached mites; it was brushed with a toothbrush into a small pan of alcohol. Although no attached ticks were observed, one appeared to be partially engorged (Figure 1).




FIGURE 1. Larval *Dermacentor variabilis* removed from carpenter bee (Photo courtesy Joe Macgown, Mississippi State University).

Subsequently, on April 21, 2010, 10 more carpenter bees (7 males, 3 females) were collected from the same location and examined for ticks. None were infested. In the southern United States, *Dermacentor variabilis* larvae are active year-round, but accelerate host-seeking activity from February to mid-April (Sonenshine *et al.* 1966, Clark *et al.* 1998). They have been collected by drag cloth in northwest Florida during February (Cilek and Olson 2000). Since flowers provide the sole source of food and most of the water for carpenter bees (Gerling *et al.* 1989), we assume the infested bee acquired the tick larvae while visiting plants low to the ground, such as clover.

Acknowledgements

The tick identification was confirmed by Dr. Richard G. Robbins (ISD/AFPMB, Walter Reed Army Medical Center), and a voucher specimen is deposited in the Mississippi Entomological Museum, Mississippi State University (accession number 80-1). This article has been approved for publication as Journal Article No. J-11879 of the Mississippi Agricultural and Forestry Experiment Station, Mississippi State University.

References

- Cilek, J. E. & Olson, M.A. (2000) Seasonal distribution and abundance of ticks (Acari: Ixodidae) in northwestern Florida. *Journal of Medical Entomology*, 37, 439–44.
- Clark, K. L., Oliver, J. H., Jr., McKechnie, D. B. & Williams, D. (1998) Distribution, abundance, and seasonal activities of ticks collected from rodents and vegetation in South Carolina. *Journal of Vector Ecology*, 23, 89–105.
- Gerling, D., Velthuis, H. H. W. & Hefetz, A. (1989) Bionomics of the large carpenter bees of the genus *Xylocopa*. *Annual Review of Entomology*, 34, 163–190.
- Hooker, W. A., Bishopp, F. C. & Wood, H. P. (1912) The life history and bionomics of some North American ticks. USDA, Bureau of Entomology, Bulletin No. 106, 214 pp.
- Leprince, D. J., Foil, L. D. & Mullen, G. (1988) Parasitism of *Tabanus americanus* by *Boophilus annulatus*. *Journal of Entomological Science*, 23, 274–275.
- Sonenshine, D. E., Atwood, E. L. & Lamb, J. T. (1966) The ecology of ticks transmitting Rocky Mountain spotted fever in a study area in Virginia. *Annals of the Entomological Society of America*, 59, 1234–1262.

Accepted by R.G. Robbins 28 Sept. 2010; published 10 Dec. 2010