

## **Morphological ontogeny of *Lopheremaeus mirabilis* (Acari: Oribatida: Plateremaeidae), and comments on *Lopheremaeus* Paschoal**

Authors: Seniczak, Stanisław, Ivan, Otilia, Kaczmarek, Sławomir, and Seniczak, Anna

Source: Systematic and Applied Acarology, 25(12) : 2147-2164

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.25.12.1>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Morphological ontogeny of *Lopheremaeus mirabilis* (Acari: Oribatida: Plateremaeidae), and comments on *Lopheremaeus* Paschoal

STANISŁAW SENICZAK<sup>1\*</sup>, OTILIA IVAN<sup>2</sup>, SŁAWOMIR KACZMAREK<sup>1</sup> & ANNA SENICZAK<sup>3</sup>

<sup>1</sup>Department of Evolutionary Biology, Faculty of Biological Sciences, Kazimierz Wielki University, Bydgoszcz, Poland

<sup>2</sup>Department of Experimental and Applied Biology, Institute of Biological Research Iași - National Institute of Research & Development for Biological Sciences, Bucharest, Romania

<sup>3</sup>Department of Natural History, University of Bergen, University Museum of Bergen, Bergen, Norway

\*Corresponding author: [stseni@ukw.edu.pl](mailto:stseni@ukw.edu.pl)

### Abstract

The morphological ontogeny of *Lopheremaeus mirabilis* (Csiszár 1962), the type species of *Lopheremaeus* Paschoal, 1988 (Plateremaeidae) is described and illustrated. The adult of this species has six pairs of notogastral setae ( $lp$ ,  $h_1$ ,  $h_2$  and  $p$ -series), epimeral and anogenital hypertrichy, and 4–5 pairs of anal setae. The cuticle of juveniles is plicate, the larva lacks the gastronotal setae  $c_3$ ,  $la$ ,  $lm$  and  $h_3$ , the nymphs are multideficient (they lack also setae of  $d$ -series) and eupheredermous, i.e. they carry the exuvial scalps of previous instars on the gastronotum. In the adult, dorsal and ventral crests are present on all femora and trochanter III, and dorsal crest is present on trochanter IV, the nymphs have also dorsal and ventral crests on genua, whereas the larva has no crests. In all instars, seta  $d$  on all genua and tibiae is present, and all tarsi have basal bulb (containing muscles) and uniformly narrow distal stalk (with only tendons) at about mid-length. This species was reported only from Bulgaria, Bosnia-Herzegovina and North Macedonia, Romania and Georgia.

**Keywords:** oribatid mites, leg setation, leg crests, juveniles, stage structure

### Introduction

*Lopheremaeus mirabilis* (Csiszár 1962) was originally proposed as *Plateremaeus mirabilis* Csiszár, 1962 based on specimens from Bulgaria (Csiszár & Jeleva 1962), but now this species is known also from Bosnia-Herzegovina, North Macedonia, Romania and Georgia (Tarman 1983; Ivan & Vasiliu 2000; Murvanidze & Mumladze 2016). Paschoal (1988) revised the Plateremaeidae, and appointed it the type species of *Lopheremaeus* Paschoal, 1988. He presented the main diagnostic characters of the genus as: apodemes poorly sclerotized or absent; bothridial seta long, setiform, with very short spines at distal part, notogaster rounded, seta  $h_1$  well developed, epimeral formula 8-7-12-4, genital and anal openings separated, with seven and four pairs of setae, respectively. However, adults of *L. mirabilis* from Romania studied herein have also three pairs of  $p$ -series setae on the notogaster and some individuals have more anal setae, which were not mentioned in Csiszár and Jeleva (1962) paper, modifying the diagnosis of *Lopheremaeus* given by Paschoal (1988). Subías (2020) included three species in *Lopheremaeus*, and one of which he considered *species inquirenda*. According to the catalogue of juvenile oribatid mites by Norton and Ermilov (2014) and further literature, the juveniles of *Lopheremaeus* are not described and any other member of the family.

The aim of this paper is to describe and illustrate the morphological ontogeny of *L. mirabilis*.

## Material and methods

### Sampling

The juveniles and adults of *L. mirabilis* used in this study were collected on 7 July 2019 by O. Ivan from a Dacian oak-hornbeam forest, located in Eastern Romania (46°40'27"N, 28°03'35"E, 280 m a. s. l.), near the southern limit of the continental bioregion where it contacts the steppe. This forest is about 100 years old (Natura 2000 site), and is composed mainly of *Quercus petraea* (Matt.) Liebl., *Q. dalechampii* Ten., *Carpinus betulus* L. and *Tilia tomentosa* Moench. We investigated population parameters of *L. mirabilis*, including density and developmental stage structure. The gender of adults is not visible through the thick, reticulate integument, and therefore metric data (in  $\mu\text{m}$ ) were based on 15 randomly selected adults, which were dissected in order to determine the gender, number of gravid females and carried eggs, and included: total length (tip of rostrum to posterior edge of notogaster, measured in lateral aspect); maximum width (widest part of notogaster, measured in dorsal aspect); length of anal and genital openings; and length of setae (measured from the appropriate plane).

### Studies of type material

We studied the morphology of a specimen of *L. mirabilis* from the Csiszár collection stored at the Hungarian Natural History Museum (HNHM). This individual is in a good condition, except for legs, which are broken. It is labelled as follows: *Plateremaeus mirabilis* Csisz., 1962; Bulgaria, Karlovo–Kalofer; 1956 June 9–11, Balogh–Csiszár. Measurements: length—512, width—301.

### Illustrations and photographs

The illustrations of *L. mirabilis* are limited to the body regions that show substantial differences between instars, including the dorsal and lateral aspect and some leg segments of the larva, tritonymph and adult, ventral regions of all instars, and the palp and chelicera of the adult. Illustrations were prepared from individuals mounted temporarily on slides in lactic acid, using the open-mount technique (Grandjean 1949). In the text and figures, we used the following abbreviations: rostral (*ro*), lamellar (*le*), interlamellar (*in*) and exobothridial (*ex*) setae, bothridium (*bo*), bothridial seta (*bs*), notogastral or gastronal setae (*c*-, *d*-, *l*-, *h*-, *p*-series), cupules or lyrifissures (*ia*, *ih*, *ips*, *iad*), scalps of larva (L), protonymph (Pn) and deutonymph (Dn), subcapitular setae (*a*, *m*, *h*), cheliceral setae (*cha*, *chb*), palp setae (*inf*, *l*, *d*, *lt*, *vt*, *ul*, *su*) and solenidion  $\omega$ , Trägårdh's organ (*Tg*), Claparède's organ (*Cl*), epimeral setae (*1a*–*c*, *2a*, *3a*–*c*, *4a*–*c*), adanal and anal setae (*ad*-, *an*-series), aggenital setae (*ag*), leg setae (*bv*, *ev*, *d*, *l*, *ft*, *tc*, *it*, *p*, *u*, *a*, *s*, *pv*, *pl*, *v*), solenidia ( $\sigma$ ,  $\phi$ ,  $\omega$ ) and famulus ( $\epsilon$ ). Terminology used follows that of Grandjean (1951, 1953) and Norton and Behan-Pelletier (2009). The species nomenclature follows Subías (2004).

For scanning electron microscopy (SEM), the mites were air-dried and coated with Au/Pd in a Polaron SC502 sputter coater and placed on Al-stubs with double-sticky carbontape. Observations and micrographs were made with a ZEISS Supra 55VP scanning electron microscope. Light photographs were prepared from individuals mounted temporarily on slides in lactic acid, using a Leica DM3000 microscope and Leica DFC420 camera.

### *Lopheremaeus mirabilis* (Csiszár, 1962)

(Figs. 1–20)

*Plateremaeus mirabilis* Csiszár, 1962: Tarman 1983.

*Lopheremaeus mirabilis*: Paschoal 1988; Subías 2004; Murvanidze and Mumladze 2016.

### Diagnosis


Adults of medium size (592–656), oval, with rounded rostrum, and polygonal pattern on prodorsum and notogaster. Bothridial seta setiform, with very short spines near distal end. Notogaster elliptical in dorsal aspect, with six pairs of notogastral setae ( $lp$ ,  $h_1$ ,  $h_2$  and  $p$ -series). Formula of epimeral setae 8-7-12-4 (hypertichy occurs), genital setae (seven pairs) in two rows (4 pairs in inner rows, 3 pairs in lateral rows), four pairs of aggenital setae and 4–5 pairs of anal setae; all short and smooth. Adanal seta  $ad_3$  inserted roughly in alignment with lateral margin of genital and anal plate, at level about midway between them, other two pairs in lateral position to anal plates. All femora and trochanter III with dorsal and ventral crests, trochanter IV with dorsal crest. Solenidion  $\phi_1$  on tibia I on large apophysis, covering basal dorsal part of tarsus I, famulus  $\epsilon$  on tarsus I short, semispherical. Seta  $d$  present on all genua and tibiae, slightly removed from respective solenidion. All tarsi with basal bulb and uniformly narrow distal stalk about mid-length, with three claws and short setae  $p$ .

Prodorsum in juveniles reticulate, prodorsal setae  $ro$  and  $le$  of medium size, setae  $in$  and  $ex$  short. Bothridial seta clavate in larva, setiform in nymphs. Larva with eight pairs of gastronotal setae ( $c_3$ ,  $la$ ,  $lm$  and  $h_3$  absent), nymphs with same number ( $d$ -series lost,  $p$ -series appears). In larva, crests on legs not evident, nymphs with dorsal and ventral crests on all femora, genua and trochanter III, and dorsal crest on trochanter IV. On all genua and tibiae seta  $d$  present, slightly removed from respective solenidion. All tarsi with basal bulb and uniformly narrow distal stalk at about mid-length, with one claw and short setae  $p$ .

### Morphology of adult

Adult (Figs. 1–8) similar to that described by Csiszár and Jeleva (1962), but see Remarks. Mean length (range) of females  $633.3 \pm 13.5$  (606–656,  $N=9$ ) and mean width (range)  $374.6 \pm 13.5$  (352–400), and mean length (range) of males  $611.0 \pm 11.9$  (592–624,  $N=6$ ) and mean width (range)  $384.7 \pm 42.1$  (352–468). Setae  $ro$  and  $le$  (Figs. 1a, 2, 3a, 5a–c, 6a, 6d, Table 1), and bothridial seta (Figs. 1c, 5a–c, 6a–c) as in Csiszár and Jeleva (1962). Notogaster reticulate (Figs. 1a, 1b, 3a, 5, 6a–c, 7a, 8a), setae (6 pairs,  $lp$ ,  $h_1$ ,  $h_2$  and  $p$ -series) strongly curved inwards, smooth and long (Table 1),  $p$ -series slightly thinner than others (Figs. 1a, 2, 3a, 5d, 7a, 8a). Subcapitular setae  $h$ , and  $m$  slightly longer than  $a$ , all smooth. Formula of epimeral setae 8-7-12-4 (I–III hypertrichous, variation not observed); setae in first row of epimere I clearly longer than others, all smooth (Fig. 2). Genital setae (7 pairs) in two rows, inner setae (4 pairs) and lateral setae (3 pairs) of similar length, all smooth. Aggenital setae (4 pair) similar to genital setae. Adanal seta  $ad_3$  inserted roughly in alignment with lateral margin of genital and anal plate, at level about midway between them; all strongly curved and smooth. Anal setae (4–5 pairs) similar to genital setae. Chelicera relatively slim, seta  $cha$  longer than  $chb$ , both barbed, in some individuals second seta  $chb$  present (Figs. 3b, 3c). Palp relatively short and thin, solenidion  $\omega$  separated from seta  $acm$  (Fig. 3d). Formula of palp setae (and solenidion): 0-2-1-3-9(1). Setae  $sup$ ,  $inf$ ,  $l''$  and  $d$  finely barbed, other setae and eupathidia shorter, smooth. Leg segments protected by retrotecta. All femora and trochanter III with dorsal and ventral crests, trochanter IV with dorsal crest (Figs. 1a, 2, 3a, 4, 5b, 5d, 7a–c, 8b–d). Most leg setae relatively long, strongly curved and smooth, most distal setae on tarsi slightly curved or almost straight and finely barbed. Solenidion  $\phi_1$  on tibia I long, on large apophysis, covering large dorsal part of tarsus I; tarsal solenidia  $\omega_1$  and  $\omega_2$  clearly shorter, other solenidia short. Famulus  $\epsilon$  semispherical (Figs. 4a, 7d, 8b). Seta  $d$  present on all genua and tibiae, removed from the respective solenidion (Fig. 4). All tarsi with basal bulb (containing muscles) and uniformly narrow distal stalk (with only tendons) at about mid-length, with three claws and short setae  $p$  (Figs. 4, 7d, 8b). Some parts of leg segments and leg setae covered with granular cerotegument. Formulae of leg setae (and solenidia, from trochanter to tarsus): I–1-6-4(1)-5(2)-20(2); II–1-5-4(1)-5(1)-15(2); III–3-5-3(1)-4(1)-15; IV–3-5-3-4(1)-12.





**FIGURES 1–2.** *Lopheremaeus mirabilis*, adult, legs partially drawn, scale bars 50  $\mu\text{m}$ . 1. (a) Dorsal aspect, (b) central part of notogaster, (c) bothridial seta (b, c, enlarged). 2. Ventral aspect.

#### Remarks


Females and males investigated herein are slightly larger than those studied by Csiszár and Jeleva (1962) — holotype length 530, width 312, sex not investigated — and have three pairs of setae of *p*-series, which were overlooked by these authors. The topotype specimen from HNHN is smaller (512 x 301) than holotype, has the notogastral setae of *p*-series (strongly curved, visible in ventral and posterior aspects) and dorsal crest on trochanter III. The adults from Romania have more crests on the legs, 4–5 pairs of anal setae (versus four pairs in Csiszár and Jeleva 1962), but not observed by Csiszár and Jeleva (1962). These authors mentioned in *Plateremaeus* Berlese, 1908 crispins (or retroecta) on legs and 4–5 pairs of anal setae, but this genus was further divided in *Plateremaeus* and *Lopheremaeus* (Paschoal 1988). In our individuals, the shape of leg setae, solenidion  $\phi_1$  on tibia I and famulus  $\epsilon$  on tarsus I are similar as in Csiszár and Jeleva (1962). All these characters widen the diagnosis of *Lopheremaeus* given by Csiszár and Jeleva (1962).

#### Description of juvenile stages

Larva oval in dorsal aspect (Figs. 9a, 10a) and unpigmented. Prodorsum subtriangular, prodorsal setae *ro* and *le* of medium size (Table 1), *in* and *ex* short. Mutual distance between setal pairs *le* and *in* about two times longer than between pair *ro*. Seta *le* inserted closer to *ro* than to *in*. Bothridium large, rounded, bothridial seta long, clavate and barbed. Central part of prodorsum reticulate, posterior part punctate, and ridge present between bothridium and central part of prodorsum (Figs. 9b, 10a). Some parts of body and most setae covered with granular cerotegument.

Gastronotum of larva plicate, relatively flat, with horizontal edge and eight pairs of setae, including *h*<sub>2</sub> inserted lateral to medial part of paraproctal valves (Figs. 10a, 11a, 12a), setae *c*<sub>3</sub>, *la*, *lm* and *h*<sub>3</sub> not observed. Setae *c*<sub>1</sub> and of *c*<sub>2</sub> short (Table 1), other setae of medium size, all curved, inserted on small apophyses and covered with granular cerotegument, but setae of *d*-series with thicker

granular cerotegument than other setae, and appearing barbed. Cupule *ih* lateral to anterior part of anal valves, other cupules and gland opening not observed in plicate integument. Paraproctal valves (segment PS) glabrous.




**FIGURE 3.** *Lopheremaeus mirabilis*, adult. (a) Lateral aspect, legs partially drawn, scale bar 50 µm; mouthparts, right side, scale bars 10 µm; (b) chelicera, (c) part of chelicera of another individual, (d) palp.

All tarsi with thick basal bulb and uniformly narrow distal stalk at about mid-length, with one claw and short setae *p* (Figs. 9a, 13). Crests on legs not evident. Solenidion  $\phi_1$  on tibia I on large apophysis, covering large dorsal part of tarsus I, other solenidia short or of medium size, famulus  $\varepsilon$  on tarsus I relatively long, with apical globule. Most distal setae on tarsi long, slightly curved or nearly straight, other setae long or of medium size and curved; all smooth. Seta *d* present on all genua and tibiae, removed from the respective solenidion. Some parts of leg segments and setae covered with granular cerotegument.


Shape and colour of nymphs (Figs. 9c, 9d) as in larva, but prodorsal seta *ro* and *le* relatively longer (Table 1), strongly curved inwards, and *le* inserted closer to *ro* than in larva, setae *in* and *ex* short. Bothridium rounded, with posterolateral edge, bothridial seta setiform, covered with granular cerotegument. In protonymph, anterior and medial part prodorsum reticulate, and lateral and posterior parts punctate. Gastronotum with eight pairs of setae: *p*-series appears in protonymph (Fig. 11b) and remains in deutonymph and tritonymph (Figs. 14a, 14b), and *d*-series lost and remains absent through rest of ontogeny. In all nymphs, setae *c*<sub>1</sub> and *c*<sub>2</sub> short, *h*<sub>1</sub> long, and other setae of medium size (Table 1). Dorsal part of gastronotum flat, with exuvial scalps of previous instars, other gastronotal setae in marginal position on gastronotum (Figs. 15b, 18, 19b). Seta *h*<sub>1</sub> curved upwards and can be helpful in carrying exuvial scalps. Genital valves of protonymph with one pair of setae, three pairs added in deutonymph (one pair in inner row and two pairs in outer row), and one pair added in tritonymph in inner row (Figs. 1b, 14a, 14b); all short and usually covered with granular

cerotegument. In deutonymph, one pair of aggenital setae appearing and remaining through ontogeny. Paraproctal valves of protonymph (segment AD) and deutonymph (segment AN) glabrous, but in tritonymph 4–5 pairs of anal setae present, all short and smooth. In protonymph, cupule *ips* lateral to anal valves, other cupules and gland opening not observed, possibly obscured by granular cerotegument in this and other nymphs (Figs. 11b, 14a, 14b).

In all nymphs, hypertrichy present in epimeral region, variation not observed. In protonymph, one pair of setae added on epimere II, and one pair on epimere III, compared to larva (Figs. 16a, 16b), in deutonymph three pairs of setae added on epimere III, and one pair added on epimere IV (Figs. 17b, 17a), compared to protonymph. In tritonymph, four pairs of setae added on epimere I, three pairs added on epimere II, one pair added on epimere III, and two pairs added on epimere IV, compared to deutonymph (Figs. 17a, 17b).



**FIGURE 4.** *Lopheremaeus mirabilis*, leg segments of adult (part of femur to tarsus), right side, seta on the opposite side not illustrated, but indicated in the legend, scale bar 20  $\mu$ m. (a) Leg I, tarsus (*pl'*); (b) leg II; (c) leg III; (d) leg IV; (e) part of femur and genu IV (slightly enlarged).



**FIGURE 5.** *Lopheremaeus mirabilis*, adult, SEM micrographs. (a) Dorsal view, (b), lateral view, (c) frontal view, (d) posteroventral view.

In nymphs, dorsal and ventral crests present on trochanter III, all femora and genua, and dorsal crest present on trochanter IV (Figs. 12b, 18, 19, 20). Solenidion  $\phi_1$  on tibia I on large apophysis, covering large dorsal part of tarsus I, other solenidia short or of medium size, famulus  $\epsilon$  on tarsus I semispherical. On all genua and tibiae, seta *d* present, removed from respective solenidion (Fig. 20). All tarsi with basal bulb and uniformly narrow distal stalk at about mid-length, with one claw and short setae *d*. Most distal setae on tarsi long, slightly curved or nearly straight and finely barbed, other setae long or of medium size, curved and smooth. Some parts of leg segments and leg setae covered with granular cerotegument.

#### *Summary of ontogenetic transformations*

In all juveniles of *L. mirabilis*, the prodorsal setae *ro* and *le* are of medium size, while *in* and *ex* are short; by contrast, in the adult seta *ex* is of medium size, but shorter than *ro* and *le*. In all instars, the opening of the bothridium is rounded, but the bothridial seta is clavate in the larva, and setiform in other instars. The larva has eight pairs of gastronotal setae (*c*-, *d*-series, *lp*, *h*<sub>1</sub>, *h*<sub>2</sub>), and the nymphs have also eight pairs (*p*-series appears, *d*-series is lost in protonymph), but the notogaster of the adult loses the *c*-series, such that six pairs of notogastral setae remain. The formula of gastronotal setae of *L. mirabilis* is therefore 8-8-8-8-6 (from larva to adult). Formulae of epimeral setae are: 3-1-2 (larva, including scaliform *lc*), 3-2-3-1 (protonymph), 3-2-6-2 (deutonymph), 7-5-7-4 (tritonymph) and 8-7-12-4 (adult). Formula of genital setae is 1-4-5-7 (protonymph to adult), aggenital setae is 1-1-4 (deutonymph to adult), and segments PS-AN is 03333-0333-0(4-5)(4-5). Seta *d* is present on genua and tibiae of all instars. The ontogeny of leg setae and solenidia is given in Table 2.


**TABLE 1.** Measurements of some morphological characters of juvenile stages and adult of *Lopheremaeus mirabilis* (mean measurements of 3–7 juveniles and 10 adults in  $\mu\text{m}$ ); Nd—not developed.

Morphological characters	Larva	Protonymph	Deutonymph	Tritonymph	Adult
Body length	281	371	468	579	631
Body width	150	208	281	310	337
Length of prodorsum	104	142	159	215	208
Length of: seta <i>ro</i>	25	38	54	64	79
seta <i>le</i>	14	34	52	59	85
seta <i>in</i>	2	2	3	4	8
seta <i>ex</i>	3	3	4	5	53
seta <i>bs</i>	80	89	104	116	149
seta <i>c1</i>	4	7	11	17	Lost
seta <i>c2</i>	4	6	10	16	Lost
seta <i>da</i>	27	Lost	Lost	Lost	Lost
seta <i>dp</i>	28	Lost	Lost	Lost	Lost
seta <i>lp</i>	32	20	21	22	83
seta <i>h1</i>	27	88	102	117	61
seta <i>h2</i>	15	34	44	56	63
seta <i>p1</i>	Nd	19	25	32	65
seta <i>p3</i>	Nd	16	23	30	63
genital opening	Nd	32	49	72	88
anal opening	28	61	84	117	120

#### *Distribution, ecology and biology*

According to Subías (2004, 2020), *L. mirabilis* has an eastern Mediterranean distribution, and this species was reported only from Bulgaria, Bosnia-Herzegovina and North Macedonia, Romania and Georgia (Csiszár & Jeleva 1962, Tarman 1983, Ivan & Vasiliu 2000, Murvanidze & Mumladze 2016). This species inhabits alpine meadows, forests and urban soils (Murvanidze & Mumladze 2016). It was collected in Romania for the first time in 1992 in a small number (Ivan & Vasiliu 2000), but was misidentified as *Lopheremaeus laminipes* (Berlese 1916). Our additional collection and the comparison with a specimen from the Csiszár collection allowed us to identify these individuals as *L. mirabilis*. Thus, this is the first record of *L. mirabilis* in Romania, and represents the most northern record of the species.

In our study, the density of *L. mirabilis* in the soil of Dobrina forest (Romania) varied between May and July, being highest in June 2014 (6.4 individuals per 500  $\text{cm}^3$ ), and lowest in May 1992 (0.4 individuals per 500  $\text{cm}^3$ , Table 3). In most samples, adults dominated the juveniles, comprising 41–64% of individuals. In the sample with the most abundant juveniles (July 2019), the juveniles dominated (64% of individuals) the adults, and the stage structure of *L. mirabilis* was the following: seven larvae, four protonymphs, four deutonymphs, three tritonymphs and 10 adults. Among 15 dissected adults, nine were females and six males, but we rather not like to give the sex ratio of this species because these mites were collected in different dates. Most females (78%) were gravid, carrying 1–3 (mostly two) relatively large eggs, each 247–271 x 181–199, and one of which constitutes about 40% of the total body length of females.




**FIGURE 6.** *Lopheremaeus mirabilis*, adult, SEM micrographs. (a) Anterior part, dorsal view, (b) bothridium and bothridial seta, (c) bothridial setae and notogaster, frontal view, (d) rostrum, frontal view.

### Comparison of morphology of *Lopheremaeus mirabilis* with congener and remarks


The adult of *L. mirabilis* is morphologically similar to that of *L. laminipes*, but Csiszár and Jeleva (1962) recognized the latter species as distinct mainly by the lighter cerotegument and notogastral sculpture, without dense black punctation, and more posterior location of all notogastral setae in the latter species. However, in the figure of *L. laminipes* drawn by Mahunka and Mahunka-Papp (1995) the location of posterior setae on the notogaster is similar to that of *L. mirabilis* as shown by Csiszár and Jeleva (1962) and specimens studied herein, which suggests that the distinction of these species needs more investigation.

The homologies of notogastral setae in the adult of *L. mirabilis* are inconsistently reported in the literature. Csiszár and Jeleva (1962) observed only three pairs of notogastral setae, and labeled none of them. Paschoal (1988) revised the Plateremaeidae and used the old Grandjean's (1951) "*Dometorina* nomenclature" of notogastral setae, but he treated the notogastral setae  $r_2$  and  $r_3$  as homologues of  $lp$  and  $lm$ , respectively, which is inconsistent with Grandjean's (1951) unideficient nomenclature. In our opinion, setae  $r_2$  and  $r_3$  are located posterior to porose areas  $A1$  and  $A2$ , respectively, and should be labeled as  $h_3$  and  $h_2$ , respectively. In the case of *L. mirabilis*, the adult has no porose areas, seta  $h_3$  is absent in the larva, and it is difficult to judge if this seta appears in the protonymph because normally this seta occupies the lateral position, as seta  $lp$ , which is present in the larva. There is no trace that  $h_3$  appears in the protonymph and  $lp$  is lost, and therefore we consider seta  $lp$  in all nymphs and adult of *L. mirabilis*.





**FIGURE 7.** *Lopheremaeus mirabilis*, adult, SEM micrographs. (a) Legs III and IV, dorsal view, (b) legs I and II, dorsal view, (c) femur and part of genu III, lateral view, (d) tarsus I, lateral view (arrow points famulus  $\epsilon$ ).



**FIGURE 8.** *Lopheremaeus mirabilis*, adult, photo, scale bars 50  $\mu$ m. (a) Notogaster, dorsal view, (b) part of leg I, lateral view (arrow points famulus  $\epsilon$ ), (c) femur and genu III, lateral view, (d) femur IV, dorsal view (setae broken).

**TABLE 2.** Ontogeny of leg setae (Roman letters) and solenidia (Greek letters) of *Lopheremaeus mirabilis*.

Leg	Trochanter	Femur	Genu	Tibia	Tarsus
<b>Leg I</b>					
Larva	–	$d, bv''$	$(l), d, \sigma$	$(l), v', d, \phi_1$	$(fi), (tc), (p), (u), (a), s, (pv), (pl) \varepsilon, \omega_1$
Protonymph	–	$(l)$	–	–	$\omega_2$
Deutonymph	$v'$	–	$v'$	$v'', \phi_2$	–
Tritonymph	–	$v''_1$	–	–	$(it)$
Adult	–	$v''_2$	–	–	$l'', v'$
<b>Leg II</b>					
Larva	–	$d, bv''$	$(l), d, \sigma$	$l', v', d, \phi$	$(fi), (tc), (p), (u), (a), s, (pv), \omega_1$
Protonymph	–	$(l)$	–	$l''$	–
Deutonymph	$v'$	–	$v'$	$v''$	$\omega_2$
Tritonymph	–	$v''$	–	–	$(it)$
Adult	–	–	–	–	–
<b>Leg III</b>					
Larva	–	$d, ev'$	$l', d, \sigma$	$v', d, \phi$	$(fi), (tc), (p), (u), (a), s, (pv)$
Protonymph	$v'_1$	–	–	–	–
Deutonymph	$l'$	$l'_1$	$v'$	$l', v''$	–
Tritonymph	–	–	–	–	$(it)$
Adult	$v'_2$	$l'_2, v'$	–	–	–
<b>Leg IV</b>					
Protonymph	–	–	–	–	$fi'', (p), (u), (pv)$
Deutonymph	$v'_1$	$d, ev'$	$d, l'$	$v', d, \phi$	$(a), s$
Tritonymph	$v'_2$	$l'_1$	$v'$	$l', v''$	$(tc)$
Adult	$v'_3$	$l'_2, v'$	–	–	–


Note: structures are indicated where they are first added and are present through the rest of ontogeny; pairs of setae in parentheses, dash indicates no additions.

The presence of epimeral hypertrichy and crests on some leg segments support the membership of *L. mirabilis* in Plateremaeidae. Epimeral hypertrichy also occurs in *Plateremaeus*, *Allodamaeus* Banks, 1947, *Paralopheremaeus* Paschoal, 1988 and *Calipteremaeus* Paschoal, 1988; and in *Paralopheremaeus* small crests also are present on some leg segments (Paschoal 1988). Epimeral hypertrichy also occurs in *Hermannia reticulata* Thorell, 1871 and *H. scabra* (Koch 1879), but in *L. mirabilis* and *H. reticulata* hypertrichous are epimeres I–III, whereas in *H. scabra* epimeres III and IV (Seniczak *et al.* 2017a, b).


Seniczak and Seniczak (2011) and Seniczak *et al.* (2012, 2016) compared the morphological ontogeny of some species of Plateremaeoidea, and the juveniles of *L. mirabilis* are most similar to those of *Gymnodamaeus bicostatus* (Koch 1836), which have plicate cuticle. However, the larva of *G. bicostatus* has 11 pairs of gastronotal setae ( $h_3$  absent), whereas that of *L. mirabilis* has eight pairs ( $c_3, la, lm$  and  $h_3$  absent). The nymphs of the former species have 10 pairs of gastronotal setae ( $lp, c$ -,  $h$ - and  $p$ -series), whereas those of the latter species have eight pairs ( $c_1, c_2, lp$ -series,  $h_1, h_2$  and  $p$ -series). The adult of *G. bicostatus* retains five pairs of notogastral setae ( $h_1, h_2$  and  $p$ -series), whereas that of *L. mirabilis* retains six pairs ( $lp, h_1, h_2$  and  $p$ -series). The nymphs of both species carry the exuvial scalps, but *L. mirabilis* has long, curved upwards seta  $h_1$ , which can be helpful in carrying the scalps, whereas in *G. bicostatus* this seta is short (Seniczak & Seniczak 2011). *Aleurodamaeus*





*setosus* (Berlese 1883), *Arthrodamaeus italicus* (Berlese 1916), *A. mongolicus* Bayartogtokh & Weigmann, 2005 and *Licnobelba latiflabellata* (Paoli 1908) have also long, curved upwards seta  $h_1$ , which can be helpful in carrying the exuvial scalps (Seniczak *et al.* 2012, 2016).



**FIGURE 9.** *Lopheremaeus mirabilis*, photo, dorsal view. (a) Larva, (b) anterior part of body of larva, (c) tritonymph without exuvial scalps, (d) prodorsum of tritonymph. Scale bars (um): a —100; b, d —50 um; c —200.




**FIGURE 10–11.** *Lopheremaeus mirabilis*, legs partially drawn, scale bars 20  $\mu$ m. 10. (a) Larva, dorsal aspect, (b) edge of gastronetum (enlarged). 11. Ventral part of hysterosoma, (a) larva, (b) protonymph.




**TABLE 3.** Density (individuals/500cm<sup>3</sup>), stage structure and vertical distribution of *Lopheremaeus mirabilis* in the soil of Dobrina forest (Romania); L—larva, Pn—protonymph, Dn—deutonymph, Tn—tritonymph, Juv—juveniles, Ad—adult, Olf—litter and fermentation layer, Ah—humus layer, C—constancy in five samples (in%).

Date of sampling	Juveniles						Adult	Total	Indiv/500cm <sup>3</sup>	Presence in Olf/Ah	C
	L	Pn	Dn	Tn	Juv	%					
May 1992	0	0	0	0	0	0	2	2	0.4	1/1	20
July 1992	0	0	0	0	0	0	4	4	0.8	2/2	40
June 2014	0	5	4	4	13	41	19	32	6.4	29/3	40
May 2019	0	1	2	2	5	45	6	11	2.2	11/0	20
July 2019	7	4	4	3	18	64	10	28	5.6	25/3	40
Total	7	10	10	9	36	47	41	77	3.1 <sup>1</sup>	16.6/1.8 <sup>1</sup>	32 <sup>1</sup>


<sup>1</sup>Mean value.




**FIGURES 14–15.** *Lopheremaeus mirabilis*, legs partially drawn, scale bars 50 μm. 14. Ventral part of hysterosoma, (a) deutonymph, (b) tritonymph. 15. Tritonymph, (a) dorsal aspect, (b) reticulation of prodorsum, (c) edge of gastrnotum (b, c, enlarged).




**FIGURES 16–17.** *Lopheremaeus mirabilis*, epimeral regions, partially drawn, scale bars 50 μm. 16. (a) Larva, (b) protonymph. 17. (a) Deutonymph, (b) tritonymph.



**FIGURE 18.** *Lopheremaeus mirabilis*, tritonymph with exuvial scalps, dorsal view. (a) Whole body, (b) prodorsum and gastronetum, (c) prodorsum, (d) legs III and IV.



**FIGURE 19.** *Lopheremaeus mirabilis*, tritonymph. (a) Without exuvial scalps, dorsal view, (b) with exuvial scalps, lateral view, (c) ventral view, (d) hypostomal region.



**FIGURE 20.** *Lopheremaeus mirabilis*, leg segments of tritonymph (part of femur to tarsus), right side, seta on the opposite side not illustrated, but indicated in the legend, scale bar 20  $\mu$ m. (a) Leg I, tarsus ( $pl'$ ); (b) leg II; (c) leg III; (d) leg IV.



## Acknowledgements

We thank Prof. Dr. Roy A. Norton (State University of New York, Syracuse, New York, USA) and anonymous reviewer for thorough review of the manuscript and very helpful suggestions that improved the scientific value of this paper. We also thank Dr. László Dányi, the curator of the Soil Zoology Collection in the Hungarian Natural History Museum, Budapest, for loan of a specimen of *L. mirabilis* for our study, and are grateful to Eng. Irene Heggstad (Department of Earth Science, University of Bergen) for her professional help with SEM images. The ecological part of our investigation was supported by a grant of Romanian Ministry of Research and Innovation through Program 1, contract no. 22PFE / 2018. This study was done under the program of the Polish Minister of Science and Higher Education "Regional Initiative of Excellence" in 2019–2022 (Grant No. 008/RID/2018/19).

## References

- Banks, N. (1947) On some Acarina from North Carolina. *Psyche*, 54(2), 110–141.  
<https://doi.org/10.1155/1947/70181>
- Bayartogtokh, B. & Weigmann, G. (2005) New and little known species of oribatid mites of the genera *Arthrodamaeus* and *Fuscozetes* (Arachnida: Acari: Oribatida) from Mongolia. *Species Diversity*, 10, 75–84.  
<https://doi.org/10.12782/specdiv.10.75>
- Berlese, A. (1883) Escursione in Sicilia. Acarofauna Sicula. Ia serie. *Bollettino della Società entomologica italiana*, 15, 212–220.
- Berlese, A. (1908) Elenco di generi e specie nuove di Acari. *Redia*, 5(1), 1–15.
- Berlese, A. (1916) Centuria prima di Acari nuovi. *Redia*, 12(1), 19–67.
- Csiszár, J. & Jeleva, M. (1962) Oribatid mites (Acari) from Bulgarian soils. *Acta Zoologica Academiae Scientiarum Hungaricae*, 8(3–4), 273–301.
- Grandjean, F. (1949) Observation et conservation des tres petits Arthropodes. *Bulletin du Muséum National d'Histoire Naturelle*, Series 2, 3, 363–370.
- Grandjean, F. (1951) Sur deux espèces du genre "*Dometorina*" n.g. et les mœurs de "*D. plantivaga*" (Berl.) (Acariens, Oribates). *Bulletin de la Société zoologique de France*, 75, 224–242.
- Grandjean, F. (1953) Essai de classification des Oribates (Acariens). *Bulletin de la Société zoologique de France*, 78, 421–446.
- Ivan, O. & Vasiliu, M. (2000) Oribatid fauna (Acari, Oribatida) in Romanian *Quercus* forests. *Anuarul Muzeului Național al Bucovinei*, 15, 67–116.
- Koch, C.L. (1936) Deutschlands Crustaceen, Myriapoden und Arachniden. *Friedrich Pustet*, Regensburg, Bd. 4–9.
- Koch, L. (1879) Arachniden aus Sibirien und Novaja Zemlja, eingesammelt von der Schwedischen Expedition im Jahre 1875. *Kongliga Svenska Vetenskaps Akademiens Handlingar*, Stockholm, 16(5), 1–136.
- Mahunka, S. & Mahunka-Papp, L. (1995) The oribatid species described by Berlese (Acari). *Hungarian Natural History Museum*, Budapest, 325 pp.
- Murvanidze, M. & Mumladze, L. (2016) Annotated checklist of Georgian oribatid mites. *Zootaxa*, 4089(1), 1–81.  
<https://doi.org/10.11646/zootaxa.4089.1.1>
- Natura (2000) Dobrina forest ROSCI0335, standard form (pdf). (In Romanian)  
<http://biodiversitate.mmediu.ro/rio/natura2000/static/pdf/rosci0335>
- Norton, R.A. & Behan-Pelletier, V.M. (2009) Suborder Oribatida. In: Krantz, G.W., Walter, D.E. (Eds.), *A manual of Acarology, 3rd Edition*. Texas Tech University Press, Lubbock, pp. 430–564.
- Norton, R.A. & Ermilov, S.G. (2014) Catalogue and historical overview of juvenile instars of oribatid mites (Acari: Oribatida). *Zootaxa*, 3833, 1–132.  
<https://doi.org/10.11646/zootaxa.3833.1.1>
- Paoli, G. (1908) Monografia del genere *Dameosoma* Berl. e generi affini. *Redia*, 5, 31–91.
- Paschoal, A.D. (1988) A revision of the Plateremaeidae (Acari: Oribatei). *Revista Brasileira de Zoologia*, 3(6) (1987), 327–356.  
<https://doi.org/10.1590/S0101-81751986000200001>

- Seniczak, S., Ayyıldız, N. & Seniczak, A. (2012) Setal losses in the dorsal hysterosoma of Plateremaeoidea (Acari: Oribatida) in the light of ontogenetic studies. *Journal of Natural History*, 46(7–8), 411–451.  
<https://doi.org/10.1080/00222933.2011.640464>
- Seniczak, S. & Seniczak, A. (2011) Ontogenetic studies of three species of Gymnodamaeidae (Acari: Oribatida) with a focus on regressions of hysterosomal setae. *Journal of Natural History*, 45(5–6), 361–391.  
<https://doi.org/10.1080/00222933.2010.534188>
- Seniczak, S., Seniczak, A. & Coulson, S.J. (2017a) Morphological ontogeny, distribution, and descriptive population parameters of *Hermannia reticulata* (Acari: Oribatida: Hermannidae), with comments on Crotonioidea. *International Journal of Acarology*, 43(1), 52–72.  
<https://doi.org/10.1080/01647954.2016.1229812>
- Seniczak, S., Seniczak, A. & Coulson, S.J. (2017b) Morphological ontogeny and distribution of *Hermannia scabra* (Acari: Oribatida: Hermannidae) in Svalbard and descriptive population parameters. *Acarologia*, 57(4), 877–892.  
<http://doi:10.24349/acarologia/20174214>
- Seniczak, S., Seniczak, A. & Kaczmarek, S. (2016) Morphological ontogeny, distribution and ecology of *Arthrodamaeus italicus* and *A. mongolicus* (Acari: Oribatida: Gymnodamaeidae). *International Journal of Acarology*, 42(3), 174–192.  
<https://doi.org/10.1080/01647954.2016.1151931>
- Subías, L.S. (2004, updated in 2020) Listado sistemático, sinonímico y biogeográfico de los Ácaros Oribátidos (Acariformes, Oribatida) del mundo (1758–2002). *Graellsia*, 60 (número extraordinario), 3–305. 15ª actualización, 527 pp. (accessed February 2020).  
<https://doi.org/10.3989/graellsia.2004.v60.iExtra.218>
- Tarman, K. (1983) Catalogus faunae Jugoslaviae III/4 Acarina Oribatei. Consilium Academicarum Scientiarum Rei Publicae Socialistae Foederativae Jugoslaviae. Ljubljana, *Academia Scientiarum et Artium Slovenica*, 3, 1–61.
- Thorell, T. (1871) Om Arachnider fran Spitzbergen och Beeren-Eiland. *Öfversigt af Kongliga Vetenskaps-Akademiens Förhandlingar*, 28, 83–702.

*Submitted: 29 Jul. 2020; accepted by Maka Murvanidze: 1 Sep. 2020; published: 13 Nov. 2020*