

Forms and Variability of Phosphorus in the Baltic Sea--A Challenge to Ecosystem Modeling

Author: Karlsson, O. Magnus

Source: Air, Soil and Water Research, 3(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/ASWR.S6123>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

OPEN ACCESS

Full open access to this and thousands of other papers at <http://www.la-press.com>.

Forms and Variability of Phosphorus in the Baltic Sea—A Challenge to Ecosystem Modeling

O. Magnus Karlsson

Department of Earth Sciences, Uppsala University, Villavägen 16, SE-752 36 Uppsala, Sweden.

Corresponding author email: magnus.karlsson@ivl.se

Abstract: Knowledge of the partitioning between the dissolved and particulate phases of nutrient elements is a key factor in aquatic ecosystem modeling since partitioning regulates the availability to demand ratio of the nutrient in question. This is seldom taken into account in environmental monitoring programs. In this paper, the occurrence and variability of particulate and dissolved phosphorus were studied in the coastal zone of the Baltic Sea. The particulate fraction (PF) of total phosphorus (TP) concentration in coastal waters from some forty stations along the east coast of Sweden was, on average, 0.33. Dissolved inorganic phosphorus (DIP) was a poor predictor of total dissolved phosphorus (DP) representing only 20%–30% of this fraction. Sensitivity analyses showed that the value of PF had a significant impact on modeled predictions of TP concentration in the water on a Baltic sub-basin scale, whereas an applied coastal model was insensitive to variations in PF. Hence, this study encourages further sampling efforts on the partitioning of phosphorus in the open waters of the Baltic Sea.

Keywords: phosphorus, particulate fraction, ecosystem modeling, Baltic Sea

Air, Soil and Water Research 2010:3 79–93

doi: [10.4137/ASWR.S6123](https://doi.org/10.4137/ASWR.S6123)

This article is available from <http://www.la-press.com>.

© the author(s), publisher and licensee Libertas Academica Ltd.

This is an open access article. Unrestricted non-commercial use is permitted provided the original work is properly cited.

Introduction

Phosphorus is probably the most studied plant nutrient in aquatic sciences. It is often described as the nutrient that limits the growth and biomass of algae.¹ Numerous regressions have been constructed to link phosphorus, especially total phosphorus (TP), to variables such as algal chlorophyll, algal weight and productivity, and Secchi depth.^{2–7} Phosphorus in natural waters can be divided into three component parts: dissolved inorganic phosphorus (DIP), dissolved organic phosphorus (DOP) and particulate phosphorus (PP). The sum of DIP and DOP is termed dissolved phosphorus (DP), and the sum of all phosphorus components combine to give TP. Soluble and particulate phosphorus are operationally differentiated by whether they pass through a 0.45 µm filter.⁸ The “soluble” fraction does not necessarily contain only dissolved phosphorus forms. The filter excludes most particles, but colloidal phosphorus may be present in the filtered fraction.

DIP largely consists of orthophosphate (PO_4). Orthophosphate can be taken up directly by algae, and the concentration of this fraction indicates the amount of phosphorus that is immediately available for algal growth. DIP is quickly regenerated and is therefore a poor predictor of nutrient status in aquatic systems.⁹ A number of organic phosphorus molecules have been identified that would fall within the DOP category but two main classes seem to predominate DOP in natural waters.⁸ The first category includes low molecular weight compounds, which are apparently derived from algal and bacterial metabolism, and are not considered as being directly biologically available. The second group of molecules includes colored large molecular weight compounds, eg, phosphorus bound in humid complexes, which release orthophosphate in the presence of ultraviolet light. These compounds may form a pool of phosphorus available for uptake by biota.^{10–12}

The Baltic Sea has been suffering from eutrophication for decades. Remedial measures have, until recently, been focused on reducing the nitrogen load. However, recent findings using both modeling^{13,14} and empirical approaches^{15,16} have led to a shift in the abatement strategy, whereas the phosphorus input to the Baltic proper is nowadays recognized as important to reduce.^{17,18} Empirical data on the differentiation between PP and DP in the Baltic Sea are scarce,

since most environmental monitoring programs only measure TP and DIP.¹⁴ This fact potentially limits the predictive power of dynamic mass balance models when applied in the Baltic Sea. Comprehensive studies in boreal glacial lakes^{19,20} have found that DP and PP roughly represent 50% each of TP. To use this order of magnitude when differentiating between DP and PP has been successful in modeling phosphorus turnover in the different sub-basins of the Baltic Sea.¹⁴ Nevertheless, since data from the outer Oslo fjord in vicinity of the Baltic Sea,²¹ for example, suggests that as much as 90% of TP is DP, it is warranted to test the hypothesis that the PF value for phosphorus found in lakes also prevails in the brackish Baltic Sea. It is well known that salinity affects the aggregation and sedimentation of particles in estuaries, eg, in relation to the concept of the zone of maximum turbidity.²² It is also possible that salinity influences the partitioning between dissolved and particulate phases of phosphorus, as has been found for trace metals.²³

This study was designed to investigate the occurrence and variability of different physical fractions of phosphorus in Swedish Baltic coastal waters and to investigate the sensitivity associated with the PF value when modeling TP concentrations at different ecosystem scales.

Materials and Methods

Study area

Between 2008 and 2010, 88 samples from surface and bottom water (depth ranging from 5 to 50 meters) were collected from 32 stations (App. 1) situated in the northwestern Baltic proper and the south-western Bothnian Sea (Fig. 1). Samples were collected from all seasons but samples from the production period (May–September) predominated. The study area contains a gradient in water salinity because of the limited exchange of water between the Baltic Sea and the Atlantic, with values ranging from 6 PSU in the south towards 4 PSU in the north.²⁴ Moreover, a gradient in nutrient concentrations is also present because of differences in population densities, with TP levels in the south averaging around 20 µg L⁻¹ decreasing to 10 µg L⁻¹ in the northern part of the study area.¹⁴ In general, river mouths create an east–west gradient in salinity and nutrient levels. As reference material, water samples were collected from the oligotrophic Lake Vättern in central Sweden




Figure 1. Map of the study area with sampling stations marked with dots.

and the Särö archipelago on the Swedish west coast where water salinity is considerably higher (around 20 PSU).

Water sampling and analysis

Water samples were collected with a Ruttner sampler and stored cold in two 1-L plastic bottles. When returned to the laboratory, part of the sampling volume (normally 1 L) was filtered through a 0.45 μm Whatman GF/C glass microfiber filter for later analysis of suspended particulate matter (SPM) and suspended organic matter (SOM). Filters used

for suspended matter analysis had been dried at 550 °C and pre-weighed. Another part of the sampling volume (normally 500 mL) was filtered through a 0.45 μm Whatman cellulose acetate filter for later analysis of particulate phosphorus. The filtrate was saved as well as the remaining unfiltered water volume. The chemical analyses were performed at Erken Laboratory, Uppsala University. The analyses were done according to Swedish standard methods.²⁵

The water salinity and temperature were measured online in the field using an YSI INC Model 30 M handheld salinity, conductivity and temperature

device. The Secchi depth was measured using a standard Secchi disk.

Data Treatment

The definition of PF is:

$$PF = \frac{C_{part}}{C_{tot}} \quad (1)$$

where C_{tot} is the total (unfiltered) concentration and C_{part} is the particulate concentration.²⁶

C_{part} was calculated in two ways: (1) as the measured PP concentration on the filters and (2) as the difference between measured TP concentration in unfiltered water (TP) and filtered water (DP). The DP concentration was directly measured as TP in the filtrate (DP1). DP was also calculated as TP-PP (DP2). Correspondingly, PF was calculated directly from PP/TP (PF1) and from (TP-DP)/TP (PF2).

The variability within the coastal area was studied through analyses of ten replicates each from three areas. The variation was expressed using the coefficient of variation (CV), defined as the ratio between the standard deviation and the arithmetic mean value of a sample.

To investigate how different values of PF for phosphorus may affect the mass balance for TP, and thereby the water concentration of TP in coastal areas and sub-basins of the Baltic Sea, a sensitivity analysis was performed where different values of PF were used and the resulting prediction of the target variable the TP-concentration in the surface water was simulated. This was done using the simulation software Stella[®]. Two types of model for TP turnover were applied on a coastal ecosystem and a sub-basin of the Baltic Sea, respectively. First, a simplified version (Fig. 2a) of a validated mass-balance model for TP-turnover in Baltic coastal areas⁶ was applied in one semi-enclosed coastal area, the Kallrigafjärden Bay. This area has previously been modeled using a similar approach.²⁷ It has an extensive background dataset and typical characteristics of Swedish Baltic coastal areas concerning, for example, water turnover. The other applied model for sensitivity analysis was a crude phosphorus budget for the whole Baltic Sea divided into sub-basins (Fig. 2b) based on nutrient budget calculations presented in Wulff et al.²⁸ and Håkanson and Bryhn.¹⁴

To put the results from the sensitivity analysis of PF into perspective, a similar analysis was performed with the riverine input of TP as the sensitivity parameter. The variability in the yearly riverine load of TP was estimated using a dataset (App. 2) covering 35 rivers during the period 1997–2006 from the national Swedish monitoring program of estuaries provided by the Sveriges Lantbruksuniversitet (SLU; Swedish University of Agriculture).

Results

Basic statistics for the measured water variables are presented in Table 1. Most measured variables had a wide range. For example, TP varied between 9 and 170 $\mu\text{g L}^{-1}$ with a mean value of 25 $\mu\text{g L}^{-1}$, and PP varied between 2 and 108 $\mu\text{g L}^{-1}$ with a mean value of 9 $\mu\text{g L}^{-1}$. The average value of PF calculated from Equation 1 for each sample was 0.33.

A summary of the different DP and corresponding PF values are given in Table 2. The results in Table 2 show that the mean DP concentration was lower (10 $\mu\text{g L}^{-1}$) when measured in the filtrate (DP1) compared to the calculated DP2 (average 15 $\mu\text{g L}^{-1}$). Similarly, the calculated PF values differed. The mean PF1 was 0.33, while the mean PF2 was 0.51. The DIP concentration was considerably lower (mean = 3 $\mu\text{g L}^{-1}$) compared to the DP concentration, indicating that DOP and/or colloidal forms of phosphorus constituted a major part of total DP-concentration in the analyzed samples.

The measured average CV for different water variables are presented in Table 3. DIP had the largest inherent variability, with a CV of 0.28, followed by PP with a CV of 0.16, whereas TP and DP had a rather low CV of 0.07.

Table 4 shows a correlation matrix between different phosphorus fractions and other studied physical and chemical characteristics. A strong positive correlation was found between PP and TP ($r^2 = 0.94$). PP was also positively correlated with SPM and SOM ($r = 0.96$ and 0.93 , respectively). A strong positive correlation ($r = 0.99$) was also found between the SPM and SOM of suspended matter (Table 4). The correlations between TP and PP and SPM and SOM, respectively, after logarithmic transformation are presented in Figure 3.

Considering the mean PF values of 0.33 and 0.51 found in this study depending on the analytical




Figure 2. Outlines of the applied simplified versions of models for **a)** Baltic coastal areas and **b)** the sub-basins of the Baltic Sea. Comprehensive descriptions of the models are given in Håkanson and Eklund (2007), Håkanson and Bryhn (2008) and Karlsson et al (2010). **Abbreviations:** BP, Baltic proper; BS, Bothnian Sea; BB, Bothnian Bay.

technique applied, the global average PF value of 0.56¹⁴ and the reported value of 0.10 from the Oslo fjord,²¹ it was found reasonable to run the sensitivity analysis with the following values of PF: 0.5, 0.25 and 0.75 respectively. The choice of PF value had

a relatively large impact on the prediction of the TP concentration in the surface water of the Baltic basin proper whereas the sensitivity in the modeled coastal area was insignificant ($<0.5 \mu\text{g L}^{-1}$) (Fig. 4). The steady-state value (100 years) of the

Table 1. Basic statistics of measured variables, $n = 60$.

	Water temperature (°C)	Salinity (psu)	Secchi depth (m)	SPM (mg L ⁻¹)	LOI (mg L ⁻¹)	TP (μg L ⁻¹)	PP (μg L ⁻¹)	PO4-P (μg L ⁻¹)	PF
Mean	12	5.4	4.8	4.6	2.4	25	9	2.4	0.33
Median	13	5.2	4.5	3.2	1.4	22	6	1.4	0.33
Sta. Dev.	6	2.4	2.1	8.1	6.3	20	14	6.3	0.17
Max	22	17.2	10.5	65	50	170	108	50	0.66
Min	-0.1	2.9	1.8	1.2	0.5	9	2	0.5	0.06

Table 2. Comparison between different DP- and PF-values, DP1 = measured TP-concentration in the filtrate, DP2 = TP-PP, PF1 = PP/TP, PF2 = (TP-DP1)/TP, n = 63.

Variable	Mean
TP ($\mu\text{g L}^{-1}$)	23
PP ($\mu\text{g L}^{-1}$)	8
DP1 ($\mu\text{g L}^{-1}$)	10
DP2 ($\mu\text{g L}^{-1}$)	15
DIP ($\mu\text{g L}^{-1}$)	3
PF1	0.33
PF2	0.51

TP concentration in the Baltic basin proper was calibrated to $19 \mu\text{g L}^{-1}$ applying a PF value of 0.5. Changing PF to 0.25 increased the predicted TP concentration to $23 \mu\text{g L}^{-1}$, whereas a PF value of 0.75 resulted in a lower TP concentration of $16 \mu\text{g L}^{-1}$ (Fig. 4). In a simulation of Kallrigafjärden Bay, the predicted steady-state (60 months) TP concentration was around $13 \mu\text{g L}^{-1}$ regardless of which PF value that was applied (Fig. 4).

Analysis of the SLU dataset (App. 2) regarding 35 Swedish river estuaries showed that a typical CV for the yearly TP transport is 0.40. A sensitivity simulation with the Baltic sub-basin model using this value to generate 300 normally distributed values of the total riverine TP load within the Baltic Sea resulted in a distribution in the TP concentration of the surface water presented in Figure 5. The confidence interval, calculated as $\pm 1.96 \times$ the standard deviation (SD), for the predicted TP concentration was wide (6, 30). This means that setting up a phosphorus budget in the Baltic Sea using input data from one single year, as often is done in international assessments of the ecological status of the Baltic Sea,²⁹ would contain large uncertainties.

Table 3. Coefficients of variation (CV) for SPM, LOI and different phosphorus fractions based on 10 replicates from three areas.

Variable	CV
SPM	0.13
LOI	0.13
TP	0.07
PP	0.16
DP	0.07
DIP	0.28

Discussion

The mean value of PF for phosphorus found in this study, 0.33, was higher than previously reported from the outer Oslo fjord in the north-east Atlantic,²¹ but was lower than what has been reported from European boreal glacial lakes^{19,20} and the productive Chesapeake Bay on the east coast of the United States.¹⁴ This could possibly be interpreted as an indication that there is a general gradient in PF, with higher values recorded in shallower, nutrient-rich and productive lakes and coastal areas, following a decreasing trend towards the open sea where the SPM concentration generally is lower because of lower primary production and less resuspension.³⁰ It should be mentioned though, that the data set presented in this study is limited and that the values reported should be considered as indicative and seen as a first step towards empirical quantification of phosphorus partitioning in the Baltic Sea.

Water samples were collected all year round but there was a bias in the sampling matrix, with more samples from the production period (May–September) compared to the winter period. When the dataset was disaggregated by season, the PF value was higher during the production period (0.4) compared to the period October to February (0.2). This may be one of the causes of the deviation from the PF value reported by Håkanson and Bryhn,¹⁴ which reflects summer conditions. Moreover, the DIP concentration was higher during the winter months, probably reflecting a mineralization of phosphate from seston that not was directly incorporated into new algae biomasses. This pattern is typical of the seasonal dynamics of nutrients in the Baltic Sea.³¹ No significant differences appeared when comparing samples from surface waters with bottom water samples. This can be explained by the fact that the water mass was relatively uniform in the sampled areas, without any marked thermo- or haloclines. Salinity does not seem to be an important factor for the partitioning of phosphorus, as seen for trace metals.²³

The organic content of SPM was surprisingly uniform, with an average value for SOM of 47% ww and a CV of 24%. This number is very close to what has been reported from fine sedimentary matter caught in traps in lakes³² but higher than found in recent coastal accumulation sediments, which typically have an organic content (SOM) of 15%.³³ The lower

Table 4. Kendall rank correlation coefficients. *r*-values marked bold are significant at $P < 0.05$, $n = 60$.

	Sampling depth (m)	Water temperature (°C)	Salinity (psu)	Secchi depth (m)	SPM (mg L ⁻¹)	SOM (mg L ⁻¹)	TP (µg L ⁻¹)	PP (µg L ⁻¹)	DIP (µg L ⁻¹)	DP (µg L ⁻¹)
Sampling depth (m)	1.00	0.00	-0.06	-0.08	-0.15	-0.13	-0.14	-0.18	0.14	0.01
Water temperature (°C)		1.00	0.13	-0.06	0.24	0.07	-0.31	-0.01	-0.76	-0.17
Salinity (psu)			1.00	0.07	0.49	0.18	-0.12	-0.14	0.07	0.00
Secchi depth (m)				1.00	-0.12	-0.33	-0.66	-0.45	-0.03	-0.62
SPM (mg L ⁻¹)					1.00	0.99	0.91	0.93	-0.10	-0.22
LOI (mg L ⁻¹)						1.00	0.94	0.96	-0.08	-0.23
TP (µg L ⁻¹)							1.00	0.94	0.12	0.70
PP (µg L ⁻¹)								1.00	-0.09	-0.07
DIP (µg L ⁻¹)									1.00	0.49
DP (µg L ⁻¹)										1.00




Figure 3. Observed correlations between log TP and log PP and log SPM and log LOI, respectively, ($n = 60$).




Figure 4. Predicted TP-concentrations (mg m⁻³) in surface waters when varying the PF-values in a) the Kallrigafjärden Bay and b) the Baltic proper.




Figure 5. Sensitivity analysis using the Baltic Sea sub-basin model. The target variable is TP ($\mu\text{g/l}$) in the Baltic Proper and the TP-loading has been varied ($\text{CV} = 0.40$, $n = 300$). The box-and-whisker plot shows the mean, ± 1 standard deviation and $\pm 1.96 \cdot$ standard deviation.

organic content in recent surface sediments compared to the suspended matter in the overlying water column can partly be explained by mineralization of easy degradable organic matter and partly by dilution with minerogenic matter from resuspension processes occurring during strong wind events.^{34,35} The average value of the SPM concentration found in this study (4 mg L^{-1}) is of the same order of magnitude as earlier reported in Pustelnikov³⁶ (average 3.3 mg L^{-1}) and Håkanson³⁰ (average 2.5 mg L^{-1}).

As seen from Table 2, a difference appeared between DP measured directly in the filtrate and DP calculated from TP-PP, making it possible to calculate PF in two ways. So the question is: which analytical method is best for determining PF? Measuring the particulate fraction as the matter collected on the filter is the traditional method of determining the particulate fraction.^{37,38,25,19,20} It has the advantage that a rather large volume of water can be filtered, which reduces the risk of getting samples with concentrations of the actual substance that are below the detection limit of the analytical method. This can be a problem when performing analyses on the filtrate, especially with substances with high affinity with particles. On the other hand, the analytical methods are consistent

when identical analyses are performed on unfiltered and filtered water, respectively. The results from this study also indicate that the inherent variability in phosphorus analyses may be lower for water samples compared to filter analyses (Table 3).

DIP, which is a standard parameter in environmental monitoring programs, was a poor predictor of DP since, on average, it only represented 20%–30% of total DP (Table 2). Hence, the data presented in this study suggests that DIP cannot be used as an estimator of the non-settling phase of phosphorus in mass balance modeling. Similar conclusions have been drawn by Håkanson and Bryhn¹⁴ and Håkanson and Stenström-Khalili.³⁹

The rather strong correlations found in this study (eg, between TP and PP and SPM and LOI, respectively (Table 4, Fig. 3)) is, to a large extent, explained by the wide range in the dataset, which, in turn, depends on the fact that one sampling excursion was undertaken during a significant algae bloom. If this data point was removed from the dataset, the r -values in Table 4 dropped markedly but the correlation was still significant ($P < 0.05$). Nevertheless, these regressions should be treated as indicative and they need to be further elaborated.

Varying the PF value had a marked impact on the prediction of TP concentration in the Baltic basin proper but not on the prediction of the TP concentration in the coastal estuary of Kallrigafjärden Bay (Fig. 4). The main reason for this is the generally fast water turnover in coastal areas, which gives internal processes, eg, sediment–water exchange, little importance for the nutrient mass balance. On the other hand, in the basin of the Baltic proper and the other large basins of the Baltic Sea where the water turnover is slow (within the range of decades), the settling of particles, although also a slow process, is important to the overall mass balance. A similar sensitivity to PF value when modeling on a Baltic Sea sub-basin scale is indicated in Håkanson and Bryhn,¹⁴ whereas other Baltic Sea models (eg,^{13,40,41}) have a different parameterization of the sedimentation processes, making model comparisons irrelevant.

Not surprisingly, the variation in the riverine load of TP had a large impact on the modeled TP concentration in the Baltic basin proper (Fig. 5). A similar sensitivity to the allochthonous input has

also been recognized in various lake ecosystem models.^{32,42} The calculated CV for the yearly riverine load of TP to the Baltic Sea of 0.40 infers that an average of 16 years, transport data, at least, a required to establish a typical mean value for the riverine load if one accepts an error of 20%.⁴³ Wulff et al⁴⁴ advocated that the conditions in the Baltic Sea are slowly getting less eutrophic, based on calculations from riverine load data for the years 1994–2006. The calculations above show that such a conclusion, drawn from a rather short period, may be premature.

Conclusions

The indicative PF value obtained in this study is lower but of the same magnitude as that reported earlier in boreal lakes. PP measured as the remains on the filter after filtering is not equivalent with PP measured as the difference between TP in unfiltered and filtered water. DIP was a poor predictor of DP, representing only a small fraction of the total dissolved phase. DIP also had the largest inherent variability of measured phosphorus fractions. A strong correlation was found between TP and particulate phosphorus (PP) in the sampling matrix. This study demonstrates that an accurate PF value is necessary for obtaining high predictive power in phosphorus mass balance models of the Baltic Sea. Since the empirical basis for this is weak, more attention should be given to determining of the particulate fraction of phosphorus in future sampling programs of open waters of the Baltic Sea.

Disclosures

This paper is unique and not under consideration by any other publication, and has not been published elsewhere. The author and peer reviewers report no conflicts of interest. The author confirms that he has permission to reproduce any copyrighted material.

Acknowledgements

The author thanks Erken Laboratory for performing the chemical analyses, the Swedish University of Agriculture for sharing river transport data and Dan Lindgren for preparing the map. I also thank Gunnar Andersson, Per Jonsson, Staffan Karlsson, Gustaf Karlsson Medin, Mikael Malmaeus, Ingvar Niklasson, Emil Rydin, Krister Seleskog and Anders Wadman for assistance in the field work.

References

1. Schindler DW, Hecky RE, Findlay DL, et al. Eutrophication of lakes cannot be controlled by reducing nitrogen input: Results of a 37-year whole-ecosystem experiment. *Proceedings of the National Academy of Sciences*. 2008; 105(32): 11254–8.
2. Vollenweider RA. Sichttiefe und Production. *Verhandlungen der Internationalen Vereinigung der Limnologie*. 1958;13:142–3.
3. Dillon PJ, Rigler FH. The phosphorus-chlorophyll relationship in lakes. *Limnology and Oceanography*. 1974;19:767–73.
4. Evans MS, Arts MT, Robarts RD. Algal productivity, algal biomass, and zooplankton biomass in a phosphorus-rich, saline lake: deviations from regression model predictions. *Canadian Journal of Fisheries and Aquatic Science*. 1996;53:1048–60.
5. Sagert S, Jensen, DK, Henriksen P, Rieling T, Schubert H. Integrated ecological assessment of Danish Baltic Sea coastal areas by means of phytoplankton and macrophytobenthos. *Estuarine, Coastal and Shelf Science*. 2005;63: 109–18.
6. Håkanson L, Eklund JM. A dynamic mass-balance model for phosphorus fluxes and concentrations in coastal areas. *Ecol Res*. 2007;22:296–320.
7. Malmaeus JM, Karlsson OM, Lindgren D, Eklund J. The optimal size of dynamic phosphorus models for Baltic coastal areas. *Ecological Modelling*. 2008;216:303–15.
8. Carlson RE, Simpson J. A coordinator's guide to volunteer lake monitoring methods. *North American Lake Management Society*. 1996.
9. Dodds WK. Misuse of inorganic N and soluble reactive P concentrations to indicate nutrient status of surface waters. *Journal of the North American Benthological Society*. 2003;22:171–81.
10. Huang B, Hong H. Alkaline phosphatase activity and utilization of dissolved organic phosphorus by algae in subtropical coastal waters. *Marine Pollution Bulletin*. 1999;39:205–11.
11. Vidal M, Duarte CM, Agusti S. Dissolved organic nitrogen and phosphorus pools and fluxes in the Central Atlantic Ocean. *Limnology and Oceanography*. 1999;44:106–15.
12. Nausch M, Nausch G. Bioavailability of dissolved organic phosphorus in the Baltic Sea. *Marine Ecology Progress Series*. 2006;321:9–17.
13. Savchuk OP, Wulff F. Modelling regional and large-scale response of Baltic Sea ecosystems to nutrient reductions. *Hydrobiologia*. 1999;393: 35–43.
14. Håkanson L, Bryhn AC. *Eutrophication in the Baltic Sea*. Berlin: Springer Verlag; 2008.
15. Larsson U, Hajdu S, Walve J, Elmgren R. Baltic Sea nitrogen fixation estimated from the summer increase in upper mixed layer total nitrogen. *Limnology and Oceanography*. 2001;46:811–20.
16. Wasmund N, Nausch G, Schneider B, Nagel K, Voss M. Comparison of nitrogen fixation rates determined with different methods: a study in the Baltic Proper. *Mar Ecol Prog Ser*. 2005;297:23–31.
17. Boesch D, Hecky R, O'Melia C, Schindler DW, Seitzinger S. Eutrophication of Swedish Seas. *Swedish Environmental Protection Agency*. Technical report; 2006. No. 5509, Sweden.
18. SEPA. Övergödningen av Sveriges kuster och hav (In Swedish). *Swedish Environmental Protection Agency*. Technical report; 2006. No. 5587, Sweden.
19. Håkanson L, Boulion V. *The Lake Foodweb*. Leiden: Backhuys Publishers; 2002.
20. Bryhn AC, Hessen DO, Blenckner T. Predicting particulate pools of nitrogen, phosphorus and organic carbon in lakes. *Aquatic Sciences*. 2007;69: 484–94.
21. Aure J, Danielssen DS. Ytre Oslofjord. Hydrografi og næringsalter over terskeldyp. 1995–1998 (In Norwegian). *Havforskningsinstituttet*. Technical report; 1999. No. 785, Norway.
22. Bowden KF. Turbulence and mixing in estuaries. In: Kennedy VS, editor. *The Estuary as a Filter*. Orlando: Acad. Press; 1984:15–26.
23. Turner A. Trace-metal partitioning in estuaries: importance of salinity and particle concentration. *Marine Chemistry*. 1996;54:27–39.
24. Kullenberg G. 1981. Physical oceanography. In: Voipio A. ed. *The Baltic Sea*. Amsterdam: Elsevier Oceanography Series; 1981;30:135–75.
25. SS-EN 872:2005 for SPM, SS-EN 15169:2007 (modified for water) for SOM, SS-EN ISO 6878:2005 for DIP, SS 02 81 27 for TP and SS-EN ISO 15681-1:2005 for PP.



26. Johansson H, Lindström M, Håkanson L. On the modelling of the particulate and dissolved fractions of substances in aquatic ecosystems—sedimentological and ecological interactions. *Ecological Modelling*. 2001; 137:225–40.
27. Karlsson OM, Malmaeus JM, Josefsson S, Wiberg K, Håkanson L. Application of a mass-balance model to predict PCDD/F turnover in a Baltic coastal estuary. *Estuarine, Coastal and Shelf Science*. 2010;88: 209–18.
28. Wulff F, Rahm L, Hallin AK, Sandberg J. A nutrient budget model of the Baltic Sea. In: Wulff F, Rahm L, Larsson P. eds. *A Systems Analysis of the Baltic Sea*. Heidelberg: Springer-Verlag Ecological studies. 2001;148:353–73.
29. HELCOM. Ecosystem Health of the Baltic Sea. *Helsinki Commission*. Technical report; 2010. No. 122, Finland.
30. Håkanson L. *Suspended particulate matter in lakes, rivers and marine systems*. New Jersey: The Blackburn Press; 2006.
31. Eilola K. On the dynamics of organic nutrients, nitrogen and phosphorus, in the Baltic Sea. *Swedish Meteorological and Hydrological Institute*. Technical report; 2009. No. 99, Sweden.
32. Håkanson L, Peters RH. 1995. *Predictive limnology*. Amsterdam: SPB Academic Publishing; 1995.
33. Håkanson L, Rosenberg R, 1985. Praktisk kustekologi (In Swedish). *Swedish Environmental Protection Agency*. Technical report; 1985. No. 1987, Sweden.
34. Eckhell J, Jonsson P, Meili M, Carman R. Storm influence on the Accumulation and Lamination of Sediments in Deep Areas of the Northwestern Baltic Proper. *Ambio*. 2000;29:238–45.
35. Persson J, Jonsson P. Historical development of laminated sediments—an approach to detect soft sediment ecosystem changes in the Baltic Sea. *Marine Pollution Bulletin*. 2000;40:122–34.
36. Pustelnikov OS. Geochemical features of suspended matter in connection with recent processes in the Baltic Sea. *Ambio*. 1977;5:157–62.
37. Benoit G, Oktay-Marshall SD, Cantu A, et al. Partitioning of Cu, Pb, Ag, Zn, Fe, Al and Mn between filter-retained particles, colloids, and solution in six Texas estuaries. *Mar Chem*. 1994;45:307–36.
38. Pohl C, Hennings U. The effect of redox processes on the partitioning of Cd, Pb, Cu and Mn between dissolved and particulate phases in the Baltic Sea. *Mar Chem*. 1999;65:41–53.
39. Håkanson L, Stenström-Khalili MI. Uncertainties in Data and Spurious Correlations Related to the Redfield Ratio. *International Review of Hydrobiology*. 2009;94:338–51.
40. Vichi M, Ruardij P, Baretta JW. Link or sink: a modelling interpretation of the open Baltic biogeochemistry. *Biogeosciences Discussions*. 2004;1: 79–100.
41. Schernewski G, Neumann T. The trophic state of the Baltic Sea a century ago: a model simulation study. *Journal of Marine Systems*. 2005;53: 109–24.
42. Bryhn AC, Girel C, Paolini G, Jacquet S. Predicting future effects from nutrient abatement and climate change on phosphorus concentrations in Lake Bourget, France. *Ecological Modelling*. 2010;221:1440–50.
43. Håkanson L. Sediment sampling in different aquatic environments, statistical aspects. *Water Resour Res*. 1984;20:41–6.
44. Wulff F, Humborg C, Rodriguez Medina M, Mörtz CM, Savchuk O, Sokolov A. Revision of the country allocation of nutrient reductions in the Baltic Sea Action Plan—Section A: Hydrological adjusted riverine loads and atmospheric loads from different countries averaged for 2000–2006. *Baltic Nest Institute*. Technical Report; 2009. Sweden.



Supplementary Data

Data from the sampling sites

Station	Sampling depth (m)	Water temperature (°C)	Salinity (PSU)	Secchi depth (m)	SPM (mg L ⁻¹)	LOI (mg L ⁻¹)	TP (µg L ⁻¹)	PP (µg L ⁻¹)	PO4-P (µg L ⁻¹)	DP (µg L ⁻¹)
Gräsö	8	10.8	4.7	7	4.5	1.8	10	4	2	8
Örskär	15	10.9	4.5		4.5	1.8	9	4	0.1	6
Öregrundsgrepen	3.5	12.3	4.5	6	5.5	2.2	11	6	0.3	7
Kalligafjärden	1.5	14.5	3.4	2.5	6.7	2.7	22	12	0.5	8
Tvären sw	5	15	5		2.5	1.2	26	6	9	23
Tvären dw	40	12	5.5		2.7	1.2	26	6	7	16
Höggarnsfjärden dw	20	15.2	2.9	2.5	2.9	1.3	23	5	2	13
Höggarnsfjärden sw	5	14.7	2.9	2.5	3.7	1.4	27	5	1	9
Siarö sw	5	17	4.8		3.7	1.8	20	8	1	5
Siarö dw	30	16	4.8		3	1.4	18	9	1	5
Kallskär sw	10	17	4		2.9	1.5	13	5	1	7
Kallskär dw	50	15	4		3.2	1.4	10	5	1	5
Svartlögfjärden sw	2	17.5	3.7		3.4	1.7	14	8	2	6
Svartlögfjärden dw	20	17.1	3.7		3.5	1.7	16	8	4	5
Välårö sw	5	16.5	5.1		5.5	2.2	27	10	1	10
Välårö dw	15	16.4	5.1		4	1.7	20	8	8	14
Epskär	5	13	5.2	5	7.3	3	30	14	2	14
Gustaf Dahлёn 1 dw	50	19.5	5.4	5	3.2	1.9	17	7	1	8
Gustaf Dahлёn 1 sw	25	20.2	5.2	5	3.4	2	18	8	1	11
Gustaf Dahлёn 2 dw	30	19.9	5.5	5	2.8	1.7	22	10	1	9
Gustaf Dahлёn 2 sw	10	19.7	5.4	5	3	1.5	19	8	2	8
Gustaf Dahлёn 3	15	19.4	5.5	6	2.5	1.4	19	10	1	9
Gustaf Dahлёn 4	5	21.1	5.1	5	2.5	1.1	24	11	1	10
Gustaf Dahлёn 5	2	21.7	5.2	2.5	5	1.7	35	12	8	17
Gustaf Dahлёn algae	0.5	21.5	5.2	0.5	64.5	50	170	108	3	
Öregrundsgrepen	5	16.3	4.1	3.5	4.1	1	17	4	1	6
Kalligafjärden	3	16.3	3.4	2.3	2.4	1.1	27	7	1	9
Ägnöfjärden sw	5	15.2	5	4.5	2	0.9	22	4	1	11
Ägnöfjärden dw	25	14	5.1	4.5	1.5	0.9	22	2	3	16
Trälhavet sw	5	13.7	3.6	4.5	3.2	1.3	20	5	1	9
Trälhavet dw	35	13.7	3.6	4.5	2.7	1.2	27	5	1	9
Furusund*	5	6.2	5.5	6	3.35	1.68	15	5	1.7	6
Stussviken*	5	11.1	6.2	6.5	2.09	1.16	22	6	6	12
Särö 1	5	21.6	17.2	6	7.8	3.2	15	5	1	7
Särö 2	5	21.6	16.4	8	11.8	2.5	11	4	1	6
Lake Vättern				17	2.3	0.5	5	3	<1	<5
Gårdsfjärden*	0.5		4	2.5	4.48	1.77	30	7	4.9	9

(Continued)



Station	Sampling depth (m)	Water temperature (°C)	Salinity (PSU)	Secchi depth (m)	SPM (mg L ⁻¹)	LOI (mg L ⁻¹)	TP (µg L ⁻¹)	PP (µg L ⁻¹)	PO4-P (µg L ⁻¹)	DP (µg L ⁻¹)
Edeboviken sw	2	7.1	5.9	4	1.8	1.1	23	7	6	
Edeboviken dw	9	7.1	5.9		2.2	1.1	22	6	5	
Källsöfjärden sw	13	7.2	6.2		1.9	0.9	22	6	7	
Källsöfjärden dw	2	7.3	6.14	4	3.3	1.2	22	6	7	
Medholmsviken sw	2	7.3	6.2	4.5	2	0.8	19	7	6	
Medholmsviken dw	8	7.3	6.2		2.4	0.9	18	5	6	
Hargsviken sw	6	6	6		2.1	1.2	23	8	5	
Hargsviken dw	2	6	6	4.2	3.3	1.2	23	8	4	
Singöfjärden sw	5	-0.1	5.3		3.4	1.5	20	3	11	
Singöfjärden dw	47	0.1	5.3		3.2	1.3	21	3	12	
Östhammarsfjärden sw	2	6.1	4.5	1.8	5.1	5.1	46	29	8	
Östhammarsfjärden dw	7	6.1	4.5		6.2	5	50	28	8	
Kobbjärden	5	7.4	5.8	10.5	2.6	1.3	21	3	10	
Kudoxafjärden	5	7.1	5.9	9	1.4	0.7	20	3	9	
Gräsköfjärden	5	7.2	5.8	9	1.5	0.7	19	4	8	
Edöfjärden sw	5				1.2	0.5	24	2	14	
Edöfjärden sw	5				2.8	1.1	32	3	21	
Edöfjärden dw	36				3.8	1.5	23	3	14	
Stussviken sw	5	0.3	6.1		2.3	0.9	28	2	18	
Stussviken dw	25	0.3	6.4		2.6	0.9	28	2	19	
Älgöfjärden sw	5	3.5	3.9	3	4.3	2	28	18.4	9	
Älgöfjärden dw	35	1.3	5.7		3.6	1.3	31	5.72	21	
Edöfjärden sw	5	3.8	4.3	3	6.4	3.3	32	21	9	
Edöfjärden dw	35	0.6	5.7		3.4	1.2	27	4.08	17	

Note: *Average of 10 replicates.



Replicate sampling at three stations.

Station	Sampling depth (m)	Water temperature (°C)	Salinity (PSU)	Secchi depth (m)	SPM (mg L ⁻¹)	LOI (mg L ⁻¹)	TP (µg L ⁻¹)	PP (µg L ⁻¹)	PO4-P (µg L ⁻¹)	DP (µg L ⁻¹)
Furusund	5	6.2	5.5	6	3.9	1.9	14	6	2	7
Furusund	5	6.2	5.5	6	3.5	1.7	14	5	1	6
Furusund	5	6.2	5.5	6	3.1	1.5	14	5	2	7
Furusund	5	6.2	5.5	6	3.4	1.6	14	6	1	6
Furusund	5	6.2	5.5	6	3.2	1.4	19	6	1	6
Furusund	5	6.2	5.5	6	3.6	1.8	15	5	2	6
Furusund	5	6.2	5.5	6	2.9	1.5	14	5	3	6
Furusund	5	6.2	5.5	6	3.3	1.8	17	5	1	6
Furusund	5	6.2	5.5	6	3.3	1.9	16	5	2	6
Furusund	5	6.2	5.5	6	3.3	1.7	16	5	2	6
Mean	5.0	6.2	5.5	6.0	3.3	1.7	15.3	5.3	1.7	6.2
Gårdsfjärden	5	13.6	4	2.5	4.3	2	30	10	4	10
Gårdsfjärden	5	13.6	4	2.5	3.5	1.3	28	12	4	9
Gårdsfjärden	5	13.6	4	2.5	3.9	1.6	28	6	5	8
Gårdsfjärden	5	13.6	4	2.5	3.3	1.6	31	5	4	8
Gårdsfjärden	5	13.6	4	2.5	4.2	1.8	31	5	4	8
Gårdsfjärden	5	13.6	4	2.5	4.7	1.9	31	6	4	9
Gårdsfjärden	5	13.6	4	2.5	5.2	1.9	31	7	7	9
Gårdsfjärden	5	13.6	4	2.5	4.8	1.6	26	6	5	8
Gårdsfjärden	5	13.6	4	2.5	5.7	2	28	7	6	9
Gårdsfjärden	5	13.6	4	2.5	5.2	2	32	7	6	10
Mean	5.0	13.6	4.0	2.5	4.5	1.8	29.6	7.2	4.9	8.8
Stussviken	5	11.1	6.2	6.5	2.1	1.2	22	6	5	12
Stussviken	5	11.1	6.2	6.5	2.1	1.1	21	6	6	13
Stussviken	5	11.1	6.2	6.5	2.5	1.4	21	6	6	13
Stussviken	5	11.1	6.2	6.5	1.8	1	22	5	6	13
Stussviken	5	11.1	6.2	6.5	2.4	1.5	22	6	6	12
Stussviken	5	11.1	6.2	6.5	1.7	0.9	21	6	5	12
Stussviken	5	11.1	6.2	6.5	1.9	1.1	21	6	5	13
Stussviken	5	11.1	6.2	6.5	2.3	1.1	24	6	9	12
Stussviken	5	11.1	6.2	6.5	1.9	1.1	21	6	5	12
Stussviken	5	11.1	6.2	6.5	2.2	1.2	22	5	7	12
Mean	5.0	11.1	6.2	6.5	2.1	1.2	21.7	5.8	6.0	12.4

TP-transport (tonnes yr⁻¹) in 35 Swedish rivers from the SLU-database, calculated standard deviations and CV-values for selected periods.

	Mean transport (tonnes yr ⁻¹) 1965–2006	Stdav 1965–2006	CV 1965–2006	Mean transport (tonnes yr ⁻¹) 1997–2006	Stdav 1997–2006	CV 1997–2006
Norrström	230	253	1.10	170	65	0.38
Nyköpingsån	26	11	0.43	29	12	0.43
Motala ström	99	36	0.37	112	38	0.34
Botorpström	4	2	0.42	4	1	0.23
Emån	21	8	0.37	19	5	0.25
Alsterån	5	2	0.33	5	1	0.21
Ljungbyån	7	5	0.71	4	1	0.28
Gothemsån	7	3	0.49	7	4	0.63
Lyckebyån	5	2	0.38	5	2	0.32
Mörrumsån	20	7	0.37	24	7	0.29
Helgeån	53	19	0.37	56	21	0.38
Skivarpsån	6	4	0.69	4	2	0.43
Kävlingeån	28	11	0.38	29	10	0.35
Råån	6	3	0.44	6	2	0.32
Forsmarksån	2	1	0.66	2	0	0.30
Dalälven	188	59	0.31	173	67	0.39
Gavleån	21	9	0.43	16	6	0.35
Ljusnan	114	48	0.42	85	43	0.50
Delångersån	7	2	0.35	6	2	0.40
Ljungan	62	84	1.35	94	162	1.73
Indalsälven	124	38	0.31	100	52	0.52
Ångermanälven	167	54	0.32	138	45	0.32
Gideälven	21	6	0.31	19	7	0.35
Lödge älv	17	7	0.40	17	7	0.38
Öre älv	29	12	0.40	30	14	0.45
Umeälven	165	49	0.30	146	52	0.35
Rickleån	10	3	0.32	10	4	0.37
Skellefteälven	50	13	0.27	39	13	0.33
Piteälven	82	30	0.36	78	37	0.47
Alterälven	5	1	0.26	4	1	0.31
Luleälven	162	62	0.38	120	29	0.24
Råne älv	24	7	0.28	22	7	0.30
Töreälven	5	2	0.52	4	1	0.26
Kalixälven	220	72	0.33	200	78	0.39
Torne älv	318	115	0.36	328	124	0.38
Mean			0.44			0.40



Publish with Libertas Academica and every scientist working in your field can read your article

"I would like to say that this is the most author-friendly editing process I have experienced in over 150 publications. Thank you most sincerely."

"The communication between your staff and me has been terrific. Whenever progress is made with the manuscript, I receive notice. Quite honestly, I've never had such complete communication with a journal."

"LA is different, and hopefully represents a kind of scientific publication machinery that removes the hurdles from free flow of scientific thought."

Your paper will be:

- Available to your entire community free of charge
- Fairly and quickly peer reviewed
- Yours! You retain copyright

<http://www.la-press.com>