

Memory Functions in Recreational Pistol Sport Shooters: Does Lead Matter?

Authors: Asa-Mäkitaipale, Sanna, Jehkonen, Mervi, Uitti, Jukka, and Vilkki, Juhani

Source: Environmental Health Insights, 3(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/EHI.S894>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Memory Functions in Recreational Pistol Sport Shooters: Does Lead Matter?

Sanna Asa-Mäkitaipale^{1,2}, Mervi Jehkonen³, Jukka Uitti^{4,5} and Juhani Vilkki²

¹South Karelian Central Hospital, Department of Neurology, Lappeenranta, Finland. ²University of Helsinki, Department of Psychology, Helsinki, Finland. ³University of Tampere, Department of Psychology, Tampere, Finland. ⁴Finnish Institute of Occupational Health, Tampere, Finland. ⁵Clinic of Occupational Medicine, Tampere University Hospital, Tampere, Finland.

Abstract

Objective: The aim of our study was to examine the memory functions of pistol sport shooters using powder charges when exposure to lead is expected to be considerably lower than in occupational circumstances.

Methods: A neuropsychological battery of memory and intelligence tests was administered to 20 sport shooters and 20 controls whose mean ages (SDs) were 55 (9.6) and 54 (9.3) years respectively. Memory functions were evaluated with three subtests of the Wechsler Memory Scale - Revised (WMS-R) and an incidental memory test. Intelligence was assessed with four subtests of the Wechsler Adult Intelligence Scale - Revised (WAIS-R). The level of alcohol consumption and depression were examined in both groups. Blood lead level was determined among the shooters.

Results: The shooters performed worse than the controls in the tests of incidental and logical memory. The groups did not differ in intelligence, mood or alcohol consumption. The mean (SD) blood lead level of the sport shooters was 0.52 µmol/L (0.40), responding 10.76 µg/dl (8.28).

Conclusions: Low lead exposure in recreational shooting conditions may impair verbal memory. Therefore it is important to ensure that lead exposure is prevented among those shooting for sport.

Keywords: cognitive function, lead exposure, sport shooting

Introduction

Shooting with lead-containing bullets on indoor firing ranges is known to expose people to lead.^{1–3} On being fired an unjacketed lead bullet emits some lead fumes and dust which become airborne due to the gun's muzzle blast.^{4,5} The primary exposure of lead occurs when the shooter inhales the fumes in the proximity of the gun.⁶ Lead exposure in indoor firing ranges can also occur through inhalation of lead fumes and particulates suspended in the range air. The lead exposure is also dependent on the structure of the indoor range and the ventilation systems.⁵ Lead exposure can be examined in blood lead level, which is known to be elevated among trainees, instructors, police officers and controllers^{1–3,5,6} and among recreational shooters.^{7,8} Lead exposure is more intensive in closed shooting ranges than in open ranges, where shooting is normally not considered to be hazardous to sport shooters' health.⁵

Numerous studies have demonstrated that occupational lead exposure is associated with low neuropsychological test performance.^{9–23} Long-term environmental exposure to high levels of lead has persistent detrimental effects on cognitive functions in older adults.²⁴ High lead exposure may damage the central nervous system.²⁵ High tibial lead is associated with small total brain volume.^{26,27} Lead may influence cognitive functions by decreasing of brain volume.²⁸

Lead may affect memory, executive functions, reasoning, visual and motor functions.^{12–20,21} Memory functions are known to be vulnerable, not least to lead exposure which may have effects both on verbal and visual memory.^{14,18–20} Lead may impair verbal memory even at occupational exposures and have progressive effects later when exposure has finished.²⁰ Chronic lead exposure affects encoding, storage and retrieval of verbal information.²³ Lead may impair the ability to remember large verbal entities,^{14,18} verbal short-term or long-term memory.^{19,20,23} In visual memory, the effects of lead exposure can be

Correspondence: Mervi Jehkonen, University of Tampere, Department of Psychology, FIN-33014 University of Tampere, Finland. Fax: +358 3 3551 7345; Email: mervi.jehkonen@uta.fi



Copyright in this article, its metadata, and any supplementary data is held by its author or authors. It is published under the Creative Commons Attribution By licence. For further information go to: <http://creativecommons.org/licenses/by/3.0/>.

seen later, with long-term effects.²⁰ The effects of lead have been found in visual retention memory.^{18,20} So far it has not been demonstrated at what level of lead in the blood the cognitive functions begin to be impaired in adults.²⁹ In children blood lead concentrations below 10 µg/dl (responding 0.48 µmol/l) are associated with impaired intelligence,^{30–32} even as low blood lead levels as 7.5 µg/dl (responding 0.36 µmol/l) have been found to decrease children's intellectual functions because of environmental exposure.³³

The aim of this study was to examine sport shooters' memory functions since lead exposure is known to affect memory. To the best of our knowledge there is very little evidence of the neuropsychological effects caused by lead exposure in sport shooters using powder charges. We wanted to examine if lead impairs the memory functions of sport shooters as evaluated in clinical neuropsychological memory tests. We assumed that sport shooters would perform worse than controls in memory tests as a result of lead exposure.

Participants and Methods

Participants

The study included 20 sport shooters and 20 controls. The study population consisted of pistol sport shooters who practised on indoor or outdoor firing ranges, mainly withunjacketed lead bullets. All sport shooters from two local shooting clubs

who volunteered to participate in the study were contacted. The sport shooters had to meet the following inclusion criteria: a) at least 10 years of continuous sport shooting with lead bullets in indoor or outdoor shooting ranges, and b) male gender. The control group consisted of volunteers, who were similar to the shooters regarding to gender, age and years of education. The controls did not have any known lead exposure. The eligibility of respondents was confirmed by a subsequent questionnaire. The potential participants were excluded if any of the following criteria was met: a) neurological and/or psychiatric disease, b) alcohol or drug abuse, or c) other exposure to heavy metals and/or solvents.

Methods

Each eligible respondent participated in a single two-hour neuropsychological examination. The participants completed questionnaires on personal background, alcohol consumption and shooting habits. The analysis of blood lead level was carried out only for the sport shooters.

Neuropsychological examination

A trained examiner blind to participants' exposure administered neuropsychological tests which have been found to be useful in occupational studies. The clinical tests are shown in Table 1. Reasoning was examined with the Similarities, Digit Symbol, Block Design and Picture Completion subtests of the

Table 1. Neuropsychological tests.

Cognitive domain	Tests	Possible range
Verbal reasoning	Similarities ¹	0–34
Visual reasoning	Digit Symbol ¹	0–93
	Block Design ¹	0–51
	Picture Completion ¹	0–22
	Digit Span (a + b) ¹	0–28
Verbal memory	a) forward: 0–14	
	b) backward: 0–14	
	Logical Memory (Story A) ²	0–25
	Verbal Learning ²	0–24
	Incidental memory test ³	0–20
Visual memory	Visual Retention ²	0–41

Notes: ¹The subtests of the Wechsler Adult Intelligence Scale - Revised. ²The subtests of the Wechsler Memory Scale - Revised. ³The incidental memory test,³⁶ see Methods.

Wechsler Adult Intelligence Scale—Revised³⁴ (WAIS-R). Memory was tested with the Logical Memory (Story A), Verbal Learning and Visual Retention subtests of the Wechsler Memory Scale - Revised³⁵ (WMS-R), with an incidental memory test³⁶ and with the Digit Span subtest of the WAIS-R.³⁴

In the incidental memory test³⁶ the Similarities Subtest of the WAIS-R was used as an orienting task. After the participant had answered the first 10 questions, in which he tried to discover the similarity between two words, he was asked, without previous warning, to recall the word pairs or words included in the questions in any order. After the free recall, the first word of each pair in which the participant did not recall both of the words in the free recall was given as a cue for the recall of the second word. The total number of correct words in the free and cued recall formed the score.

The level of depression was assessed with the Beck Depression Inventory³⁷ (BDI). The range of its total score was 0–63. Alcohol consumption was examined by the Alcohol Use Disorders Identification Test³⁸ (AUDIT) ranging from 0 to 40.

Analysis of blood lead level

Sport shooters participated in a blood lead (PbB) analysis. Blood samples were collected one week after the neuropsychological examination. An amount of 5–10 mL of whole blood in heparinised vacuum tube for trace element analysis was collected. Blood lead concentration was determined with an electrothermal atomic absorption spectrophotometric method³⁹ in the biomonitoring laboratory of the Finnish Institute of Occupational Health.

Data analysis

Statistical Package for the Social Sciences (SPSS-PC version 11.5) was used for the data analysis.

In the background variables (age, years of education, the BDI and the AUDIT) the group differences were studied with t-tests for independent groups. In the intelligence tests and in the immediate memory tests, the sums of raw scores were used, but in the delayed recalls of the memory tests percent variables. In the tests of logical memory and visual retention the percent variable was the delayed recall score as a percentage of the immediate recall score. In the verbal learning test the delayed recall score as a percentage of the third (last) immediate recall score was used. The difference between the two groups was tested with a multivariate analysis of covariance (MANCOVA) separately on the four intelligence and the nine memory variables, with age as a covariate. Correlations between the psychological test variables, the blood lead level and shooting habits were studied with Pearson's correlation coefficients. The significance level of statistical analysis was set at $p \leq 0.05$.

Results

The sport shooters did not differ from the controls on the background variables (Table 2). The sport shooters had spent an average of seven hours on indoor or outdoor firing ranges during the last month. The range of years of sport shooting was 11 to 46 (mean = 27.4, SD = 11.6).

The results of the blood lead analysis and the amount of bullets fired during the last month are given in Table 3.

The mean blood lead level of the sport shooters was 0.52 $\mu\text{mol/L}$ (responding 10.76 $\mu\text{g/dl}$), which is higher than the normal level 0.3 $\mu\text{mol/L}$ (responding 6.21 $\mu\text{g/dl}$) of non-exposed people.^{40,41} The shooters' blood lead level was higher the more bullets they had fired during the last month or during the last year ($r = 0.705$, $p = 0.001$; $r = 0.551$, $p = 0.012$).

Table 2. The characteristics of the participants.

	Sport shooters	Controls	Comparison of groups	
	Mean [SD; Range]	Mean [SD; Range]	t-value	p
Age (years)	55.0 [9.6; 30–68]	54.0 [9.3; 30–67]	0.44	0.67
Years of education	12.7 [3.6; 7–20]	13.3 [4.4; 8–23]	0.49	0.63
BDI	5.6 [4.7; 0–17]	5.1 [4.0; 0–12]	0.32	0.75
AUDIT	4.7 [3.4; 0–13]	5.7 [4.1; 0–13]	0.84	0.40

Abbreviations: SD, standard deviation; p, significance; BDI, Beck Depression Inventory;³⁷ range 0–63, AUDIT, Alcohol Use Disorders Identification Test;³⁸ range 0–30.

Table 3. The blood lead level and the number of shot bullets and their correlations.

	Mean	SD	r	p
Blood lead ($\mu\text{mol/l}$)	0.52	0.40		
Bullets fired during the last month	326.00	509.28	0.705	0.001
Bullets fired during the last year	4645.00	4908.37	0.551	0.012

Abbreviations: SD, standard deviation; r, Pearson correlation coefficient; p, significance.

The results of the neuropsychological tests are shown in Table 4. The difference between the two groups on the memory variables was significant [MANCOVA $F(9,29) = 5.021$, $p \leq 0.001$]. The sport shooters' performance was markedly poorer than that of the controls on the incidental memory test and on the immediate and delayed verbal logical memory. The groups did not differ significantly on the intelligence tests [MANCOVA $F(4,34) = 1.577$, $p = 0.203$]. The correlations between the psychological test scores and the blood lead level or the shooting habits did not reach statistical significance.

Discussion

The aim of this study was to find whether sport shooters' lead exposure has an adverse effect on

memory functions. The sport shooters performed significantly worse than the controls on two verbal memory tests, namely on the immediate and delayed Logical Memory test and the incidental memory test. No other significant differences were found in the verbal or visual memory tests. The groups did not differ on the intelligence tests, on mood or alcohol consumption. To the best of our knowledge there is very little evidence so far about neuropsychological effects caused by lead exposure on firing ranges.

In our study the blood lead levels among sport shooters were slightly higher than the reference limit for non-exposed people. In Finland the reference level of non-exposed people is $0.3 \mu\text{mol/L}$ (responding $6.21 \mu\text{g/dl}$) determined by Finnish Institute of Occupational Health. The level is in

Table 4. Comparison of the neuropsychological test scores between shooters and controls tested with MANCOVA.

Neuropsychological test	Shooters		Controls		F-value	p
	Mean	SD	Mean	SD		
A. Memory tests (WMS-R)						
Incidental memory ¹	13.85	1.93	15.60	1.39	13.71	≤0.001
Logical Memory, immediate recall	10.40	3.25	13.50	3.33	4.32	0.021
Logical Memory, delayed recall ²	73.61	11.05	84.80	11.69	5.06	0.011
Verbal Learning, immediate recall	14.60	4.54	15.35	4.34	0.16	0.689
Verbal Learning, delayed recall ²	101.52	27.69	105.08	23.58	0.14	0.713
Digit Span, forward	7.20	0.37	8.00	0.38	2.84	0.071
Digit Span, backward	6.95	0.33	6.55	0.37	0.84	0.620
Visual Retention, immediate recall	34.95	3.49	33.20	4.98	2.82	0.072
Visual Retention, delayed recall ²	89.26	14.59	90.36	21.24	1.61	0.215
B. Intelligence tests (WAIS-R)						
Similarities	28.25	2.92	28.65	3.27	0.16	0.691
Picture Completion	18.10	1.83	18.45	2.01	0.25	0.624
Block Design	33.70	8.70	32.05	9.93	0.84	0.365
Digit Symbol	44.55	9.67	48.25	10.54	1.12	0.298

Abbreviations: SD, standard deviation; p, significance; WMS-R, Wechsler Memory Scale - Revised; see possible ranges in Table 1, ¹Vilkkilä et al. 1992, ²See Methods; WAIS-R, Wechsler Adult Intelligence Scale - Revised; see possible ranges in Table 1.

balance with other papers.^{40,41} The shooters' mean blood lead level was lower than that found in earlier studies concerning shooting with lead bullets^{2,6,7} but still higher than that found among Swedish police officers.³ In spite of the low average blood lead level in this study, a significant correlation was found between the numbers of bullets fired during the last month or year and the blood lead levels of the sport shooters. This may indicate that the amount lead fumes inhaled in the proximity of the gun in firing ranges mainly increases the PbB level. This observation is supported by Demmeler et al,⁸ where rounds shot each month highly correlated with PbB concentrations of 129 recreational indoor-shooters. In our study it was assumed that the PbB levels of the controls did not exceed the reference limit for non-exposed people.

In studies reporting the detrimental effects of occupational lead exposure on verbal learning,^{16,42} the PbB levels were considerably higher than those of the present study. Peak tibial lead has been associated with poor performance in verbal memory.¹⁹ On the other hand, chronic occupational lead exposure, irrespective of the current PbB level, has been demonstrated to impair encoding and retrieval of verbal information in the Rey Auditory Verbal Learning Test^{19,20,23} and in the Logical Memory subtest of the WMS-R.¹⁴ Similar verbal memory deficits have been reported with cumulative environmental lead exposure irrespective of recent PbB levels.²⁴ We did not examine the cumulative lead exposure, which would be an interesting topic for further studies. In the present study we focused on the current effects of continuous lead exposure on contemporary memory functions. According to our results, low recreational lead exposure is associated with impaired retrieval of verbal material in sport shooters. The groups did not differ in the Verbal Learning subtest of the WMS-R, which may indicate that encoding is not impaired in sport shooters and that the cueing in this test created an advantageous context to them for the retrieval of verbal associations.

There are contradictory results concerning the effect of low lead exposure on cognitive functions. On the one hand, relatively low blood lead levels (mean = 30.8 µg/100 ml; responding 1.49 µmol/l) due to occupational exposure have been associated with neuropsychological deficits, for example with executive and visuospatial dysfunctions.²¹ On the other hand, it has been documented that very low blood lead levels (mean = 5.4 µg/100 ml;

responding 0.26 µmol/l) do not result in cognitive deficits in individuals who have formerly been occupationally exposed to lead.⁴³ Interestingly, our study indicated that even a low, slightly elevated blood lead level is associated with impaired verbal memory after long-term exposure to low lead levels. The memory dysfunction was not caused by deficient verbal reasoning, because the performance of the sport shooters and the controls on the reasoning tests was almost equal. In our study the lead levels of the sport shooters were considerably lower than in other studies, which have documented neuropsychological defects caused by occupational lead exposure.⁴²

One limitation of this study is that it was not possible to conduct the blood lead level analysis for the control group, although it is highly likely that the lead exposure of the controls did not exceed the normal level. It is worth noting that all sport shooters had practised shooting for many years. Although the blood lead level does not indicate the effect of long-term cumulative lead exposure, our results suggest that long-term recreational lead exposure results in memory impairment. To summarize, recreational lead exposure was associated with verbal memory deficits examined in clinical neuropsychological tests. In future studies it would be interesting to examine whether recreational lead exposure has equally progressive effects on verbal or visual memory as occupational lead exposure after that exposure ceases.

Disclosure

The authors report no conflicts of interest.

References

1. Lösel H. Does lead pose a special danger to sport shooters. *UIT Journal*. 1992a;3:8–11.
2. Abudhaise BA, Alzoubi MA, Rabi AZ, et al. Lead exposure in indoor firing ranges: environmental impact and health risk to range users. *Int J Occup Med Environ Health*. 1996;9:323–39.
3. Löfstedt H, Seldén A, Storéus L, et al. Blood lead in Swedish police officers. *Am J Ind Med*. 1999;35:519–22.
4. Fischbein A, Nicholson WS, Weisman I. Comparative lead emissions from conventional and jacketed ammunition. *Am Ind Hyg Assoc J*. 1980;41:525–7.
5. Lösel H. Does lead pose a special danger to sport shooters. Part 2. *UIT Journal*. 1992b;4:8–13.
6. Valway SE, Martyny JW, Miller JR, et al. Lead absorption in indoor firing range users. *AJPH*. 1989;79:1029–32.
7. Gulson BL, Palmer JM, Bryce A. Changes in blood lead of recreational shooter. *Sci Total Environ*. 2002;293:143–150.
8. Demmeler M, Nowak D, Schierl R. High blood lead levels in recreational indoor-shooters. *Int Arch Occup Environ Health*. 2009;82:539–49.

9. Hänninen H, Hernberg S, Mantere P, et al. Psychological performance of participants with low exposure to lead. *J Occup Med*. 1978;20:683–9.
10. Stollery B, Banks H, Broadbent D, et al. Cognitive functioning in lead workers. *Br J Ind Med*. 1989;46:698–707.
11. Stollery B, Broadbent D, Banks H, et al. Short term prospective study of cognitive functioning in lead workers. *Br J Ind Med*. 1991;48:739–49.
12. Schwartz BS, Bolla KI, Stewart W, et al. Decrements in neurobehavioral performance associated with mixed exposure to organic and inorganic lead. *Am J Epidemiol*. 1993;137:1006–21.
13. White RF, Diamond R, Proctor S, et al. Residual cognitive deficits 50 years after lead poisoning during childhood. *Br J Ind Med*. 1993;50:613–22.
14. Lindgren KN, Masten VL, Ford DP, et al. Relation of cumulative exposure to inorganic lead and neuropsychological test performance. *Occup Environ Med*. 1996;53:472–7.
15. Muldoon SB, Cauley JA, Kuller LH, et al. Effects of blood lead levels on cognitive function of older women. *Neuroepidemiology*. 1996;15:62–72.
16. Bleecker ML, Lindgren KN, Ford DP. Differential contribution of current and cumulative indices of lead dose to neuropsychological performance by age. *Neurology*. 1997;48:639–45.
17. Hänninen H, Aitio A, Kovala, et al. Occupational exposure to lead and neuropsychological dysfunction. *Occup Environ Med*. 1998;55:202–9.
18. Daniell WE, Claypoole KH, Checkoway H, et al. Neuropsychological function in retired workers with previous long term occupational exposure to solvents. *Occup Environ Med*. 1999;56:93–105.
19. Stewart WF, Schwartz BS, Simon D, et al. Neurobehavioral function and tibial and chelatable lead levels in 543 former organolead workers. *Neurology*. 1999;52:1610–7.
20. Schwartz BS, Stewart WF, Bolla KI, et al. Past adult lead exposure is associated with longitudinal decline in cognitive function. *Neurology*. 2000;55:1144–50.
21. Schwartz BS, Lee BK, Lee GS, et al. Associations of blood lead, dimercaptosuccinic acid-chelatable lead, and tibia lead with neurobehavioral test scores in South Korean lead workers. *Am J Epidemiol*. 2001;153:453–64.
22. Barth A, Schaffner AW, Osterode W, et al. Reduced cognitive abilities in lead-exposed men. *Int Arch Occup Environ Health*. 2002;75:394–8.
23. Bleecker ML, Ford DP, Lindgren KN, et al. Differential effects of lead exposure on components of verbal memory. *Occup Environ Med*. 2005;62:181–7.
24. Shih RA, Glass TA, Bandeen-Roche K, et al. Environmental lead exposure and cognitive function in community-dwelling older adults. *Neurology*. 2006;67:1556–62.
25. Sillbergeld E. Neurological perspective on lead toxicity. In Needleman H, ed. *Human Lead Exposure*. Boca Raton, FL: CRC Press. 1992; p. 89–103.
26. Stewart WF, Schwartz BS, Davatzikos C, et al. Past adult lead exposure is linked to neurodegeneration measured by brain MRI. *Neurology*. 2006;66:1476–84.
27. Stewart W, Schwartz B. Effects on lead on the adult brain: A 15-year exploration. *Am J Ind Med*. 2007;50:729–39.
28. Schwartz BS, Chen S, Caffo B, et al. Relations of brain volumes with cognitive function in males 45 years and older with past lead exposure. *Neuroimage*. 2007;37:633–41.
29. Goodman M, LaVerda N, Clarke C, et al. Neurobehavioral testing in workers occupational exposed to lead: systematic review and meta-analysis of publications. *Occup Environ Med*. 2002;59:217–23.
30. Canfield RL, Henderson CR, Cory-Slechta DA, et al. Intellectual impairment in children with blood lead concentrations below 10 µg per deciliter. *N Engl J Med*. 2003;16:1517–26.
31. Jusko TA, Henderson Jr CR, Lanphear BP, et al. Blood lead concentrations <10 µg/dL and child intelligence at 6 years of age. *Environ Health Perspect*. 2008;2:243–8.
32. Surkan PJ, Zhang A, Trachtenberg F, et al. Neuropsychological function in children with blood lead levels <10 µg/dL. *Neurotoxicology*. 2007;28:1170–7.
33. Lanphear BP, Hornung R, Khoury K, et al. Low-level environmental lead exposure and children's intellectual function: an international pooled analysis. *Environ Health Perspect*. 2005;7:894–9.
34. Wechsler D. Wechsler Adult Intelligence Scale - Revised. New York: The psychological Corporation 1981.
35. Wechsler D. Wechsler Memory Scale - Revised. New York: The Psychological Corporation 1987.
36. Vilkkii J, Holst P, Öhman J, et al. Cognitive test performances related to early and late computed tomography findings after closed-head injury. *J Clin Exp Neuropsychol*. 1992;14:518–32.
37. Beck AT, Ward CH, Mendelson M, et al. Inventory for measuring depression. *Arch Gen Psychiatry*. 1961;4:561–71.
38. Saunders JB, Aasland OG, Babor TF, et al. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption-II. *Addiction*. 1993;88:791–804.
39. Subramanian KS, Meranger A. A rapid electrothermal atomic absorption spectrophotometric method for cadmium and lead in human whole blood. *Clin Chem*. 1981;27:1866–71.
40. Grandjean P, Nielsen GD, Jørgensen PJ, Horder M. Reference intervals for trace elements in blood: significance of risk factors. *Scand J Clin Lab Invest*. 1992;52:321–37.
41. Hense HW, Filipiak B, Novak L, Stoeppler M. Nonoccupational determinants of blood lead concentrations in a general population. *Int J Epidemiol*. 1992;21:753–62.
42. Meyer-Baron M, Seeber A. A meta-analysis for neurobehavioural results due to occupational lead exposure with blood lead concentrations <70 µg/100 ml. *Arch Toxicol*. 2000;73:510–8.
43. Winker R, Barth A, Ponocny-Seliger E, et al. No cognitive deficits in men formerly exposed to lead. *Wien Klin Wochenschr*. 2005;117:755–60.