

Erratum

Source: The Condor, 110(3): 597

Published By: American Ornithological Society

URL: https://doi.org/10.1525/cond.2008.110.3.erratum


The BioOne Digital Library (https://bioone.org/) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (https://bioone.org/subscribe), the BioOne Complete Archive (https://bioone.org/archive), and the BioOne eBooks program offerings ESA eBook Collection (https://bioone.org/esa-ebooks) and CSIRO Publishing BioSelect Collection (https://bioone.org/esa-ebooks) and CSIRO Publishing BioSelect Collection (https://bioone.org/csiro-ebooks).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.


Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commmercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ERRATUM



In Condor 110/2 (May 2008), Figure 2 on p. 290 of the paper "Reevaluating marine diets of Surf and White-winged Scoters: interspecific differences and the importance of soft-bodied prey," by Eric M. Anderson et al. did not print correctly due to a combination of insufficient figure resolution and composition of the original figure. The correct figure appears below. The authors and editors regret the error.






FIGURE 2. (A) Lengths of whole bivalves (n=104) consumed by scoters collected in Puget Sound, Washington, in 2005–2006. (B) Averages (\pm SE) across this and past studies (number of studies within each bar) for the mean and maximum lengths of bivalves in esophagus + gizzard contents for Surf and White-winged Scoters. (C) Average percent of diet across this and past studies comprised of hard-bodied prey (mollusks) for Surf and White-winged Scoters. In (C), medians and means are represented by solid and dashed lines, respectively; boxes indicate the 25th and 75th percentiles, and whiskers the 10th and 90th percentiles (the circle is an outlier). *P<0.05, *P<0.01.

The Condor, Vol. 110, Number 3, page 597. ISSN 0010-5422, electronic ISSN 1938-5129. © 2008 by The Cooper Ornithological Society. All rights reserved. Please direct all requests for permission to photocopy or reproduce article content through the University of California Press's Rights and Permissions website, http://www.ucpressjournals.com/reprintInfo.asp. DOI: 10.1525/cond.2008.110.3.erratum