

On the Status of Some Species of *Cheirostylis* Blume (Orchidaceae) from India

Author: Bhattacharjee, Avishek

Source: Candollea, 67(1) : 31-36

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2012v671a3>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

On the status of some species of *Cheirostylis* Blume (Orchidaceae) from India

Avishek Bhattacharjee

Abstract

BHATTACHARJEE, A. (2012). On the status of some species of *Cheirostylis* Blume (Orchidaceae) from India. *Candollea* 67: 31-35. In English, English and French abstracts.

Cheirostylis chinensis var. *glabra* Bhaumik & M. K. Pathak and *Cheirostylis seidenfadeniana* C. S. Kumar & F. N. Rasm. (Orchidaceae) are treated as synonyms of *Cheirostylis moniliformis* (Griff.) Seidenf. and *Cheirostylis parvifolia* Lindl. respectively. The taxonomic status of *Cheirostylis munnacampensis* A. N. Rao as a synonym of *Cheirostylis yunnanensis* Rolfe is confirmed and the status of *Cheirostylis mohanramii* Chaturv., Moakum & C. S. Kumar is also discussed.

Key-words

ORCHIDACEAE – *Cheirostylis* – India – Taxonomy

Résumé

BHATTACHARJEE, A. (2012). Sur le statut de quelques espèces de *Cheirostylis* Blume (Orchidaceae) originaires d'Inde. *Candollea* 67: 31-35. En anglais, résumés anglais et français.

Cheirostylis chinensis var. *glabra* Bhaumik et M. K. Pathak et *Cheirostylis seidenfadeniana* C. S. Kumar & F. N. Rasm. (Orchidaceae) sont traités comme synonymes de *Cheirostylis moniliformis* (Griff.) Seidenf. et *Cheirostylis parvifolia* Lindl. respectivement. Le statut taxonomique de *Cheirostylis munnacampensis* A. N. Rao comme synonyme de *Cheirostylis yunnanensis* Rolfe est confirmé et celui de *Cheirostylis mohanramii* Chaturv., Moakum & C. S. Kumar est discuté.

Address of the author: Central National Herbarium, Botanical Survey of India, A.J.C.B. Indian Botanic Garden, Howrah- 711 103, India. E-mail: avibsi@rediffmail.com

Submitted on May 11, 2010. Accepted on July 21, 2011.

Edited by P. Perret & P. Bungener

Cheirostylis Blume (*Orchidaceae*) is a genus of terrestrial, sometimes lithophytic, rarely epiphytic orchids characterized by fleshy, usually moniliform rhizome with internodal rhizoids, connate sepals forming a sepaline tube, hypochile of lip with 1 to several parallel seriate appendages and column with 2 stelids and 2 well-separated stigma lobes. PRIDGEE & al. (2003) and CHEN & al. (2009) reported the occurrence of about 50 species of *Cheirostylis* in the world distributed in tropical Africa, through South-east Asia, Japan, Indonesia and Pacific Islands to Australia and placed the genus under the subtribe *Goodyerinae* Ridl. of the tribe *Cranichideae* Endl. belonging to the subfamily *Orchidoideae* Lindl. Based on a study of type materials along with fresh, pickled and herbarium specimens of relevant taxa of *Cheirostylis* revealed the conspecificity of recently described *C. chinensis* var. *glabra* Bhaumik & M. K. Pathak and *C. seidenfadeniana* C. S. Kumar & F. N. Rasm. with earlier existing *C. moniliformis* (Griff.) Seidenf. and *C. parvifolia* Lindl. respectively. CHEN & al. (2009) recently treated *C. munnacampensis* A. N. Rao as a synonym of *C. yunnanensis* Rolfe without providing any clarification. This status is confirmed and the one of *C. mohanramii* Chaturv., Moakum & C. S. Kumar is also discussed (under *C. griffithii* Lindl.).

Cheirostylis moniliformis (Griff.) Seidenf. in Dansk Bot. Ark. 32: 69. 1978.

= *Goodyera moniliformis* Griff., Itin. Pl. Khasiyah Mts.: 143, no. 679. 1848.

Typus: BHUTAN “Bootan”: Griffith A 679 (holo-: K [K000 387624], photo!; iso-: OXF).

= *Cheirostylis chinensis* var. *glabra* Bhaumik & M. K. Pathak in Bull. Bot. Surv. India 47: 183. 2006. **Typus:** INDIA. **Arunachal Pradesh:** Dibang Valley district, Bejari, 150 m, 5.III.2004, Bhaumik & Tham 104752[A] (holo-: CAL!; iso-: ASSAM!, 2 sheets [B] [C]), syn. nov.

Specimens examined. – INDIA. **Arunachal Pradesh:** Lohit District, Minzong-yasang, 3.X.1985 [flowered on 11.III.1986 at Sessa], A. N. Rao 21463, 21489 (Orchid Herbarium Tipi); West Kameng District, 2 km from Munna Camp (towards Dirang), 1491 m, 16.IV.2006, A. Bhattacharjee 34819 A, 34819 B (CAL).

Note. – BHAUMIK & PATHAK (2006) described *C. chinensis* var. *glabra* based on collections from Arunachal Pradesh and distinguished it from the type variety in having glabrous bracts and ovary, acuminate petals and epichile with 2 broad green patches at the base. But these authors overlooked the earlier described *C. moniliformis* which is identical (Fig. 1) with their variety. Thus, *C. chinensis* var. *glabra* is treated here as a heterotypic synonym of *C. moniliformis*.

Cheirostylis parvifolia Lindl. in Edwards’s Bot. Reg. 25: 19. 1839.

Typus: SRI LANKA “Ceylon”: Loddiges s.n. (holo-: K [K000718267], photo!).

= *Cheirostylis seidenfadeniana* C. S. Kumar & F. N. Rasm. in Nordic J. Bot. 7: 409. 1987. **Typus:** INDIA. **Kerala:** Ponmudi, Trivandrum Dist., 950 m, 25.X. 1983, C. Sathish Kumar CU 36960 (holo-: TBGT; iso-: C, CALI), syn. nov.

Distribution. – India: Kerala, Maharashtra (fide JAYAWEEA, 1981; PUNEKAR, 2002); Sri Lanka.

Note. – SATHISH KUMAR & RASMUSSEN (1987) distinguished *C. seidenfadeniana* from *C. parvifolia* on the basis of epichile with entire lobules and 2 conspicuous tufts of hairs at its base. The authors mentioned that the material of *C. parvifolia* was very scarce from India and they could consult only 3 old herbarium specimens including the type (*Loddiges s.n.*, K) which was in very bad condition. However, it has been found that the holotype including its associated illustrations of flower and dissected floral parts (on the type sheet) is sufficient for its purpose and there is no need of epitypification at present. During the present work, critical study of the live specimens of *C. seidenfadeniana* provided by Dr. C. Sathish Kumar (first author of *C. seidenfadeniana*) revealed that the minute tufts of hairs at the base of epichile can be seen only in live, pickled or very well preserved specimens but not in the old herbarium specimens. The present study also reveals that the margin of the epichile lobules of *C. seidenfadeniana* varies (Fig. 1) from nearly entire to 2-4 lacerate which clearly indicates that *C. seidenfadeniana* and *C. parvifolia* are the same species. Thus, *C. seidenfadeniana* is treated here as synonym of *C. parvifolia*. In spite of author’s personal visit to CALI and TBGT, the holotype and isotype(s) of *C. seidenfadeniana* could not be located as claimed in the protologue. Dr. Olof Ryding, Curator of Vascular Plants of C also confirmed the unavailability of any type specimen of *C. seidenfadeniana* in C. Surprisingly, a specimen bearing the same collection number and field data (excepting the date which is prior to publication of *C. seidenfadeniana*) as that of the holotype and isotypes of *C. seidenfadeniana* has been located at K. In absence of all the designated types (holotype and isotypes), the K-specimen may be selected as lectotype (if there is no doubt that it was actually used while describing *C. seidenfadeniana*) or as neotype of *C. seidenfadeniana*. For the time being the lecto- or neotypification has not been done in the present paper with a hope that the types may be found in near future. This rare species is presently grown and conserved in the Orchidarium of Tropical Botanic Garden and Research Institute, Thiruvananthapuram, Kerala, India by Dr. Sathish Kumar and his co-workers.




Fig. 1. – Comparison plate: '*Cheirostylis chinensis* var. *glabra* Bhaumik & M. K. Pathak'. **A.** Labellum; **B.** Petal; **C.** Column with anther-cap – *Cheirostylis moniliformis* (Griff.) Seidenf.; **D-E.** Labellum; **F.** Petal; **G.** Column – '*Cheirostylis seidenfadeniana* C. S. Kumar & F. N. Rasm.'; **H.** Labellum – *Cheirostylis parvifolia* Lindl.; **I-J.** Labellum. [A-C: after BHAUMIK & PATHAK, 2006; D, F-G: A. Bhattacharjee 34819 A, CAL; E: A. Bhattacharjee 34819 B, CAL; H: after SATHISH KUMAR & RASMUSSEN, 1987; I: Sathish Kumar s.n., TBGT, spirit; J: A. Bhattacharjee 38142 A, CAL]

Cheirostylis yunnanensis Rolfe in Bull. Misc. Inform. Kew 1896: 201. 1896.

Typus: CHINA. Yunnan: Mengtse, IV.1893, *Hancock* 25 (holo-: K, photo!).

= *Cheirostylis munnacampensis* A. N. Rao in Nordic J. Bot. 8: 340. 1988 (synonymised by CHEN & al., 2009). **Typus:** INDIA. Arunachal Pradesh: Munna Camp (W. Kameng Dist.), 2000 m, 16.V.1984, A. N. Rao 14567 (holo-: Orchid Herbarium Tipi!; iso-: Orchid Herbarium Tipi!).

Distribution. – India: Arunachal Pradesh, Sikkim (fide PEARCE & CRIBB, 2002), West Bengal; China; Myanmar; Thailand; Vietnam.

Other specimens examined. – INDIA. Arunachal Pradesh: Tipi, 28.III.2002, A. N. Rao 30862 (Orchid Herbarium Tipi). **West Bengal:** Darjeeling District, Mahananda Wildlife Sanctuary, 1 km away from Kalijhora towards Setikhola, 12.III.2008, T. K. Paul 43698 (CAL).

Note. – While describing *C. munnacampensis*, RAO (1988) discriminated it from *C. yunnanensis* in having oblong petals, hypochile with 7 pairs of calli and stelids almost two times longer than the rostellar arms instead of spatulate petals, hypochile with 3 or 4 pairs of calli and stelids almost as long as the rostellar arms found in *C. yunnanensis*. But *C. yunnanensis* is a highly variable species and shows variation in the shape of petals, number of calli inside the hypochile, relative length of stelids and rostellar arms. The shape of petals varies from obliquely oblong to spatulate to obliquely elliptic-lanceolate. SEIDENFADEN (1978) observed obtuse apices of the petals, whereas in the present study both obtuse and acute apices of petals have been found. During the present study 3-9 pairs of calli observed inside the hypochile of *C. yunnanensis*. The relative length of stelids and rostellar arms also varies in different specimens of *C. yunnanensis*, but the stelids are always longer than the rostellar arms. Considering these ranges of variation (Fig. 2), *C. munnacampensis* is confirmed as a heterotypic synonym of *C. yunnanensis*.

Cheirostylis griffithii Lindl. in J. Proc. Linn. Soc., Bot. 1: 188. 1857.

Lectotypus (designated by SEIDENFADEN, 1978): INDIA. Meghalaya: Mamloo, *Griffith s.n.* (K-LINDL, photo!).

= *Cheirostylis mohanramii* Chaturv., Moaakum & C. S. Kumar in Nagaland Univ. SAP-Seminar Leaflet: [4]. 2009, nom. nud.

Distribution. – India: Arunachal Pradesh (fide CHOWDHURY, 1998), Meghalaya, Nagaland, Sikkim (fide LUCKSOM, 2007), Uttaranchal (fide DEVA & NAITHANI, 1986), West Bengal; Bangladesh; China; Myanmar; Nepal; Pakistan; Thailand.

Specimens examined. – INDIA. Meghalaya: East Khasi Hills District, Mawphlong forest, 7.XI.1966, S. K. Katak 37155 (ASSAM); Mawsmi, Khasi Hills, 19.XII.1972, P. K. Hajra 51876 (ASSAM). **Nagaland:** Naga Hills, 1935, N. L. Bor s.n. (DD). **West Bengal:** Darjeeling District, Kumai, near Jaldhaka river, 1219 m, XI.1894, *Pantling* 353 (CAL).

Note. – The name *C. mohanramii* Chaturv., Moaakum & C. S. Kumar was assigned to a collection from Aotsakilimi, Zunheboto district of Nagaland. It was claimed as a new species with a published photograph of an inflorescence (with three flowers) and brief information on its habit, habitat and phenology in a leaflet during a seminar organized by Department of Botany, Nagaland University on 20th April, 2009. The leaflet is also available online at «http://www.nagauniv.org.in/menu/events/SAP_SEMINAR_20April2009.pdf» and a photo-plate of *C. mohanramii* has also been deposited in CAL. However, the name *C. mohanramii* does not fulfill the conditions of an effective and valid publication as it lacks Latin diagnosis or description and type(s). After the study of dissected floral parts (photographs) of '*C. mohanramii*' sent by Dr. C. Sathish Kumar, it is found identical to *C. griffithii*. In this circumstances *C. mohanramii* does not require to be validly published.

Acknowledgements

I express my sincere thanks to the Director, Botanical Survey of India for providing facilities and encouragement. I gratefully acknowledge my supervisors Dr. H. J. Chowdhery, Botanical Survey of India (BSI), NRC, Dehradun and Mr. R. K. Bhakat, Department of Botany and Forestry, Vidyasagar University, Midnapore for their guidance. I am also thankful to Mr. Paul Ormerod, Queensland, Australia, Dr. K. N. Gandhi, Harvard University Herbarium, Cambridge, Dr. P. Venu, Central National Herbarium, BSI, Howrah for their valuable suggestions and Dr. C. Sathish Kumar, Tropical Botanic Garden and Research Institute, Thiruvananthapuram for providing pickled as well as some live specimens of *C. parvifolia*. Thanks are also due to Dr. P. Lakshminarasimhan, former Indian Botanical Liaison Officer (IBLO), Kew and Dr. V. Prasad, IBLO, Kew for sending images of the types consulted during the study and Dr. A. N. Rao, State Forest Research Institute, Itanagar for necessary permission to consult the types of *C. munnacampensis* at Orchid Herbarium Tipi.

References

- BHAUMIK, M. & M. K. PATHAK (2006). A new variety of *Cheirostylis chinensis* Rolfe var. *glabra* (Orchidaceae) from Dibang Valley, Arunachal Pradesh, India. *Bull. Bot. Surv. India* 47: 183-184.
- CHEN, S., S. W. GALE, P. J. CRIBB & P. ORMEROD (2009). *Cheirostylis* Blume. In: WU, Z. Y., P. H. RAVEN & D. Y. HONG (ed.), *Fl. China* 25: 57-63. Science Press and Missouri Botanical Garden Press.




Fig. 2. – '*Cheirostylis munnaecampensis* A. N. Rao'. **A.** Flower; **B.** Petal; **C.** Labellum; **D.** Column with anther-cap – *Cheirostylis yunnanensis* Rolfe; **E.** Flower; **F-G.** Petal; **H.** Labellum (partly fragmented); **I.** Portion of hypophylla with appendages (magnified); **J.** Column.

[**A-D:** after RAO, 1988; **E-J:** T. K. Paul 43698, CAL, dissected from dry and pressed specimen]

CHOWDHURY, H. J. (1998). *Orchid flora of Arunachal Pradesh*. Bishen Singh Mahendra Pal Singh.

DEVA, S. & H. B. NAITHANI (1986). *The Orchid Flora of North West Himalaya*. Print & Media Associates.

JAYAWEEERA, D. M. A. (1981). Orchidaceae. In: DASSANAYAKE, M. D. & F. R. FOSBERG (ed.), *A revised handbook to the Flora of Ceylon 2*: 5-368. Amerind Publishing Co. Pvt. Ltd.

LUCKSOM, S. Z. (2007). *The orchids of Sikkim and North East Himalaya*. Concept.

PEARCE, N. R. & P. J. CRIBB (2002). Orchids of Bhutan. *Fl. Bhutan* 3(3). Royal Botanic Garden, Edinburgh and Royal Government of Bhutan.

PRIDGEON, A. M., P. J. CRIBB, M. W. CHASE & F. N. RASMUSSEN (2003). *Genera Orchidacearum 3, Orchidoideae (Part 2), Vanilloideae*. Oxford University Press.

PUNEKAR, S. (2002). Report of rare terrestrial orchid species *Cheirostylis parvifolia* Lindl. from Dajipur Wildlife Sanctuary, Kolhapur District of Maharashtra State, India. *J. Econ. Taxon. Bot.* 26: 105-107.

RAO, A. N. (1988). Two new species of *Cheirostylis* (Orchidaceae) from Arunachal Pradesh, India. *Nordic J. Bot.* 8: 339-340.

SATHISH KUMAR, C. & F. N. RASMUSSEN (1987). *Cheirostylis seidenfadeniana* sp. nov. (Orchidaceae) from India. *Nordic J. Bot.* 7: 409-411.

SEIDENFADEN, G. (1978). Orchid genera in Thailand VI. Neottioideae Lindl. *Dansk Bot. Ark.* 32(2).

