

Psydrax merrillii (Rubiaceae), a new combination for a Polynesian tree species

Author: Mouly, Arnaud

Source: Candollea, 79(2) : 295-298

Published By: The Conservatory and Botanical Garden of the City of Geneva (CJBG)

URL: <https://doi.org/10.15553/c2024v792a6>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Psydrax merrillii (Rubiaceae), a new combination for a Polynesian tree species

Arnaud Mouly

Abstract

MOULY, A. (2024). *Psydrax merrillii* (Rubiaceae), a new combination for a Polynesian tree species. *Candollea* 79: 295–298. In English, English abstract. DOI: <http://dx.doi.org/10.15553/c2024v792a6>

Canthium merrillii (Setch.) Christoph. is a tree restricted to the Polynesian archipelagos of Samoa, Tonga, and Wallis and Futuna. The paleotropical genus *Psydrax* Gaertn., which belongs to the tribe *Vanguerieae* of the family *Rubiaceae*, comprises unarmed shrubs bearing stamens with anthers reflexed on the filaments, whereas the species of the genus *Canthium* Lam. are characterized by being spinescent shrubs. *Canthium merrillii* (Setch.) Christoph. is therefore transferred to *Psydrax* based on the absence of spines. A description and an illustration of the species are provided.

Keywords

RUBIACEAE – *Psydrax* – Polynesia – New combination – Taxonomy

Address of the author:

Université de Franche-Comté, UMR CNRS 6249 Chrono-Environnement, 16 route de Gray, 25030 Besançon cedex, France.
E-mail: arnaud.mouly@univ-fcomte.fr

Submitted on June 16, 2024 – Accepted on October 21, 2024 – First published online on November 20, 2024
ISSN: 0373-2967 (print), 2235-3658 (online) – Published by the Conservatoire et Jardin botaniques de Genève
Open access article under Creative Commons Attribution Licence (CC BY 4.0)

Introduction

Canthium merrillii (Setch.) Christoph. (Rubiaceae, *Vanguerieae*), originally described as *Plectronia merrillii* Setch. (SETCHELL, 1924), is a tree up to 15 m high restricted to the Polynesian archipelagos of Samoa, Tonga, and Wallis and Futuna, where it thrives in dry littoral forests to cloud forests from sea level to 700 m of elevation.

Canthium Lam. is a genus characterized by its spinescent habit (LANTZ & BREMER, 2004), mainly occurring in continental and insular tropical Asia and East Tropical Africa. In contrast, the species belonging to *Psydrax* Gaertn. are unarmed and have stamens with anthers reflexed on the filaments. *Canthium merrillii* is a completely unarmed tree displaying stamens as mentioned for *Psydrax*. Moreover, the species shows further typical characters of the Pacific *Psydrax*: corymbiform inflorescence, style ended in a stylar head complex, and fruits with verrucose pyrenes compressed at the insertion of the hilum (MEYER, 2017).

WHISTLER (2004) already made the combination in the genus *Psydrax*, cited the basionym and its author but without reference to its place of valid publication, which renders it not validly published under ICN Art. 41.5 (TURLAND et al., 2018). The valid combination is made below by giving a full and direct reference to the place of valid publication of the basionym.

Materials and methods

This contribution is mainly based on the revision of digital images of specimens kept at UC, especially those belonging to Setchell's herbarium. Collections from NOU and P were also studied, as well as digital images of specimens from AK, BISH, K, and US.

New combination

Psydrax merrillii (Setch.) Whistler ex Mouly, **comb. nov.** (Fig. 1).

- = *Plectronia merrillii* Setch. in Pap. Dept. Marine Biol. Carnegie Inst. Wash. 20: 49. 1924. = *Canthium merrillii* (Setch.) Christoph. in Bull. Bernice P. Bishop Mus. 128: 201. 1935. **Holotypus:** AMERICAN SAMOA. **Tutuila Island:** Tutuila, fl., III.1922, *Setchell [leg. Babcock & Sütupe]* 423 (UC [UC220154] image!; iso-: UC [UC220155, UC469852] image!).
- *Psydrax merrillii* (Setch.) Whistler, Rainforest Trees Samoa: 143. 2004 [comb. inval.].

Tree, up to 15 m high, branches cylindrical to laterally compressed, sometimes subquadrangular, unarmed, glabrous; bark brown-reddish then grey. *Stipules* 4.5–5.5 × 3.5–4 mm,

triangular, connate at base, flat or slightly to deeply awned, persistent on young shoots. *Leaves* opposite, slightly coriaceous; lamina 8–16(–20) × 5–11(–12) cm, elliptic-ovate, base rounded to cuneate, apex obtuse, minutely mucronate; secondary nerves 5–7, slightly prominent underneath, tertiary nerves visible; domatia common, pocket-like, at the axil of secondary or tertiary nerves; petiole 0.9–1.3 cm long. *Inflorescences* 3–4 × 3–4 cm, axillary, paired at nodes, corymbiform, 15–30 flowers, axes thin and straight, topped by short navicular bracts, solitary or in pairs. *Flowers* hermaphroditic, fragrant; pedicel 2–8 mm long; hypanthium c. 2 mm in diam., turbinate, glabrous; calyx 5-lobed, glabrous outside, tube 0.4–0.8 mm long, lobes 0.5 mm long, triangular with obtuse apex; corolla cylindrical to campanulate, 5-lobed, pure white, turning yellow as it ages, tube 4–4.5 × 1.8–2.5 mm, mouth with a ring of moniliform trichomes, lobes 3.5–4 mm long, triangular-lanceolate, reflexed, thick, margin slightly involute dorsally, papillose internally; stamens inserted at the corolla-lobes sinuses, exerted, filament 1.5–2 mm long, anthers 2.5–3 mm long, oblong, attached medio-basally and dorsally, reflexed; style elongated, 8–9 mm long, stigmatic head 1.5–2 mm long, cylindrical-ovoid, papillose; ovary bilocular, a single ovule per locule. *Fruits* 8–11 mm in diam., 3.5–4 mm thick, discoid, dark green to black when mature and fleshy in vivo, verrucose in sicco; pyrenes 2, c. 8–9.5 mm long, subcylindrical, thick, verrucose, crescent moon shaped. *Seeds* c. 7 mm long.

Distribution and ecology. – The plant is restricted to Upolu and Savai'i in Samoa, Tutuila in American Samoa, Tafahi in Tonga, and Futuna in Wallis and Futuna. It thrives from the dry littoral forests near the coast to the cloud forests at elevations of 5 to 700 m.

Additional specimens examined. – **AMERICAN SAMOA. Tutuila Island:** Pago Pago, 426 m, 18.III.1930, *Diefenderfer* 13 (BISH); s.l., VI–VII.1920, *Setchell [leg. Sütupe]* 518 (UC [UC215601]); West of Aoloaifou, 300 m, 1.X.1976, *Whistler* 3672 (US). **SAMOA. Savai'i Island:** Gataivai (inland 1 mile), 50 m, 20.VIII.1968, *Bristol* 2269 (BISH); Taga, 5 m, 6.X.1931, *Christophersen* 2825 (BISH, P, US), 2835 (BISH); Salailua-Lataitai, 10 m, 19.X.1931, *Christophersen* 2860 (BISH); *ibid.*, 17.IX.1931, *Christophersen & Hume* 2619 (BISH, P, US); Asau, just east of the Potlatch Lumber Mill, 20 m, 11.III.1974, *Whistler* 1787 (BISH). **Upolu Island:** Ridge above Malololelei, 700 m, 14.VIII.1929, *Christophersen* 242 (BISH); Vaea, 350 m, 28.VIII.1929, *Christophersen* 459 (BISH, P, US); near Afulilo waterfall in the Le Mafa area, 350 m, 11.II.1974, *Whistler* 1571 (US); Ridge South of Mount Fao, 400 m, 2.I.1976, *Whistler* 3241 (US). **TONGA. Tafahi Island:** s.l., 530–540 m, 25.VII.1978, *Buelow* 1176 (AK); summit of the island, 19.XI.1987, *Whistler* 6255 (US); between Vai Kona and Vai Langita, 26.XI.1987, *Whistler* 6366 (US). **WALLIS AND FUTUNA. Futuna Island:** Nuku, Singave, 35 m, 1.VII.1974, *Kirch s.n.* (BISH); Nuku, Singave, along ridge leading to Osiiti, 30–40 m, 19.VII.1974, *Kirch* 160 (BISH); Mont Puke, 14.XI.2008, *Munzinger et al.* 5302 (K, NOU, P).




Fig. 1. – *Psydrax merrillii* (Setch.) Whistler ex Mouly. **A.** Flowering branch; **B.** Stipules; **C.** Flower in longitudinal section; **D.** Detail of fruiting branch; **E.** Fruit in lateral view.
[Setchell 423, UC] [Drawing: A. Mouly]

Acknowledgements

The author thanks Ana Penny at the University and Jepson Herbaria, University of California, Berkeley for providing digitized specimens.

References

- LANTZ, H. & B. BREMER (2004). Phylogeny inferred from morphology and DNA data: characterizing well-supported groups in Vanguerieae (Rubiaceae). *Bot. J. Linn. Soc.* 146: 257–283.
- MEYER, J.-Y. (2017). *Guide des plantes de Wallis et Futuna*. Au Vent des Iles, Pirae, Tahiti.
- SETCHELL, W.A. (1924). American Samoa: Part I. Vegetation of Tutuila Island. *Pap. Dept. Marine Biol. Carnegie Inst. Wash.* 314: 1–175.
- TURLAND, N.J., J.H. WIERSEMA, F.R. BARRIE, W. GREUTER, D.L. HAWKSWORTH, P.S. HERENDEEN, S. KNAPP, W.-H. KUSBER, D.-Z. LI, K. MARHOLD, T.W. MAY, J. MCNEILL, A.M. MONRO, J. PRADO, M.J. PRICE & G.F. SMITH (2018). International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. *Regnum Veg.* 159.
- WHISTLER, W.A. (2004). *Rainforest trees of Samoa. A guide to the common native and naturalized lowland and foothill forest trees of the Samoan Archipelago*. Isle Botanica, Hawaii.