



Native Hosts and Parasitoids Associated with *Anastrepha fractura* and Other *Anastrepha* Species (Diptera: Tephritidae) in the Brazilian Amazon

Authors: Dutra, Vivian Siqueira, Ronchi-Teles, Beatriz, Garcia, Marcos Vinicius Bastos, Adaime, Ricardo, and Silva, Janisete Gomes

Source: Florida Entomologist, 96(1) : 270-273

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.096.0144>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

NATIVE HOSTS AND PARASITOIDS ASSOCIATED WITH *ANASTREPHA FRACTURA* AND OTHER *ANASTREPHA* SPECIES (DIPTERA: TEPHRITIDAE) IN THE BRAZILIAN AMAZON

VIVIAN SIQUEIRA DUTRA¹, BEATRIZ RONCHI-TELES², MARCOS VINICIUS BASTOS GARCIA³, RICARDO ADAIME⁴,
AND JANISETE GOMES SILVA^{5*}

¹Graduate Program in Entomology, Instituto Nacional de Pesquisas da Amazônia, Caixa Postal 2223, 69080-971
Manaus, Amazonas, Brazil

²Coordenação de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia, Caixa Postal 2223, 69080-971
Manaus, Amazonas, Brazil

³Embrapa Amazônia Ocidental, Rodovia AM 10 km 29, 69011-970 – Manaus, Amazonas, Brazil

⁴Embrapa Amapá, Rodovia Juscelino Kubitschek, km 5, 68903-419 – Macapá, Amapá, Brazil

⁵Departamento de Ciências Biológicas, Universidade Estadual de Santa Cruz, Rodovia Ilhéus-Itabuna km 16,
45650-000 - Ilhéus, Bahia, Brazil

*Corresponding author; E-mail: jgs10@uol.com.br

The Brazilian Amazon region comprises an exceptionally-high diversity of angiosperms, among which approximately 180 described species, both native and exotic that can be potential fruit fly hosts (Silva & Ronchi-Teles 2000). Currently, 60 *Anastrepha* species have been recorded in the Brazilian Amazon region, of which about 30 species are endemic (Trindade & Uchôa 2011; Zucchi et al. 2011).

Here we report on new host/fruit fly/parasitoid associations for some *Anastrepha* species for both Brazil and the Amazon region. A total of 4,137 fruit (73.9 kg) from 40 different native and introduced plant species in 19 families were collected from 2008 through 2011 in Manaus (S 03° 06' 07" W 60° 01' 30"), Maués (S 03° 23' 01" W 57° 43' 07"), Presidente Figueiredo (S 02° 02' 04" W 60° 01' 30"), and São Gabriel da Cachoeira (S 00° 07' 49" W 7° 05' 21") in the state of Amazonas; in Porto Velho (S 08° 45' 43" W 63° 54' 14") in the state of Rondônia, and in Boa Vista (S 02° 49' 11" W 60° 40' 24") in the state of Roraima. Fallen fruit, both ripe and ripening, were collected randomly from the ground under tree canopies within the forest. Cassava fruits were collected in an area adjacent to the forest in Manaus (Amazonas) and Porto Velho (Rondônia). Adult flies and parasitoids were reared from collected fruits following methods described in Ronchi-Teles et al. (2011). Voucher specimens were deposited at the Coleção de Invertebrados of the Instituto Nacional de Pesquisas da Amazônia.

A total of 3,470 fruit (63.3 kg) were infested yielding 7,662 puparia from which 3,073 *Anastrepha* adults (1,469 males and 1,604 females), 669 braconid parasitoids, and 17 figitid parasitoids emerged. We report *Anastrepha fractura* Stone infesting fruit of *Salacia* sp. (Celastrales: Celastraceae) in association with *Asobara anastrephae* (Muesebeck), *Doryctobracon brasiliensis* (Szépligeti), and *Opius*

bellus Gahan parasitoids for the first time (Table 1). Previously, *A. fractura* had been reported solely from Guyana (Stone 1942) and Amazonas, and the only known host and associated parasitoid were *Maquira sclerophylla* (Ducke) C.C. Berg (Rosales: Moraceae) and *Doryctobracon areolatus* (Szépligeti) (Costa et al. 2009), respectively. We also report *Anastrepha distincta* Greene infesting *Inga cinnamomea* Spruce ex Benth (Fabales: Fabaceae: Mimosoideae) in Brazil for the first time (Table 1), but which was not attacked by any of the parasitoid species herein reported. The remaining 13 *Anastrepha* species recovered during sampling (Table 1) had previously been reported infesting the hosts from which they were recovered here (Zucchi et al. 2011).

We also report *O. bellus* and *Utetes anastrephae* (Viereck) associated with *Anastrepha coronilli* Carrejo & González, and *Aganaspis pelleranoi* (Brèthes) associated with *A. coronilli* and *Anastrepha striata* Schiner for the first time in Brazil. Our results indicate that native host plants in the Amazon forest in Brazil play an important role as reservoirs of native parasitoids and corroborate previous studies in forested areas (López et al. 1999; Aluja et al. 2003; Costa et al. 2009; Ronchi-Teles et al. 2011).

We would like to thank Ana Cláudia Araújo, Felipe Gomes, Mirian Santos, Neusa Hamada, Tatiana Senra, and Weber Novaes for their help with the collections. Thanks are also due to Carter R. Miller, the anonymous reviewers and the editor for comments on an earlier version of the manuscript. This study was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq 575664/2008-8) and Rede Amazônica de Pesquisas sobre Moscas-das-Frutas (EMBRAPA Chamada 05/2006 Agrofuturo). V.S.D. was supported under a CNPq fellowship during her Ph.D.

TABLE 1. *ANASTREPHA* SPECIES, HOSTS, AND ASSOCIATED PARASITOIDS IN THE AMAZON REGION, BRAZIL.

Plant Family and Species	Hosts	Collection sites	Sample weight (kg)	Number of fruit	Number of pupae	<i>Anastrepha</i> species (n)	Parasitoid species (n)
ANARCARDIACEAE <i>Anacardium occidentale</i> L. <i>Mangifera indica</i> L. <i>Spondias mombin</i> L.	Native Introduced Native	Manaus	0.9	28	15	4 <i>A. obliqua</i>	2 <i>D. areolatus</i>
		Manaus	2.5	15	109	27 <i>A. obliqua</i>	5 <i>D. areolatus</i>
		Manaus	7.9	1,038	1,491	149 <i>A. obliqua</i>	88 <i>D. areolatus</i> 43 <i>O. bellus</i> 6 <i>U. anastrephae</i> 0
	Presidente Figueiredo		3.1	232	586	9 <i>A. antunesi</i> 104 <i>A. obliqua</i> 8 <i>A. antunesi</i> ■	92 <i>O. bellus</i> 72 <i>A. anastrephae</i> 9 <i>D. areolatus</i> 1 <i>U. anastrephae</i> 1 <i>A. pelleranoi</i> 1 <i>A. nordlanderi</i> 0 0
		São Gabriel da Cachoeira	0.3	27	227	122 <i>A. obliqua</i> 8 <i>A. antunesi</i>	
		Manaus	0.6	14	731	181 <i>A. fractura</i>	60 <i>D. areolatus</i> 3 <i>A. anastrephae</i> ■ 1 <i>D. brasiliensis</i> ■ 1 <i>O. bellus</i> ■
CELASTRACEAE <i>Salacia</i> sp. ◆	Native	Manaus	4.9	162	300	77 <i>A. turpiniae</i>	41 <i>D. areolatus</i> ■
		Manaus	2.2	213	393	109 <i>A. pickeli</i>	128 <i>D. areolatus</i> 1 <i>O. bellus</i> 2 <i>U. anastrephae</i> 3 <i>O. bellus</i> ■ 0
EUPHORBIACEAE <i>Manihot esculenta</i> Crantz	Native	Porto Velho	0.8	85	9	1 <i>A. manihoti</i> 3 <i>A. pickeli</i>	
		Manaus Maués Maués	2.1 4.0 0.8	8 11 9	17 306 139	3 <i>A. distincta</i> 108 <i>A. distincta</i> 27 <i>A. distincta</i>	0 0 0
FABACEAE (MIMOSOIDAE) <i>Inga edulis</i> Mart.	Native						
<i>Inga cinnamomea</i> Spruce ex Benth. ◆	Native						

◆ New host record
■ New *Anastrepha* parasitoid association
■ Not possible to associate parasitoids with *Anastrepha* species

TABLE 1. (CONTINUED) ANASTREPHA SPECIES, HOSTS, AND ASSOCIATED PARASITOIDS IN THE AMAZON REGION, BRAZIL.

Plant Family and Species	Hosts	Collection sites	Sample weight (kg)	Number of fruit	Number of pupae	Anastrepha species (n)	Parasitoid species (n)
MALPIGHIACEAE <i>Malpighia glabra</i> L.	Introduced	Manaus	1.3	43	79	17 <i>A. obliqua</i>	21 <i>D. areolatus</i> 12 <i>O. bellus</i>
MELASTOMATACEAE <i>Bellucia grossularioides</i> (L.) Triana	Native	Manaus Presidente Figueiredo	2.2 1.5	219 226	151 133	41 <i>A. coronilli</i> 55 <i>A. coronilli</i>	13 <i>D. areolatus</i> 11 <i>D. areolatus</i> 2 <i>O. bellus</i> ■ 1 <i>U. anastrephae</i> ■ 1 <i>A. pelleranoi</i> ■
MORACEAE <i>Pouroma cecropiaefolia</i> Mart.	Native	Manaus	4.6	304	113	35 <i>A. bahiensis</i>	9 <i>D. areolatus</i> 1 <i>A. anastrephae</i> 1 <i>O. bellus</i> ■
MYRTACEAE <i>Eugenia stipitata</i> McVaugh	Native	Manaus	3.3	31	244	52 <i>A. obliqua</i>	0
		Maués	0.3	8	52	4 <i>A. obliqua</i>	0
<i>Psidium acutangulum</i> D.C.	Native	Manaus	1.2	12	82	33 <i>A. striata</i>	0
<i>Psidium guajava</i> L.	Native	Manaus	0.7	47	254	51 <i>A. striata</i>	0
		Maués	0.5	15	16	2 <i>A. obliqua</i>	0
		Presidente Figueiredo	2.7	73	210	7 <i>A. striata</i>	0
						51 <i>A. striata</i>	10 <i>Doryctobracon</i> sp.2
						5 <i>A. turpiniae</i> ■	8 <i>A. pelleranoi</i>
						6 <i>A. striata</i>	6 <i>A. pelleranoi</i> ■
						3 <i>A. striata</i>	0
<i>Psidium guineense</i> Swart.	Native	Porto Velho	0.9	49	53	20 <i>A. striata</i>	0
		São Gabriel da Cachoeira	0.8	16	46	21 <i>A. striata</i>	0
		Manaus	0.4	15	47	1 <i>A. striata</i>	0
		Porto Velho	0.3	11	23	0	1 <i>D. areolatus</i>
<i>Syzygium jambolanum</i> (Lam.) DC.	Introduced	Manaus	0.1	50	2	0	2 <i>D. areolatus</i>
<i>Syzygium malaccense</i> L.	Introduced	Manaus	0.3	3	6	3 <i>A. obliqua</i>	0
OXALIDACEAE <i>Averrhoa carambola</i> L.	Introduced	Manaus	0.7	11	35	3 <i>A. obliqua</i>	3 <i>D. areolatus</i>
		São Gabriel da Cachoeira	0.5	65	37	19 <i>A. obliqua</i>	0

◆ New host record
■ New *Anastrepha* parasitoid association
■ Not possible to associate parasitoids with *Anastrepha* species

TABLE 1. (CONTINUED) ANASTREPHA SPECIES, HOSTS, AND ASSOCIATED PARASITOIDS IN THE AMAZON REGION, BRAZIL.

Plant Family and Species	Hosts	Collection sites	Sample weight (kg)	Number of fruit	Number of pupae	Anastrepha species (n)	Parasitoid species (n)
PASSIFLORACEAE <i>Passiflora nitida</i> Kunth	Native	Manaus	3.4	63	205	42 <i>A. curitis</i>	0
RHAMNACEAE <i>Ziziphus mauritiana</i> Lam	Introduced	Boa Vista	1.3	194	4	1 <i>A. zenildae</i>	0
SAPOTACEAE <i>Pouteria caimito</i> (Ruiz & Pav.) Radlk.	Native	Manaus	5.2	160	1,484	105 <i>A. serpentina</i> 60 <i>A. leptozona</i> 1 <i>A. leptozona</i>	23 <i>D. areolatus</i> 1 <i>D. areolatus</i> 0
		Porto Velho	0.2	1	6	13 <i>A. serpentina</i> 13 <i>A. leptozona</i>	0
		São Gabriel da Cachoeira	0.8	12	57		

◆ New host record
■ New *Anastrepha* parasitoid association
■ Not possible to associate parasitoids with *Anastrepha* species

SUMMARY

A new natural host for *Anastrepha fractura* Stone, *Salacia* sp., is reported for the first time in Brazil. Parasitoids attacking *A. fractura* are also reported. We also report *Anastrepha distincta* Greene infesting *Inga cinnamomea* in Brazil for the first time. New associations between *Anastrepha* species and parasitoids are also reported.

RESUMO

Um novo hospedeiro natural de *Anastrepha fractura* Stone, *Salacia* sp., é relatado pela primeira vez no Brasil. Parasitoides atacando *A. fractura* são também relatados. Nós também relatamos *Anastrepha distincta* Greene infestando *Inga cinnamomea* no Brasil pela primeira vez. Novas associações entre espécies de *Anastrepha* e parasitoides também são relatadas.

REFERENCES CITED

ALUJA, M., RULL, J., SIVINSKI, J., NORRBOM, A. L., WHARTON, R. A., MACÍAS-ORDOÑEZ, R., DIAZ-FLEISCHER, F., AND LÓPEZ, M. 2003. Fruit flies of the genus *Anastrepha* (Diptera: Tephritidae) and associated parasitoids (Hymenoptera) in the tropical rain forest biosphere reserve of Montes Azules, Chiapas, Mexico. *Environ. Entomol.* 32: 1377-1385.

COSTA, S. G. M., QUERINO, R. B., RONCHI-TELES, B., PENTEADO-DIAS, A. M. M., AND ZUCCHI, R. A. 2009. Parasitoid diversity (Hymenoptera: Braconidae and Figitidae) on frugivorous larvae (Diptera: Tephritidae and Lonchaeidae) at Adolpho Ducke Forest Reserve, Central Amazon Region, Manaus, Brazil. *Braz. J. Biol.* 69: 363-370.

LÓPEZ, M., ALUJA, M., AND SIVINSKI, J. 1999. Hymenopterous larval-pupal and pupal parasitoids of *Anastrepha* flies (Diptera: Tephritidae) in Mexico. *Biol. Control* 15: 119-129.

RONCHI-TELES, B., DUTRA, V. S., TREGUE-COSTA, A. P., AGUIAR-MENEZES, E. L., MESQUITA, A. C. A., AND SILVA, J. G. 2011. Natural host plants and native parasitoids associated with *Anastrepha pulchra* and other *Anastrepha* species (Diptera: Tephritidae) in Central Amazon, Brazil. *Florida Entomol.* 94(2): 347-349.

SILVA, N. M., AND RONCHI-TELES, B. 2000. Amapá, Amazonas, Pará, Rondônia e Roraima. In Malavasi, A.; Zucchi, R. A. (eds.), *Moscas-das-frutas de importância Econômica no Brasil. Conhecimento básico e aplicado*. Holos-Editora, Ribeirão Preto, São Paulo. pp. 203-209.

STONE, A. 1942. The fruit flies of the genus *Anastrepha*. *USDA Misc. Publ.* 439: 1-112.

TRINDADE, R. B. R., AND UCHÔA, M. A. 2011. Species of fruit flies (Diptera: Tephritidae) in a transect of the Amazonian Rainforest in Oiapoque, Amapá, Brasil. *Zoologia* 28: 653-657.

ZUCCHI, R. A., URAMOTO, K., AND SOUZA-FILHO, M. F. 2011. Chave ilustrada para as espécies de *Anastrepha* da região Amazônica. In R. A. Silva, W. P. Lemos and R. A. Zucchi (eds.), *Moscas-das-frutas na Amazônia Brasileira. Diversidade, Hospedeiros e Inimigos Naturais*. Embrapá Amapá, Macapá, Amapá. pp. 73-90.