



Rhodomirtus tomentosa (Myrtales: Myrtaceae): New Host Records for Metanastria gemella (Lepidoptera: Lasiocampidae) and Arna bipunctapex (Lepidoptera: Lymantriidae)

Authors: Pratt, Paul D., Makinson, Jeff, Purcell, Matthew, Pogue, Michael G., Rayamajhi, Min B., et al.

Source: Florida Entomologist, 96(2) : 641-642

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.096.0235>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***RHODOMYRTUS TOMENTOSA* (MYRTALES: MYRTACEAE): NEW HOST RECORDS FOR *METANASTRIA GEMELLA* (LEPIDOPTERA: LASIOCAMPIDAE) AND *ARNA BIPUNCTAPEX* (LEPIDOPTERA: LYMANTRIIDAE)**

PAUL D. PRATT^{1,*}, JEFF MAKINSON², MATTHEW PURCELL², MICHAEL G. POGUE³, MIN B. RAYAMAJHI¹ AND TED D. CENTER¹

¹USDA/ARS, Invasive Plant Research Laboratory, 3225 College Ave., Ft. Lauderdale, FL 33314, USA

²USDA/ARS, Australian Biological Control Laboratory, CSIRO Ecosystem Sciences, GPO Box 2583 Brisbane, Queensland, Australia, 4001

³USDA/ARS, Systematic Entomology Laboratory, Smithsonian Institution, PO Box 37012, MRC-168, Washington, DC, 20013-7012, USA

*Corresponding author; E-mail: Paul.Pratt@ars.usda.gov

Rhodomyrtus tomentosa (Aiton) Hassk. (Myrtales: Myrtaceae) is a long lived evergreen shrub with large pink flowers and edible fruit. The native distribution of *R. tomentosa* includes much of southern Asia and the Philippines (Langeland & Craddock Burks 1998). Beginning in the late 1800s *R. tomentosa* was exported to various locations worldwide, including the states of Hawaii and Florida in the USA. The exotic shrub has escaped cultivation in both locations and invaded natural areas where it displaces native vegetation (Langeland & Craddock Burks 1998). The development of a biological control program targeting *R. tomentosa* has resulted in several surveys for host specific natural enemies in the plant's native range (Krauss 1966; Winotai et al. 2005). Recently, 2 lepidopteran species were discovered feeding on *R. tomentosa* during surveys in China.

On 30 Mar 2012, 43 larvae of a lasiocampid species were observed resting as a communal group on *R. tomentosa* branches near Kau Tam Tso (N 22° 30.248' E 114° 15.045') in the Wu Kau Tang area of Hong Kong, China. Larvae regurgitated a droplet of partially digested fluid when collected or when disturbed. While branches near the resting group were defoliated, larvae were held in a plastic container (10 × 10 × 5 cm) with *R. tomentosa* leaves to confirm that they consumed the target plant. Following positive results, larvae were shipped on 2 Apr 2012 to the quarantine facility at the United States Department of Agriculture (USDA), Agriculture Research Service (ARS), Invasive Plant Research Lab (IPRL) in Fort Lauderdale, Florida USA under permit # P-09-02955. Upon arrival, larvae were placed within a ventilated clear plastic container (50 × 50 × 50 cm) provisioned exclusively with *R. tomentosa* foliage until pupation. General maintenance of the colony resulted in several instances of larval setae imbedding in human skin, although no prolonged irritation occurred. Larvae rested communally during the day and fed at night. The resulting adults mated within the cage and oviposition ensued 4 days later; dead adults were preserved for identi-

fication. Eggs were placed in several petri dishes and held at 25 °C until hatching. To confirm that *R. tomentosa* supports complete development of the species, neonates were transferred to 1 of 3 ventilated plastic containers and fed leaves of the target plant until pupation. Adults were identified as *Metanastria gemella* Lajonquiere (Lasiocampidae), a species whose geographic range includes Sundaland, Palawan, and southern China (Holloway 1987). The host range of *M. gemella* is poorly understood but Holloway (1987) postulates that this species has a similar diet breadth as its congener *Metanastria hyrtaca* Cramer, which is a broad generalist. To our knowledge, the only host of *M. gemella* reported in the literature is the plantation tree *Neolamarckia cadamba* (Roxb.) Bosser (Gentianales: Rubiaceae) in Sandakan, Malaysia (Chung et al. 2009). These data clearly demonstrate that *R. tomentosa* is a developmental host of *M. gemella* and lends evidence that the species is a generalist, feeding on species across at least 2 plant families. It should also be noted that 4 tachinid flies emerged from *M. gemella* pupae and were identified as *Blepharipa* sp. by Norman E. Woodley of the Systematic Entomology Laboratory (SEL), ARS, USDA, Smithsonian Institution, Washington, DC, USA. Voucher specimens are lodged at the SEL.

On 4 Jun 2012 ca. 100 small orange lymantriid larvae were observed feeding on *R. tomentosa* near Shitan, Guangxi Province, China (N 21° 48.273' E 108° 58.106'). Larvae were shipped under the permit listed above to the IPRL quarantine facility on 18 Jun 2012. Larvae were fed *R. tomentosa* leaves in a cage similar to that described above and remained gregarious throughout their development. When disturbed, larvae wagged their heads and thoraxes from side to side in an apparently defensive behavior. Pupation occurred within light yellow silken cocoons and the resulting adults were mated; dead adults were preserved for identification. No parasitoids were recovered from the colony. Eggs arising from the adults were light yellow and covered in the

yellow scales of the female. Neonates were transferred to several plastic cages and fed exclusively *R. tomentosa* until pupation, demonstrating that the target weed is a developmental host. Adults of the lymantriid larvae were identified as *Arna bipunctapex* Hampson, a species that occurs across Southeast Asia, from India to Taiwan and Sundaland. Hosts of this species include representatives from 12 families, although *R. tomentosa* is not among the recorded food plants (Holloway 1987; Robinson et al. 2012). Voucher specimens are lodged at the SEL.

SUMMARY

Surveys for host specific natural enemies of *Rhodomyrtus tomentosa* (Myrtales: Myrtaceae), rose myrtle, in China (mainland and Hong Kong) resulted in the discovery of 2 lepidopteran species that have not been previously reported as herbivores of the plant: *Metanastria gemella* Lajonquiere (Lasiocampidae) and *Arna bipunctapex* Hampson (Lymantriidae). Multiple generations of both species were reared on *R. tomentosa* in a quarantine laboratory, demonstrating that the plant is a developmental host. However, neither species demonstrates sufficient host specificity to be considered for biological control of *R. tomentosa* in the USA.

Key Words: biological control, herbivores, host specificity, invasive species, rose myrtle

RESUMEN

Un sondeo de los enemigos naturales específicos de *Rhodomyrtus tomentosa* (Myrtales: Myrtaceae), rosa mirto, en China (continental y Hong Kong) resultó en el descubrimiento de dos

especies de lepidópteros que no han sido previamente reportados como herbívoros de la planta: *Metanastria gemella* Lajonquiere (Lasiocampidae) y *Arna bipunctapex* Hampson (Lymantriidae). Varias generaciones de ambas especies fueron criadas sobre *R. tomentosa* en un laboratorio de cuarentena, lo que demuestra que la planta es un hospedero en que estas especies pueden desarrollarse. Sin embargo, las dos especies no demostraron una suficiente especificidad sobre el hospedero para ser considerados en el control biológico de *R. tomentosa* en los EE.UU.

Palabras Clave: control biológico, herbívoros, especificidad del hospedador, especies invasoras, rosa mirto

REFERENCES CITED

- CHUNG, A. Y. C., AJIK, M., NILUS, R., HASTIE, A., ONG, R. C., AND CHEY, V. K. 2009. New records of insects associated with Laran (*Neolamarckia cadamba*) in Sabah. Sepilok Bull. 10: 45-63.
- HOLLOWAY, J. D. 1987. The Moths of Borneo. Southene Sdn Bhd. Kuala Lumpur Malaysia. <http://www.mothsofborneo.com/>. (Accessed: 28 Aug 2012).
- KRAUSS, N. L. H. 1966. Biological control investigations on some Hawaiian weeds. Proc. Hawaiian Entomol. Soc. 19: 223-231.
- LANGELAND, K. A., AND CRADDOCK BURKS, K. 1998. Identification and biology of non-native plants in Florida's natural areas. Univ. of Florida. 165 pp.
- ROBINSON, G. S., ACKERY, P. R., KITCHING, I. J., BECCALONI, G. W., AND HERNÁNDEZ, L. M. 2010. HOSTS - A Database of the world's lepidopteran hostplants. Natural History Museum, London. <http://www.nhm.ac.uk/hosts>. (Accessed: 28 Aug. 2012).
- WINOTAI, A., WRIGHT, T., AND GOOLSBY, J. A. 2005. Herbivores in Thailand on *Rhodomyrtus tomentosa* (Myrtaceae), an invasive weed in Florida. Florida Entomol. 88: 104-105.