



## **Ground Beetle (Coleoptera: Carabidae) Populations in Commercial Organic and Conventional Potato Production**

Authors: Rondon, Silvia I., Pantoja, Alberto, Hagerty, Aaron, and Horneck, Donald A.

Source: Florida Entomologist, 96(4) : 1492-1499

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.096.0430>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## GROUND BEETLE (COLEOPTERA: CARABIDAE) POPULATIONS IN COMMERCIAL ORGANIC AND CONVENTIONAL POTATO PRODUCTION

SILVIA I. RONDON<sup>1,\*</sup>, ALBERTO PANTOJA<sup>2,3</sup>, AARON HAGERTY<sup>4,5</sup> AND DONALD A. HORNECK<sup>1</sup>

<sup>1</sup>Oregon State University, Hermiston Agricultural Research and Extension Center, 2121 S. First Street, Hermiston, OR 97838, USA

<sup>2</sup>USDA, Agricultural Research Service, Subarctic Agricultural Research Unit, Fairbanks, AK, 99709, USA

<sup>3</sup>Current address: United Nations, Food and Agriculture Organization, Regional Office for Latin America and the Caribbean, Vitacura, Santiago, Chile. Avenida Dag Hammarskjöld 3241, Vitacura Santiago, Chile

<sup>4</sup>USDA, Agricultural Research Service, Subarctic Agricultural Research Unit, Fairbanks, Alaska

<sup>5</sup>Current address: Monsanto Arizona Cotton Research Center, 749 West Ash Ave, Casa Grande, AZ 85193, USA

\*Corresponding author; E-mail: [silvia.rondon@oregonstate.edu](mailto:silvia.rondon@oregonstate.edu)

### ABSTRACT

In 2 consecutive years, ground beetles (Coleoptera: Carabidae) were sampled in commercial organic and conventional potato fields, using pit fall traps. Four conventional and 3 organic potato fields were surveyed to determine ground beetle taxa composition. In a related study, potato fields were assigned to 1 of 4 transitional systems to include: organic, no spray (usually referred to as no control and/or only OMRI certified control), IPM-conventional (directed control, after sampling) and conventional (broad spectrum pesticides) systems. Seven taxa of ground beetles were identified: *Anisodactylus kerbyi* Lindroth 1953, *Amara* sp., *Bembidion* sp., *Calosoma cancellatum* Eschscholtz 1833, *Calathus ruficollis* Casey 1920, *Calathus ingratus* Dejean 1828, and *Pterostichus adstrictus* Eschscholtz 1823. With the exception of *C. cancellatum*, which was not identified from organic fields in the first yr of this study, all taxa were collected in both organic and conventional potato fields both yr. Based on total number of specimens collected, *Bembidion* sp., *P. adstrictus* and *Amara* sp. represented 42.8% of the total specimens collected. The smallest number of ground beetles was collected from no spray and IPM-conventional systems. This study provides basic valuable information regarding beetles populations for growers making transition from conventional to organic potato production.

Key Words: biological control, conventional agriculture, generalist, ground beetles, organic agriculture, sustainable systems

### RESUMEN

En dos años consecutivos, la composición de taxa de carabidos (Coleoptera: Carabidae) fue estudiada en campos comerciales de papa orgánica y convencional. Cuatro campos comerciales convencionales de papa y tres campos comerciales orgánicos, fueron monitoreados usando trampas de caída. Trampas fueron usadas desde la siembra (mediados de abril) hasta antes de la quema del follaje (mediados de agosto). También, para determinar como los sistemas transicionales afectan las poblaciones de carabidos, cuatro sistemas fueron evaluados: orgánico, sistema orgánico de transición (donde solo se permiten aplicaciones de pesticidas aprobados por el OMRI), IPM-convencional (solo se permiten aplicaciones de pesticidas después del monitoreo), y convencional (se permite el uso de pesticidas de amplio espectro). Siete especie de carabidos fueron identificados: *Anisodactylus kerbyi* Lindroth 1953, *Amara* sp., *Bembidion* sp., *Calosoma cancellatum* Eschscholtz 1833, *Calathus ruficollis* Casey 1920, *Calathus ingratus* Dejean 1828, y *Pterostichus adstrictus* Eschscholtz 1823. Todas estas taxas fueron encontradas en campos orgánicos y convencionales, con la excepción de *C. cancellatum* que no fue encontrada en campos orgánicos en el primer año de este estudio. *Bembidion* sp., *P. adstrictus* y *Amara* sp. fueron las taxas más abundantes (42.8% del número total de taxa colectadas). Sistemas orgánico de transición y IPM-convencionales presentaron el número más bajo de carabidos colectados. Este trabajo provee información básica importante para agricultores que intentan convertir su producción convencional a orgánica o viceversa.

Palabras Clave: control biológico, agricultura convencional, generalistas, carábidos, orgánico, sistemas sostenibles

Generalist ground-dwelling predaceous arthropods are common inhabitants in agroecosystems; however, they are susceptible to changes in abiotic factors (e.g., temperature, humidity), vegetative community, prey availability, and agronomic practices such as fertilization (mainly nitrogen) and pesticide use (Witmer et al. 2003). Ground beetles (Carabidae) are a biologically diverse and ecologically important element of the ground-dwelling community. The vast majority of ground beetles are active predators (Harrison & Regnier 2003) feeding on mollusks, beetles, aphids, Hemiptera, Lepidoptera, and thrips, among others (Lovei & Sunderland 1996; Sunderland 2002; Prasad & Snyder 2004); they may also be omnivorous or partially herbivorous (Laub & Luna 1992; Clark et al. 1993; Kromp 1999).

Increasing plant diversity within agroecosystems is believed to increase predator abundance and diversity (Root 1973). Moreover, habitat manipulation can affect densities of natural enemies (Barbosa 1998; Landis et al. 2000); still at present, the connection between abundance of beneficials and pest suppression is unclear (Bommarco & Banks 2003) and in some cases, controversial (Clark et al. 2006). According to Lundgren et al. (2006), the ground beetle community can be directly or indirectly impacted by farm management practices. Ground beetle composition responds to soil amendments and soil management (Purvis & Curry 1984; Reichert & Bishop 1990). No-till fields can have higher density of predators as compared to conventional tilled fields (House & All 1981; House & Alzugaray 1989; Whitmer et al. 2003; Clark et al. 2006; Shearin et al. 2007). Döring & Kromp (2003) suggested that ground beetles are less affected under organic farming systems rather than conventional, likely because some taxa are less or more susceptible to insecticides (Brust et al. 1985; Holland & Luff 2000). Koss et al. (2005) conducted an on farm study in Washington State where they compared predator and pest communities in potato fields treated with broad spectrum, selective or organic insecticides finding minor differences. Despite all this body of knowledge, few studies have been conducted specifically to determine the influence of transitional organic systems on ground-dwelling predators (Lundgren et al. 2006). In the United States, growers face a 3-yr transition from conventional to organic production that may have an effect on the biological community composition in transitional areas (Lundgren et al. 2006). In traditional intensive conventional agriculture systems as the one in the lower Columbia Basin of eastern Oregon, there are challenges that ground-dwelling communities may face such as limited food sources (since pests are being controlled), irrigation (eastern Oregon receives 203.2 mm of water per year), crop rotation (3 yr rotation) and other cropping practices. Growers with transitional and

newly certified organic fields need better information on the impact of their practices on communities of beneficial arthropods. Therefore, the objectives of this research were to study ground beetle taxa composition and relative abundance in conventional, transitional, and organic potato farming systems in eastern Oregon.

## MATERIALS AND METHODS

### Ground Beetle Survey in Organic and Conventional Potato Areas

In 2007 and 2008, ground beetles were surveyed in first yr commercial organic ( $n = 3$ ) and conventional potato fields ( $n = 4$ ) in the Boardman and Hermiston area located east of the Cascade Mountains in Oregon (Fig. 1). Insects were collected with pitfall traps; pitfall traps are widely used to collect soil arthropods in both agricultural and natural systems (Southwood 1978; Adis 1979; Weeks et al. 1997; Hansen & New 2005; O'Rourke et al. 2008). Ten pitfall traps per field were placed in a linear transect; traps were 15.2 m apart. Traps were placed in the rows to prevent flooding from irrigation. Typically potatoes are hilled, creating deep furrows between rows, and thus



Fig. 1. Map of eastern Oregon. The lower Columbia basin is one of the richest irrigated agriculture areas in the world. Circle shows the area where field plots were located (N 45.8411° W 119.2917°; N 45.8356° W 119.6992°).

traps consisted of double 12 oz cups (Solo Cup, Highland Park, Illinois). First a cup with holes for drainage was buried at soil level. A second cup filled about  $\frac{1}{4}$  full of a water and 10% soap solution was placed inside the first cup. A piece of wood was used as a partial lid to provide "shelter" and to reduce the likelihood of flooding. Each field was 50.6 ha in size under pivot irrigation with irrigation equivalent to 762 mm per season. Fields within each group followed similar pest management practices. Organic fields were treated with azadirachtin [neem extract at 7 oz/acre (512 mL/ha)] and *Bacillus thuringiensis* subspecies *tenebrionis* at 800 L of Bt per hectare, targeting aphids complex and Colorado potato beetle (*Leptinotarsa decemlineata* Say), respectively. Conventional fields were treated with selective pesticides for noctuid larvae (several taxa) and beet leafhopper (*Circulifer tenellus* Baker). Insecticides used were pyrethroids insecticides, imidacloprid and spinosad. Both yr, all these pests were present during the time of the experiment.

Traps were serviced weekly; samples were brought to the Oregon State University Hermiston Agricultural Research and Extension Center (OSU-HAREC) irrigated agricultural entomology laboratory in Hermiston for sorting, pinning, and preliminary identification. Insects were subsequently shipped to the USDA, ARS Entomology laboratory in Fairbanks, Alaska for identification confirmation. Ground beetles were identified primarily by A. Hagerty, using Lindroth (1969), Bousquet & Laroche (1993), and Ball & Bousquet (2001) identification keys. Afterwards, identifications were confirmed by George E. Ball (University of Alberta, Edmonton, Alberta, Canada), Robert Davidson (Carnegie Museum of Natural History, Pittsburgh, Pennsylvania) and Christopher J. Marshall (Oregon State Arthropod Collection, Corvallis, Oregon). Voucher specimens were kept in OSAC, OR (<http://osac.science.oregonstate.edu/>). Subsequent identifications were completed at the OSU-HAREC entomology laboratory by comparison with identified specimens.

Traps were monitored from planting (mid-Apr) until right before vine-killing (mid-Aug). Sampling was difficult before Apr or after Aug, because the fields followed normal farming practices (i.e. tillage, weed control, potato vine-killing before harvest).

#### Small Scale Transition Plots

Potato fields under a transition program from conventional to organic farming at the OSU-HAREC (N 45° 50' 26" W 119° 17' 17") in Hermiston Oregon were sampled in 2008. The transition to organic production was initiated in 2004 on a site that had been under conventional crop production the previous seasons. In our study, treatments were: 1) organic, 2) semi-organic (only OM-

RI or Organic Materials Review Institute certified sprays), 3) IPM-conventional (following IPM program including scouting and selective use of pesticides) and 4) conventional (broad spectrum pesticides). All seeds were certified. Potatoes were planted in May of 2008 and seed was provided by Three Mile Canyon Farms (RDO, Boardman, Oregon). Only OMRI approved materials for organic production were used in organic and semi organic treatments. Semi organic treatment used OMRI materials for pest control plus commercial fertilizer for plant nutrition (<http://www.ams.usda.gov/AMSV1.0/nop>). All 4 treatments have been cropped the same since their establishment in 2004. Crop rotation was berseem clover (*Trifolium alexandrinum* L.) in summer of 2004 followed by mustard (*Brassica juncea* (L.) Czern.) in the fall of 2004. Thirty three tons of compost per acre (67.4 t/ha) was applied to all treatments except conventional. Forage peas were planted in fall of 2005 but they did not germinate because of the late planting date. Green peas were planted in the spring of 2006; compost was then applied at 20 tons per acre (44.9 t/ha). Sudan grass was planted in summer of 2006. Winter wheat was planted in fall of 2006. Blood meal and chicken manure were used to fertilize the organic treatment and conventional fertilizers were used for the semi organic, IPM-conventional and conventional treatments. Standard potato practices were carried out in the conventional treatment with pesticide sprays occurring on a weekly basis (mainly for diseases like early blight). The IPM-conventional received 50% of the treatments that the conventional plots received. The semi organic received 75% of the treatments that the conventional plots received. The organic and semi organic received no pest control. Sixteen experimental plots (0.86 × 1.27 m each) were established and the 4 management intensity treatments were randomly assigned to the plots in a randomized complete block design with 4 replications. Blocks were separated by alleyways 3.47 m wide (4 potato rows). Plots within a block were separated by 6.1 m. The alleyways received no fertilizer or pest control (organic or conventional) but were always planted (Table 1). These plots were established by the 4th author for nitrogen best management practices studies for potato production. Small plots are standard for this type of evaluations (Hutchinson & Mylavarapu 2003).

#### Evaluation of Transition Practices in Ground Beetle Populations

Pitfall traps were set to determine the possible effect of farming transition on the diversity of ground beetles. Hence, 2 pitfall-traps per block per treatment (organic, semi organic, IPM-conventional and conventional) were placed 0.25 m apart. The set up procedure was similar than

TABLE 1. TRANSITIONAL PLOTS IN HERMISTON, OREGON. SIXTEEN EXPERIMENTAL PLOTS (0.86 × 1.27 M EACH) WERE ESTABLISHED AND THE FOUR MANAGEMENT INTENSITY TREATMENTS WERE RANDOMLY ASSIGNED TO THE PLOTS IN A RANDOMIZED COMPLETE BLOCK DESIGN WITH 4 REPLICATIONS. THESE PLOTS WERE ESTABLISHED FOR NITROGEN BEST MANAGEMENT PRACTICES STUDIES FOR POTATO PRODUCTION.

Treatments or Farm practices	Year			
	2004	2005	2006	2007
Organic	Berseem clover in the summer + 33 tons of compost + blood mean and chicken manure	Forage peas + blood mean and chicken manure	Green peas in the spring + 20 tons/a of compost + blood mean and chicken manure	Peas + blood mean and chicken manure
	+ mustard in the fall		+ Sudan grass in the summer + Winter wheat in the fall	
Semi-organic	Berseem clover in the summer + 33 tons/a of compost + mustard in the fall	Forage peas + conventional fertilizer (75% less than the conventional treatment)	Green peas in the spring + 20 tons/a of compost + conventional fertilizer (75% less than the conventional treatment)	Peas + conventional fertilizer (75% less than the conventional treatment)
			+ Sudan grass in the summer + Winter wheat in the fall	
IPM-conventional	Berseem clover in the summer + 33 tons/a of compost + mustard in the fall	Forage peas + conventional fertilizer (50% less than the conventional treatment)	Green peas in the spring + Sudan grass in the summer + conventional fertilizer (50% less than the conventional treatment) + Winter wheat in the fall	Wheat + conventional fertilizer (50% less than the conventional treatment)
Conventional	Berseem clover in the summer + mustard in the fall	Peas + conventional fertilizer	+ Sudan grass in the summer + conventional fertilizer + Winter wheat in the fall	Winter wheat + conventional fertilizer

potatoes

potatoes

potatoes

potatoes



the one described above under “Ground beetle survey”. Traps were checked weekly and samples were sorted in the Hermiston Irrigated Agricultural Entomology Program Laboratory.

Data Analysis

The field survey and transitional system experiment was analyzed as repeated measures ANOVA since each trap was sampled multiple times during the season. ANOVA results are presented in footnotes to the tables (SAS 2007). The data were transformed by  $\sqrt{\text{capture} + 0.5}$  before analysis to even the mathematical function of each point in the data set. Significant differences (5% level of probability) in the mean number of adults per trap between yr, taxa, and systems were determined for taxa for which at least 50 specimens were collected during the 2 yr sampling period (O'Rourke et al. 2008). The Simpson's diversity index was used to quantify the ground beetle biodiversity. This index takes into account the number of insects present as well as abundance of each species.

RESULTS AND DISCUSSION

Table 2 lists all specimens collected during the study. Seven taxa of ground beetles from 6 genera were identified: *Anisodactylus kerbyi* Lindroth, *Amara* sp., *Bembidion* sp., *Calosoma cancellatum* Eschscholtz, *Calathus ruficollis* Casey, *Calathus ingratus* Dejean, and *Pterostichus adstrictus* Eschscholtz. With the exception of *C. cancellatum*, which was not collected in 2007, all taxa were recovered both yr. However, the relative density of specimens varied between yr with 86% of the carabids specimens recovered during 2007 as compared to 2008 (14%). This is a substantial reduction in the relative density possibly due a combination of weather conditions and manage-

ment practices. Both factors are not manageable factors in applied field studies. *Bembidion* sp., *P. adstrictus*, *A. kerbyii*, and *Amara* sp. were the most abundant taxa (Table 2), representing 90% of the total carabid collected.

There were no significant differences in *Bembidion* densities in organic versus conventional systems, but there were significant differences in individual species such as *P. adstrictus* and *Amara* populations. *Amara* spp. was more abundant in conventional areas, while *P. adstrictus* was more abundant in organic areas (Table 3). The reasons for the differences in relative population densities cannot be explained with current biological knowledge of these taxa in Oregon. Possibly, the differences observed are influenced by taxa sensitivity to management practices. Considering that *Amara* populations were higher than *P. adstrictus* early in the season (Fig. 2) we speculate that further studies should consider possible competition between taxa. The seasonal population dynamics for the 3 most abundant taxa varied. One taxon, *Bembidion* sp. displayed high densities early in the season (May 1), but decline sharply to reach the lowest population density observed by these 3 taxa (Figure 2C). All taxa were collected the first week traps were deployed, suggesting that adult carabid activity in Oregon starts before May. Contrary to *Bembidion* sp., *Amara* sp. densities were the lowest recorded early in the season with a sharp increase in densities, reaching a peak of 0.45 insects per trap per 15 days period (Fig. 2A). *Pterostichus adstrictus* maximum densities (0.4 insects per trap per 15 days) were recorded on Jul 2 (Fig. 2B). All taxa were active toward the end of the potato season when traps were removed suggesting that further studies should consider earlier and latter collecting times; however sampling outside the planting season in commercial fields poses difficulties due to standard cultural practices.

TABLE 2. TOTAL NUMBER OF GROUND BEETLES COLLECTED IN TWO COMMERCIAL FIELDS IN EASTERN OREGON, 2007-2008.

Taxa	2007		2008	
	Number	%	Number	%
<i>Anisodactylus kerbyi</i> Lindroth	62	10	38	38
<i>Amara</i> spp.	135	22	2	2
<i>Bembidion</i> spp.	231	38	26	26
<i>Calosoma cancellatum</i> Eschscholtz	0	0	3	3
<i>Calathus ruficollis</i> Casey	3	< 1	4	4
<i>Calathus ingrates</i> Dejean	29	5	6	6
<i>Pterostichus adstrictus</i> Eschscholtz	150	25	20	20
Total Carabids	610	100	99	100

ANOVA ( $F = 3.50$ ;  $df = 6, 14$ ;  $P$  value = 0.0635); OD Simpson's  $D = 0.248$ ,  $SID = 0.752$ ;  $SRI = 4.032$ . The simpson's diversity index is a measure of diversity. In ecological terms, it is often used to quantify the biodiversity of a habitat taking into account the number of species present, as well as the abundance of each species.

TABLE 3. MEAN +/- SE NUMBER OF CARABIDS COLLECTED IN THE 2007-2008 SEASONS IN CONVENTIONAL AND ORGANIC FIELDS IN EASTERN OREGON.

Site	Carabids		
	<i>Amara</i> sp.	<i>Pterostichus adstrictus</i>	<i>Bembidion</i> sp.
	Mean +/- SE	Mean +/- SE	Mean +/- SE
Organic	0.2 +/- 0.01 a	0.5 +/- 0.07 b	0.5 +/- 0.12 a
Conventional	0.5 +/- 0.07 b	0.09 +/- 0.03 a	0.4 +/- 0.06 a

Means followed by the same letter within columns are not significantly different (Fisher's LSD,  $\alpha = 0.05$ ); Site ( $F = 17.865$ ,  $df = 4$ ,  $P < 0.001$ ); *Amara* ( $F = 41.726$ ,  $P < 0.001$ ); *Pterostichus adstrictus* ( $F = 26.795$ ,  $P < 0.001$ ); *Bembidion* ( $F = 0.842$ ,  $P = 0.359$ ).




Fig. 2. Mean number of *Amara* sp. (A), *Pterostichus adstrictus* (B), and *Bembidion* sp. (C) (+ SEM) in eastern Oregon, 2007.

Previous studies comparing ground beetles in organic and conventional cropping systems reported

TABLE 4. MEAN +/- SE AND MEAN SEPARATION OF CARABIDS IN EASTERN OREGON.

Site	Ground beetles
	Mean +/- SE
Organic	0.6 +/- 0.1 ab
Semi organic	0.2 +/- 0.1 a
IPM-conventional	0.3 +/- 0.1 a
Conventional	0.7 +/- 0.2 b

Means followed by the same letter within columns are not significantly different (Fisher's LSD,  $\alpha = 0.05$ ); Site ( $F = 2.656$ ,  $df = 3$ ,  $P = 0.049$ ); total carabids ( $F = 2.656$ ,  $P = 0.049$ ).

greater abundance under organic conditions (Clark et al. 2006; Dritschild & Wanner 1980), an observation not supported by our data (Table 3). In our study, ground beetles densities were greater in conventional than in organic systems. Since organic systems do not use synthetic inputs and use cover crops, in theory, organic settings should have been more favorable for ground beetle establishment and development. Dritschilo & Wanner (1980) reported that organic farms had twice the number of ground beetles found on conventional farms, but had the same level of diversity. In eastern Oregon, the high input potato system was suspected to have an effect on ground beetle population, but this study demonstrates that may not be case (Table 4). Probably the additional disturbance under the organic and no-spray treatments affected insect establishment more than conventional settings. To our knowledge, this is the first long-term report on taxa composition and population dynamics of ground beetles from organic and transition plots in Oregon potato fields. Information on ground beetles' taxa composition, distribution, population dynamics, dispersal, and biology is needed to understand their role as predators and seed consumers in potato fields and organic systems. This study provides some of the information necessary to guide future research such as taxa composition, seasonality, and a framework for sampling. Additional research is needed to study the ecology of the dominant taxa and their relationships with cultural practices.

## ACKNOWLEDGMENTS

Special thanks to Amanda Smith, Rebekah Adams, Dustin Keys, and Nick Teraberry for collecting, sorting and pinning ground beetles. The authors also thank Greg Harris, Emilie Hagewood, and R. D. Offutt, for providing potato seed and field sites in Boardman. The authors recognize the assistance of George E. Ball (University of Alberta, Canada), Robert Davidson (Carnegie Museum of Natural History, Pittsburg, Pennsylvania), Christopher J. Marshall, (Oregon State Arthropod Collection, Corvallis, Oregon), and Derek Sikes (University of Alaska, Museum of the North) for taxonomical assistance. Thanks also to Arnold Appleby (OSU Crop and Soil Science) and Andrew Jensen (Washington Potato Commission) for reviewing early versions of the manuscript.

## REFERENCES CITED

- ADIS, J. 1979. Problems of interpreting arthropod sampling with pitfall traps. *Zool. Anz.* 202: 177-184.
- BALL, G. E., AND BOUSQUET, Y. 2001. Carabidae Latreill, 1810, pp. 32-132 *In* R. H. Arnett and M. C. Thomas [eds.], *American Beetles*, Vol. 1: Archostemata, Myxophaga, Adephaga, Polyphaga: Staphyliniformia. CRC Press, Boca Raton, London, New York, Washington, D.C.
- BARBOSA, P. 1998. Conservation biological control. Academic Press. New York. NY.
- BOMMARCO, R., AND BANKS, J. E. 2003. Scale as a modifier in vegetation diversity experiments: effects on herbivores and predators. *Oikos* 102: 440-448.
- BOUSQUET, Y., AND LAROCHELLE A. 1993. Catalogue of the Geadaphaga (Coleoptera: Trachypachidae, Rhysodidae, Carabidae including Cicindelini) of America north of Mexico. *Mem. Entomol. Soc. Canada* 167: 1-397.
- BRUST, G. E., STINNER, B. R., AND MCCARTEY, D. A. 1985. Tillage and soil insecticide effects on predator black cutworm (Lepidoptera: Noctuidae) interactions in corn agroecosystems. *J. Econ. Entomol.* 78: 1389-1392.
- CLARK, M., LUNA, J. M., STONE, N. D., AND YOUNGMAN, R. R. 1993. Habitat preference of generalist predators in reduced-tillage corn. *J. Econ. Entomol.* 28: 404-416.
- CLARK, S., SZLAVECZ, K., CAVIGELLI, M. A. AND PURRINGTON, F. 2006. Ground beetle (Coleoptera: Carabidae) assemblages in organic, no-till, and chisel-till cropping systems in Maryland. *Environ. Entomol.* 35(5): 1304-1312.
- DRITSCHILD, W., AND WANNER, D. 1980. Ground beetle abundance in organic and conventional corn fields. *Enviro. Entomol.* 9(5): 629-631.
- DÖRING, T. F., AND KROMP, B. 2003. Which carabid beetles benefit from organic agriculture? A review of comparative studies in winter cereals from Germany and Switzerland Agriculture. *Ecosyst. and Environ.* 98: 153-161.
- HANSEN, J. E., AND NEW, T. R. 2005. Use of barriers pitfall traps to enhance inventory surveys of epigeal Coleoptera. *J. Insect Conserv.* 9: 131-136.
- HARRISON, S. K., AND REGNIER, E. E. 2003. Post-dispersal predation of giant ragweed (*Ambrosia trifida*) seed in no-tillage corn, *Weed Sci.* 51: 955-964.
- HOLLAND, J. M., AND LUFF, M. I. 2000. The effects of agricultural practices on Carabidae in temperate agroecosystems. *Integ. Pest Mgt. Review* 5: 109-129.
- HOUSE, G. J., AND ALL, J. N. 1981. Carabid beetles in soybean agroecosystems. *Environ. Entomol.* 18: 302-307.
- HOUSE, G. J., AND ALZUGARAY, M. R. 1989. Influence of cover cropping and no-tillage practices on community composition of soil arthropods in North Carolina agroecosystems. *Environ. Entomol.* 10: 194-196.
- HUTCHINSON, C., AND MYLAVARAPU, R. 2003. Evaluation of nitrogen best management practices for potato production in northeast Florida, *In* R. Y. Yada [ed.], *Proc. XXVI IHC Potatoes healthy food for humanity*. Acta Horticult. 619: 279-283.
- KROMP, B. 1999. Carabid beetles in sustainable agriculture: are review on pest control efficacy, cultivation impacts, and enhancement. *Agric. Ecosys. Environ.* 74: 187-228.
- LANDIS, D. A., WRATTEN, S. D., AND GURR, G. M. 2000. Habitat management to conserve natural enemies of arthropods pests in agriculture. *Annu. Rev. Entomol.* 45: 175-201.
- LAUB, C. A., AND LUNA, J. M. 1992. Winter cover crop suppression practices and natural enemies of armyworm (Lepidoptera: Noctuidae) in no-till corn. *Environ. Entomol.* 21: 41-49.
- LINDROTH, C. H. 1969. The ground-beetles (Carabidae, Excl. Cicindelinae) of Canada and Alaska. *Entomologiska Sällskapet, Lund, Sweden*.
- LINDROTH, C. H., 1969. An analysis of the carabid beetle fauna of the refugium, pp. 195-210 *In* T. N. V. Karlstrom and G. E. Ball [eds.], *The Kodiak Island refugium: Its geology, flora, fauna, and history*. Ryerson Press; Toronto 262 p. + i-xii.
- LUNDGREN, J. G., SHAW, J. T., ZABORSKI, E. R., AND EASTMAN, C. E. 2006. The influence of organic transition systems on beneficial ground-dwelling arthropods and predation of insects and weed seeds. *Renew. Agric. Food Syst.* 21: 227-237.
- LÖVEI, G. L., AND SUNDERLAND, K. D. 1996. Ecology and behavior of ground beetles (Coleoptera: Carabidae) *Annu. Rev. Entomol.* 41: 231-256.
- O'ROURKE, M. E., LIEBMAN, M., AND RICE, M. 2008. Ground beetle (Coleoptera: Carabidae) assemblages in conventional and diversified crop rotation systems. *Environ. Entomol.* 37(1): 121-130.
- PRASAD, R. P., AND SNYDER, W. E. 2004. Predator interference limits fly egg biological control by a guild of ground-active beetles. *Bio. Control.* 31: 428-437.
- PURVIS, G., AND CURRY, J. P. 1984. The influence of weeds and farmyard manure on the activity of Carabidae and other ground-dwelling arthropods in a sugar beet crop. *J. Appl. Ecol.* 21: 271-283.
- REICHERT, S. E., AND BISHOP, L. 1990. Prey control by and assemblage of generalist predators: spiders in garden test system. *Ecol.* 71: 1441-1450.
- ROOT, R. B. 1973. Organization of a plant arthropod association in simple and diverse habitats: the fauna of collars. *Ecol. Monogr.* 43: 95-124.
- SAS INSTITUTE. 1997. SAS User's Guide. SAS Institute, Cary, NC, USA.
- SHEARIN, A. F., REBERG-HORTON, S. C., AND GALLANDT, E. R. 2007. Direct effects of tillage on the activity density of ground beetle (Coleoptera: Carabidae) weed seed predators. *Environ. Entomol.* 36: 1140-1146.



- SOUTHWOOD, T. R. E. 1978. Ecological methods with particular reference to the study of insect populations. Chapman & Hall, New York.
- SUNDERLAND, K. D. 2002. Invertebrate pest control by carabids, pp. 165-214 *In* J. M. Holland [ed.], The Agroecology of Carabid Beetles. Intercept, Andover, UK.
- WEEKS, R. D., AND MCINTYRE, N. E. 1997. A comparison of live versus kill pitfall trapping techniques using various killing agents. *Entomol. Exp. Appl.* 82: 267-273.
- WHITMER, J. E., HOUGH-GOLDSTEIN, J. A., AND PESEK, J. D. 2003. Ground-dwelling and foliar arthropods in four cropping systems. *Environ. Entomol.* 32: 366-376.