



**Chartocerus sp. (Hymenoptera: Signiphoridae) and
Pachyneuron crassiculme (Hymenoptera: Pteromalidae)
are Obligate Hyperparasitoids of Diaphorencyrtus
aligarhensis (Hymenoptera: Encyrtidae) and Possibly
Tamarixia radiata (Hymenoptera: Eulophidae)**

Authors: Bistline-East, Allison, and Hoddle, Mark S.

Source: Florida Entomologist, 97(2) : 562-566

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.097.0230>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

**CHARTOCERUS SP. (HYMENOPTERA: SIGNIPHORIDAE) AND
PACHYNEURON CRASSICULME (HYMENOPTERA: PTEROMALIDAE) ARE
OBLIGATE HYPERPARASITOIDS OF DIAPHORENCYRTUS ALIGARHENSIS
(HYMENOPTERA: ENCYRTIDAE) AND POSSIBLY TAMARIXIA RADIATA
(HYMENOPTERA: EULOPHIDAE)**

ALLISON BISTLINE-EAST AND MARK S. HODDLE*

Department Of Entomology, University Of California, Riverside, CA, 92521, USA

*Corresponding author; E-mail: mark.hoddle@ucr.edu

Supplementary material for this article in Florida Entomologist 97(2) (June 2014) is online
at <http://purl.fcla.edu/fcla/entomologist/browse>

ABSTRACT

Two species of suspected hyperparasitoids, *Chartocerus* sp. and *Pachyneuron crassiculme*, emerged from parasitized *Diaphorina citri* nymphs collected in Punjab Pakistan over 15-22 Apr 2013. Exposure tests conducted in quarantine on *D. citri* nymphs parasitized by *Tamarixia radiata* and *Diaphorencyrtus aligarhensis*, as well as unparasitized *D. citri* nymphs, confirmed that *Chartocerus* sp. and *P. crassiculme* are hyperparasitoids. Both *Chartocerus* sp. and *P. crassiculme* successfully reproduced on *D. aligarhensis*, with one instance of *P. crassiculme* reproducing on *T. radiata*. There was no emergence from unparasitized *D. citri*.

Key Words: choice test, no-choice test, quarantine

RESUMEN

Dos especies de hiperparasitoides sospechosos, *Chartocerus* sp. y *Pachyneuron crassiculme*, emergieron de ninfas parasitadas de *Diaphorina citri* recolectadas en Punjab Pakistán del 15 al 22 de abril del 2013. Las pruebas de la exposición realizada en cuarentena sobre ninfas de *D. citri* parasitadas por *Tamarixia radiata* y *Diaphorencyrtus aligarhensis*, así como ninfas de *D. citri* no parasitadas confirmaron que *Chartocerus* sp. y *P. crassiculme* son hiperparasitoides. Tanto *Chartocerus* sp. y *P. crassiculme* se reprodujeron con éxito sobre *D. aligarhensis*, con un caso de *P. crassiculme* reproducido sobre *T. radiata*. No hubo emergencia de parásitos de las ninfas de *D. citri* no parasitadas.

Palabras Clave: prueba de opción, prueba de elección, cuarentena

Asian citrus psyllid (ACP), *Diaphorina citri* Kuwayama (Hemiptera: Liviidae), was discovered in California USA in 2008. *D. citri* vectors 'Candidatus Liberibacter asiaticus', a putative causative agent of huanglongbing (HLB), a lethal disease of citrus (Hoffman et al. 2013; Wang & Trivedi 2013). HLB was detected in California in Mar 2012 (Leavitt 2012). To mitigate the threat posed by *D. citri*-HLB to California's citrus industry, a biological control program using *Tamarixia radiata* (Waterston) (Hymenoptera: Eulophidae) sourced from Pakistan was initiated (Hoddle 2012). *Diaphorencyrtus aligarhensis* (Shafee, Alam, and Agarwal) (Hymenoptera: Encyrtidae), a second parasitoid of *D. citri* also collected from Pakistan, is currently in quarantine at the University of California, Riverside (UCR). The purpose of this study was to confirm that *Chartocerus* sp. and *P. crassiculme*, both suspected hyperparasitoids, are not primary parasitoids of *D. citri*.

MATERIALS AND METHODS

Parasitized *D. citri* host material returned from Punjab Pakistan to quarantine at UCR (15-22 Apr 2013) yielded previously collected *T. radiata* and *D. aligarhensis*, along with several species of known (*Marietta leopardina* Motschulsky [Hymenoptera: Aphelinidae], *Aprostocetus* (*Aprostocetus*) sp. [Hymenoptera: Eulophidae] [Hoddle et al. 2013]) or suspected (*Chartocerus* sp. [Hymenoptera: Signiphoridae], *Pachyneuron crassiculme* Waterston [Hymenoptera: Pteromalidae] and *Psyllaphycus diaphorinae* [Hymenoptera: Encyrtidae]) hyperparasitoids.

To confirm that *Chartocerus* sp. (Fig. 1A. male, B. female) and *P. crassiculme* (Fig. 2A. male, B. female) are not primary parasitoids of *D. citri*, exposure trials using 10 sets of 4-7 *Chartocerus* sp. and 10 pairs of 1 male and 1 female *P. crassiculme* that emerged from material collected in Pakistan




Fig. 1. *Chartocerus* sp. male (A) and female (B). This figure is shown in color in the supplementary document in Florida Entomologist 97(2) (2014) online at <http://purl.fcla.edu/fcla/entomologist/browse>.

were rotated through each of 4 treatment types between 26 Apr and 24 May, 2013 in quarantine at UCR. It was not possible to reliably sex live *Chartocerus* sp., so this species was exposed in groups (assumed to contain at least 1 female each) unless a pair was otherwise observed mating. Exposure treatments consisted of: (A) nymphs parasitized by *T. radiata* ($n = 8$ replicates of 5-10 parasitized nymphs for *Chartocerus* sp. and 9 replicates of 5 parasitized nymphs for *P. crassiculme*), 5-9 days

post-exposure to *T. radiata*; (B) nymphs parasitized by *D. aligarhensis* ($n = 8$ replicates of 5-10 parasitized nymphs for *Chartocerus* sp. and 10 replicates of 5 for *P. crassiculme*), 10-14 days post-exposure to *D. aligarhensis*; (C) unparasitized third to fourth instar *D. citri* nymphs ($n = 9$ replicates of 5-10 unparasitized nymphs for *Chartocerus* sp. and 10 replicates of 5 nymphs for *P. crassiculme*); and (D) each of the 3 previously listed host types (A, B, and C) presented simulta-




Fig. 2. *Pachyneuron crassiculme* male (A) and female (B). This figure is shown in color in the supplementary document in Florida Entomologist 97(2) (2014) online at <http://purl.fcla.edu/fcla/entomologist/browse>.

neously in a choice cage ($n = 9$ replicates of 5-10 of each host type for *Chartocerus* sp. and 9 replicates of 5 of each host type for *P. crassiculme*).

Each replicate was comprised of host material for each treatment type exposed to a group of potential hyperparasitoids for 24 h each. Hosts were exposed sequentially in a different order for each replicate to prevent bias due to presentation order. Emergence rates of *T. radiata* ($n = 5$ parasitized nymphs on each of 10 cuttings) and *D. aligarhensis* ($n = 5$ parasitized nymphs on each of 10 cuttings) determined baseline mortality for primary parasitoids in the absence of hyperparasitoids. Unparasitized *D. citri* nymphs ($n = 5$ fourth instar nymphs on each of 10 plants) provided data on nymph mortality in the absence of hyperparasitoids. Mummies of *T. radiata* and *D. aligarhensis* used in exposure experiments were sourced from colonies maintained in quarantine at UCR.

Diaphorina citri nymphs parasitized by either *T. radiata* or *D. aligarhensis* for no-choice treatments were presented on small *Citrus volkameriana* cuttings. *Citrus volkameriana* seedlings grown in 114 mL Cone-tainers™ (SC7 Stubby, 3.8 cm diameter, Stewe and Sons Inc., Oregon) and infested with *D. citri* nymphs were used to expose unparasitized *D. citri* nymphs to *Chartocerus* sp. and *P. crassiculme*. Clear plastic vials (Thornton Plastic Co. 148 mL capacity, Salt Lake City, Utah) with three 12 mm diam ventilation holes covered with ultra-fine organza were inverted and placed over the top of the plant and fitted into the corresponding vial lid, which had a hole cut in the center to allow it to be fitted around the cone (Irvin et al. 2009).

Choice treatments were set up in 15 cm × 15.3 cm × 15.3 cm (h × w × d) clear plastic boxes (S&W Plastics, Riverside, California) with a 30 cm sleeve sewn from no-see-um netting (Skeeta Mosquito & Other Insect Protection Products, Bradenton, Florida). Unparasitized *D. citri* nymphs in Cone-tainers and *T. radiata*- and *D. aligarhensis*-parasitized nymphs on *C. volkameriana* cuttings in water were placed in the cage without ventilated vials on top to allow free access to all 3 host types simultaneously. After 24 h, each host type was enclosed with an inverted ventilated vial to contain all insects that emerged from each host type. All experiments were conducted in quarantine at UCR's Insectary and Quarantine facility, at 27 °C, 50% RH, and 14:10 h L:D. Replicates were observed daily after initial exposure, and total numbers of each emerged species were recorded per treatment.

RESULTS

No-choice treatments resulted in *Chartocerus* sp. reproducing successfully only on *D. aligarhensis* (Table 1). Mean emergence time for *Char-*

TABLE 1. EMERGENCE AND MORTALITY RATES FOR *CHARTOCERUS* SP. EXPOSED TO UNPARASITIZED THIRD AND FOURTH INSTAR *D. CITRI* NYMPHS, AND NYMPHS PARASITIZED BY *T. RADIATA*, AND *D. ALIGARHENSIS* IN NO-CHOICE AND CHOICE TREATMENTS.

	No-choice					Choice				
	Host	Total No. Exposed	% Host Emergence	% Parasitism	% Dead ⁵	% Missing ⁶	Total No. Exposed	% Host Emergence	% Dead ⁵	% Missing ⁶
<i>D. citri</i>		65	72.31% ¹	0.00%	9.23%	18.46%	65	81.54% ¹	13.85%	4.61%
<i>T. radiata</i>		67	67.16% ²	0.00%	26.87%	5.97%	62	66.13% ²	25.81%	8.06%
<i>D. aligarhensis</i>		60	33.33% ³	46.67% ⁴	16.67%	3.33%	65	53.85% ³	29.23%	16.92%

¹Percentage of *D. citri* adults that emerged from unparasitized nymphs.
²Percentage of *T. radiata* adults that emerged from parasitized nymphs.
³Percentage of *D. aligarhensis* adults that emerged from parasitized nymphs
⁴Percentage of hosts killed by parasitism. Actual number of host killed = 28, actual number of *Chartocerus* sp. adults emerged = 33, demonstrating 2 observed instances of superparasitism.
⁵Percentage of hosts found dead.
⁶Percentage of hosts unaccounted for at time of data collection.

tocerus sp. offspring from *D. aligarhensis* was 18.36 days \pm 2.34 (SE). *Pachyneuron crassiculme* produced progeny on *D. aligarhensis* and *T. radiata* in no-choice treatments, though parasitism was much higher on *D. aligarhensis* (Table 2). Mean emergence times for males and females were 12.83 days \pm 2.48 (SE) and 11.33 days \pm 2.05 (SE), respectively. *Pachyneuron crassiculme* had a single male emerge from *T. radiata* after 11 days. Emergence rates for control treatments of *T. radiata*, *D. aligarhensis*, and *D. citri* were 84%, 88%, and 88%, respectively (Table 3). *Chartocerus* sp. and *P. crassiculme* failed to reproduce on unparasitized *D. citri* nymphs.

Immature *D. aligarhensis* exposed to *Chartocerus* sp. in no-choice tests experienced 47% parasitism, 17% died from undetermined causes, 3% were unaccounted for, and 33% emerged as adult *D. aligarhensis*. In 20% of trials (i.e., 2 of 10 replicates) *Chartocerus* sp. exhibited superparasitism, with 11 adults emerging from 9 *D. aligarhensis* mummies in 1 replicate, and 6 adults emerging from 3 mummies in the second. In no-choice tests, immature *T. radiata* exposed to *Chartocerus* sp. exhibited 0% parasitism, 27% of mummies died from unknown causes, 6% disappeared, and 67% emerged as adult *T. radiata*.

In no-choice tests where *P. crassiculme* was exposed to immature *D. aligarhensis*, 28% of hosts were parasitized by *P. crassiculme*, 19% died from unknown causes, and 53% emerged as adult *D. aligarhensis*. On *T. radiata*, *P. crassiculme* successfully parasitized only 2% of host material (i.e., one host), 40% died from unknown causes, 7% were unaccounted for, and 51% emerged as adult *T. radiata*. Unknown mortality may be attributable to superparasitism, host feeding, or a combination of both by *P. crassiculme*.

There was no successful parasitism of any host in choice tests for either *Chartocerus* sp. or *P. crassiculme*. However, elevated mortality rates were observed for *T. radiata* (26% when exposed to *Chartocerus* sp.; 28% for *P. crassiculme*) and *D. aligarhensis* (29%; 13%). In comparison, control mortality for *T. radiata* and *D. aligarhensis* were < 13% in the absence of these hyperparasitoids. When viewed collectively, data from exposure trials demonstrates that *Chartocerus* sp. and *P. crassiculme* are obligate hyperparasitoids within the *D. citri*-*Tamarixia*-*Diaphorencyrtus* system. Immediately following the conclusion of trials, all *Chartocerus* sp. and *P. crassiculme* material was killed in quarantine and preserved in 95% ethanol. Voucher specimens were deposited in the Entomology Museum at UCR (Table 4).

Assuming *Chartocerus* sp. and *P. crassiculme* preferentially parasitize *D. aligarhensis* as these exposure trial data suggest, the frequency of *Chartocerus* sp. and *P. crassiculme* emergence in quarantine from material collected from Punjab Pakistan in April 2013 was significant in compar-

TABLE 2. EMERGENCE AND MORTALITY RATES FOR *PACHYNEURON CRASSICULME* EXPOSED TO UNPARASITIZED THIRD AND FOURTH INSTAR *D. CITRI* NYMPHS, AND NYMPHS PARASITIZED BY *T. RADIATA* AND *D. ALIGARHENSIS* IN NO-CHOICE AND CHOICE TREATMENTS.

Host	No-choice					Choice			
	Total No. Exposed	% Host Emergence	% Parasitism ⁴	% Dead ⁵	% Missing ⁶	Total No. Exposed	% Host Emergence	% Dead ⁵	% Missing ⁶
<i>D. citri</i>	50	68.00% ¹	0.00%	24.00%	8.00%	45	73.33% ¹	13.33%	13.33%
<i>T. radiata</i>	45	51.11% ²	2.22%	40.00%	6.67%	46	71.74% ²	28.26%	0.00%
<i>D. aligarhensis</i>	53	52.83% ³	28.30%	18.87%	0.00%	46	86.96% ³	13.04%	0.00%

¹Percentage of *D. citri* adults that emerged from unparasitized nymphs.
²Percentage of *T. radiata* adults that emerged from parasitized nymphs.
³Percentage of *D. aligarhensis* adults that emerged from parasitized nymphs.
⁴Percentage of *Pachyneuron crassiculme* adults that successfully emerged from parasitized hosts.
⁵Percentage of hosts found dead.
⁶Percentage of hosts unaccounted for at time of data collection.

TABLE 3. EMERGENCE RATES OF UNPARASITIZED THIRD AND FOURTH INSTAR *D. CITRI* NYMPHS AND NYMPHS PARASITIZED BY *T. RADIATA* AND *D. ALIGARHENSIS* IN CONTROL TREATMENTS NOT EXPOSED TO HYPERPARASITOIDS.

Host	Total No. Exposed	No. Adults Emerged	No. Dead Hosts ⁴	No. Missing Hosts ⁵
<i>D. citri</i>	50	44 ¹	1	0
<i>T. radiata</i>	50	42 ²	6	2
<i>D. aligarhensis</i>	52	46 ³	3	1

¹Total number of *D. citri* adults that matured from unparasitized nymphs.
²Total number of *T. radiata* adults that emerged from parasitized nymphs.
³Total number of *D. aligarhensis* adults that emerged from parasitized nymphs.
⁴Total number of hosts found dead.
⁵Total number of hosts unaccounted for at time of data collection.

TABLE 4. SPECIMEN ACCESSION NUMBERS FOR ALL SPECIES USED IN EXPOSURE TRIALS AND DEPOSITED IN THE ENTOMOLOGY MUSEUM AT THE UNIVERSITY OF CALIFORNIA RIVERSIDE.

Species	Accession No.
<i>D. citri</i> ¹	UCRC_ENT00334428
<i>T. radiata</i> ²	UCRC_ENT00334402-334418
<i>D. aligarhensis</i> ²	UCRC_ENT00334426-334427
<i>Chartocerus</i> sp. ²	UCRC_ENT00417173-00417182
<i>P. crassiculme</i> ²	UCRC_ENT00417183-00417187

¹Multiple individuals of Pakistani *D. citri* preserved in a single vial of 95% ethanol.
²Point-mounted individuals

ison to *D. aligarhensis* emergence rates. *Chartocerus* sp. (237 individuals reared), *P. crassiculme* (181), and *D. aligarhensis* (743) represented 20%, 16%, and 64% of material reared, respectively, within this complex. A total of 292 *T. radiata* were reared from April 2013 collections. Exposure trials suggest that the lower numbers of *T. radiata* obtained from Pakistan in April 2013 were not likely due to hyperparasitism.

ACKNOWLEDGMENTS

The authors would like to thank Roger Burks (UCR) for assistance with specimen mounting, photography, and identification. Serguei Triapitsyn (UCR) helped with specimen identification and deposition of voucher specimens. Anamaria Dal Molin (TAMU) identified

Chartocerus sp. Funding for this project was provided, in part, by funds issued by the California Department of Food and Agriculture’s Specialty Crops Program and the California Citrus Research Board.

REFERENCES CITED

HODDLE, M. S. 2012. Has the Asian citrus psyllid parasitoid, *Tamarixia radiata*, established in California? <http://civr.ucr.edu/blog/asian-citrus-psyllid-2/radiata-established-in-california/> (Last accessed 23 Aug 2013).
HODDLE, M. S., AND HODDLE, C. D. 2013. Classical biological control of Asian citrus psyllid with *Tamarixia radiata* in urban Southern California. *Citrograph* 4(2): 52-58.
HODDLE, C. D., HODDLE, M. S., AND TRIAPITSYN, S. V. 2013. *Marietta leopardina* (Hymenoptera: Aphelinidae) and *Aprostocetus* (*Aprostocetus*) sp. (Hymenoptera: Eulophidae) are obligate hyperparasitoids of *Tamarixia radiata* (Eulophidae) and *Diaphorencyrtus aligarhensis* (Hymenoptera: Encyrtidae). *Florida Entomol.* 96: 643-646.
HOFFMAN ET AL. 2013. Heat treatment eliminates ‘*Candidatus Liberibacter asiaticus*’ from infected citrus trees under controlled conditions. *Phytopathology* 103 (1), 15-22.
IRVIN, N. A., HODDLE, M. S., AND SUAREZ-ESPINOZA, J. 2009. The functional response of *Gonatocerus ashmeadi* and the ‘new association’ parasitoid *G. tuberculifemur* attacking eggs of *Homalodisca vitripennis*. *Environ. Entomol.* 38: 1634-1641.
LEAVITT, R. 2012. Huanglongbing confirmed in California. *Citrograph* 3(3): 8-9.
WANG, N., AND TRIVDEI, P. 2013. Citrus Huanglongbing: a newly relevant disease presents unprecedented challenges. *Phytopathology* 103(7): 652-665.