

First records of two mealybug species in Brazil and new potential pests of papaya and coffee

Authors: Culik, Mark P., Martins, David dos Santos, and Gullan, Penny J.

Source: Journal of Insect Science, 6(23) : 1-6

Published By: Entomological Society of America

URL: https://doi.org/10.1673/2006_06_23.1

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



First records of two mealybug species in Brazil and new potential pests of papaya and coffee

Mark P. Culik¹, David dos Santos Martins¹, and Penny J. Gullan²

¹ Instituto Capixaba de Pesquisa, Assistência Técnica e Extensão Rural – INCAPER, Rua Afonso Sarlo 160, CEP 29052-010, Vitória, Espírito Santo, Brasil

² Department of Entomology, University of California, One Shields Avenue, Davis, CA 95616-8584, USA

Abstract

Five mealybug (Hemiptera: Pseudococcidae) plant pest species: *Dysmicoccus grassii* (Leonardi), *Ferrisia malvastra* (McDaniel), *Ferrisia virgata* (Cockerell), *Phenacoccus tucumanus* Granara de Willink, and *Pseudococcus elisae* Borchsenius are recorded for the first time in the state of Espírito Santo, Brazil. These are the first records of *D. grassii* in Brazil, from papaya (*Carica papaya*, Caricaceae), and from coffee (*Coffea canephora*, Rubiaceae). *Ferrisia malvastra* is also newly recorded in Brazil, where it was found on *Bidens pilosa* (Asteraceae). *Ferrisia virgata* was collected from an unidentified weed and *Phenacoccus tucumanus* from *Citrus* sp. (Rutaceae). *Plotococcus capixaba* Kondo was found on pitanga (*Eugenia* cf. *pitanga*, Myrtaceae) and *Pseudococcus elisae* on *Coffea canephora*, which are new host records for these mealybugs.

Resumo

São registradas pela primeira vez a ocorrência das cochonilhas (Hemiptera: Pseudococcidae) *Dysmicoccus grassii* (Leonardi), *Ferrisia malvastra* (McDaniel), *Ferrisia virgata* (Cockerell), *Phenacoccus tucumanus* Granara de Willink e *Pseudococcus elisae* Borchsenius no estado do Espírito Santo. Destacam-se os registros de *Dysmicoccus grassii* e *Ferrisia malvastra* que são os primeiros no Brasil, sendo que *D. grassii* é o primeiro no mamoeiro (*Carica papaya*, Caricaceae) e em café conilon (*Coffea canephora*, Rubiaceae). A espécie *Ferrisia malvastra* foi encontrada em picão preto (*Bidens pilosa*, Asteraceae), *F. virgata* coletada de uma planta daninha não identificada, e *Phenacoccus tucumanus* em limoeiro galego (*Citrus* sp., Rutaceae). *Plotococcus capixaba* Kondo foi coletada em pitangueira (*Eugenia* cf. *pitanga*, Myrtaceae) e *Pseudococcus elisae* em *Coffea canephora*, sendo estas plantas consideradas, respectivamente, como novas hospedeiras dessas espécies de cochonilhas.

Keywords: *Dysmicoccus grassii*, *Ferrisia malvastra*, *Ferrisia virgata*, *Phenacoccus tucumanus*, *Planococcus minor*, *Plotococcus capixaba*, *Pseudococcus elisae*, *Coffea canephora*, *Carica papaya*, *Bidens pilosa*, *Eugenia* cf. *pitanga*

Correspondence: markculik3@yahoo.com, davidmartins@incaper.es.gov.br, pjgullan@ucdavis.edu

Received: 14.9.2005 | Accepted: 8.2.2006 | Published: 28.9.2006

Copyright: Creative Commons Attribution 2.5 (<http://creativecommons.org/licenses/by/2.5>)

ISSN: 1536-2442 | Volume 2006, Number 23

Cite this paper as:

Culik MP, Martins D dos S, Gullan PJ. 2006. First records of two mealybug species in Brazil and new potential pests of papaya and coffee. 6pp. *Journal of Insect Science* 6:23, available online: insectscience.org/6.23

Introduction

Mealybugs (Hemiptera: Pseudococcidae) are small, soft-bodied insects that feed by sucking plant sap. Adult females and nymphs are wingless and frequently covered in a white, powdery or mealy wax secretion. In addition, the margin of the body often has a series of white, lateral wax filaments that typically are most prominent posteriorly. Adult males, if present, are short-lived, non-feeding and rarely collected. Some mealybug species cause considerable economic damage to agricultural and horticultural plants (McKenzie 1967; Williams and Granara de Willink 1992; Miller et al. 2002 , 2005b). Plant damage by mealybugs results from the direct effects of sap removal and injection of toxins, as well as indirectly by honeydew contamination and associated sooty mold growth that decreases photosynthesis (Mibey 1997), and occasionally from the effects of transmitted plant viruses. Feeding damage may cause leaf yellowing, defoliation, reduced plant growth, and death of plants. The occurrence of honeydew and sooty mold may reduce the marketability of plant products such as fruits. Even if plant damage is not apparent, the mere presence of mealybugs can be a quarantine concern, adding to costs of production to prevent or eliminate their presence on plants and produce.

Papaya (*Carica papaya* L., Caricaceae) is grown throughout Brazil and cultivation of this fruit is especially important economically in the state of Espírito Santo (ES) (Alves 2003). However, information on the occurrence and distribution of mealybug species on crops such as papaya in Brazil is limited. Worldwide, at least nine species of mealybugs have been recorded as pests of papaya, of which five occur in Brazil (Culik et al. 2003) but none has been recorded previously on papaya in this country (Silva et al. 1968; Medina et al.1989). Furthermore, of 64 mealybug species recorded from Brazil (Ben-Dov et al. 2005), few have previously been identified from Espírito Santo

(Culik and Gullan 2005). Therefore, as part of efforts for development of integrated crop production, including papaya, in Brazil (Martins et al. 2003), mealybugs were collected during 2004 and 2005 from various plants in Espírito Santo to identify the species present in this area. Here we document new mealybug records from papaya and other plants in Espírito Santo.

Materials and Methods

Mealybug specimens were collected during surveys of the insect fauna of papaya orchards in Espírito Santo carried out by the Espírito Santo rural research and extension institute INCAPER (Instituto Capixaba de Pesquisa, Assistência Técnica e Extensão Rural) and when noticed on plants during fieldwork or other activities.

All mealybug specimens were slide-mounted for identification using the method outlined in Williams and Granara de Willink (1992) , except that xylene was used instead of clove oil. Voucher specimens of these insects are deposited in the arthropod collections of INCAPER, Vitória, Espírito Santo; and the Bohart Museum of Entomology, Department of Entomology, University of California, Davis, California, U.S.A.

Results

Five mealybug species not reported previously from Espírito Santo were found, as well as a new host for an additional species (Table 1). *Dysmicoccus grassii* (Leonardi) was first observed infesting *C. papaya* fruits of seven plants in a commercial orchard in December 2004 and five additional papaya fruits infested with *D. grassii* were collected from the same location a few days later (eggs, nymphs and males as well as adult female mealybugs were present on the fruit). In addition, *D. grassii* was obtained from the inflorescence of coffee plants of *Coffea canephora* (Rubiaceae) in February 2005.

Table 1. Records of mealybugs (Hemiptera: Pseudococcidae) collected from Espírito Santo (ES), Brazil

Mealybug species	Collection location	Collection date	Host Plant
<i>Dysmicoccus grassii</i> (Leonardi)	Aracruz, ES	10 Dec. 2004, 16 Dec. 2004	<i>Carica papaya</i> , fruit
<i>D. grassii</i>	Linhares, ES	21 Feb. 2005	<i>Coffea canephora</i> inflorescence
<i>Ferrisia malvastra</i> (McDaniel)	Linhares, ES	21 Feb. 2005	<i>Bidens pilosa</i> , roots
<i>Ferrisia virgata</i> (Cockerell)	Manguinhos, Serra, ES	9 Jan. 2005, 4 Apr. 2005	unidentified weed (cf. <i>Spermacoce</i> sp.) leaf/stem
<i>Phenacoccus tucumanus</i> Granara de Willink	Manguinhos, Serra, ES	24 Oct. 2004	<i>Citrus</i> sp., leaf
<i>Planococcus minor</i> (Maskell)	Sooretama, ES	1 Feb. 2005	<i>Coffea canephora</i> , inflorescence
<i>P. minor</i>	Linhares, ES	21 Feb. 2005	<i>Bidens pilosa</i> , roots
<i>Plotococcus capixaba</i> Kondo	Manguinhos, Serra, ES	28 May 2005	pitanga (<i>Eugenia</i> cf. <i>pitanga</i>), foliage
<i>Pseudococcus elisae</i> Borchsenius	Aracruz, ES	24 Feb. 2005	<i>Coffea canephora</i> inflorescence
<i>Ps. elisae</i>	Manguinhos, Serra, ES	19 Apr. 2005	unidentified weed, leaf

A single specimen of *Pseudococcus elisae* Borchsenius was identified from the inflorescence of *C. canephora* in February 2005, and an additional specimen of this species was collected from an unidentified weed in another municipality in April 2005. In February 2005, *Planococcus minor* (Maskell) was collected from the roots of *Bidens pilosa* growing in a coffee field and was also collected from the inflorescence of *C. canephora*, confirming the presence of this mealybug species as a potential pest of coffee in Espírito Santo (Santa Cecília et al. 2002).

Specimens of *Ferrisia malvastra* (McDaniel) were collected from the roots of *Bidens pilosa* (Asteraceae) growing near a coffee field and two collections of *Ferrisia virgata* (Cockerell) were made three months apart from the stem or leaves of an unidentified weed from the yard of a home. In the same yard, two specimens of *Phenacoccus tucumanus* Granara de Willink were taken from foliage of a “limão galego” tree (*Citrus* sp., Rutaceae) and *Plotococcus capixaba* Kondo was collected from the foliage of a heavily infested pitanga bush (*Eugenia* cf. *pitanga*, Myrtaceae).

Although *F. virgata* and *Ph. tucumanus* have been found previously in Brazil (Silva et al. 1968; Williams and Granara de Willink 1992), these are the first records of these species in Espírito Santo, and the first record of *Pl. capixaba* from pitanga. The records of *D. grassii* and *F. malvastra* are the first from Brazil. This is also the first report of *D. grassii* from papaya, and the first reports of *D. grassii* and *Ps. elisae* from *C. canephora*.

Discussion

Dysmicoccus grassii (synonym *D. azalon* Williams) is considered to be of Neotropical origin, but it was originally described from the Canary Islands (Williams and Granara de Willink 1992). It has been found in Africa, Europe and North America (Ben-Dov 2005a; Miller and Miller 2002; Miller et al. 2005b) and recently was reported for the first time in southern Asia, including on *Theobroma cacao* (Williams 2004). Although *D. grassii* is widely distributed in the Neotropics, it has not previously been recorded in Brazil. *D. grassii* is polyphagous with agriculturally important hosts including mango, pineapple, coffee, and cacao (Ben-Dov 2005a) and it is considered to be a pest of bananas in the Canary Islands (Williams and Granara de Willink 1992) and in Nigeria (Matile-Ferrero and Williams 1995). Watson and

Kubiriba (2005) have suggested that it could become more widespread in Africa.

The present work is the first record of papaya and *C. canephora* as hosts for *D. grassii*. On *C. canephora*, this mealybug was found in the inflorescence. *D. grassii* is similar taxonomically to the species *D. texensis* Tinsley (also recorded as *D. bispinosus* Beardsley) which occurs from the Neotropics through to the southern USA on a range of hosts (Ben-Dov 2005a), including on the roots of coffee in Brazil (Williams and Granara de Willink 1992; Santa-Cecília et al. 2002). As far as we know, *D. grassii* has never been collected on the roots of its host plants. On *C. papaya*, *D. grassii* was only observed on fruits and other parts of the plants did not appear to be infested. The infestation of *D. grassii* appeared to be concentrated near the peduncle of the fruits, and yellowing and rot in the area of the peduncle of the unripe (green), mealybug infested papaya fruit also was noted. Eggs, nymphs and males as well as adult female mealybugs were observed on the infested fruit indicating the suitability of papaya as a host for this mealybug. Adult females of *D. grassii*, as observed in this study (Figure 1A), are pinkish, densely covered with white, powdery wax and have 16 pairs of short, thick wax filaments (less than a quarter of the length of the body) laterally on the body and one pair of long terminal wax filaments (about half as long as the body).

Ferrisia virgata (striped mealybug) is found throughout the world on a very broad range of host plants including species in many agriculturally important families such as Cucurbitaceae, Fabaceae (=Leguminosae), Musaceae, Myrtaceae, Arecaceae (=Palmae), Rubiaceae, Rutaceae, and Solanaceae (Ben-Dov 2005a). *F. virgata* is a common pest on many plants, is a vector of *Cacao swollen-shoot badnavirus* (CSSV) in Africa and *Piper yellow mottle virus* (PYMV) in India (Bhat et al. 2003; Ben-Dov 2005a), and is reported to be a pest of *C. papaya* in Micronesia (Nafus et al. 1999). This species is widespread in Brazil and has been noted previously in the states of Bahia, Pará, Paraíba, Rio Grande do Norte, Rio de Janeiro, and São Paulo (Silva et al. 1968) but apparently it is not known to be an important pest in this country (Anonymous 2003). Many other potential host plants of *F. virgata* (mango, papaya, pineapple etc.) were present at the site in Manguinhos where *F. virgata* was collected in this study but, although observations were made on some of these potential hosts and several other mealybug species were




Figure 1. Typical appearance of adult female mealybugs: A, *Dysmicoccus grassii* on papaya, Aracruz, ES, December 2004; B, *Phenacoccus tucumanus* from citrus, Serra, ES, 24 October 2004 (photograph, MPC).

found on such plants (Culik and Gullan 2005), *F. virgata* was observed only on a herbaceous weed in Manguinhos. However, the species was also found on the foliage of a mixed group of potted ornamentals in Vitória, ES, in April 2005. The appearance of live *F. virgata* (Miller et al. 2005a ; Osborne 2005) is distinctive: adult females generally appear grayish, covered with white powdery wax and a pair of dark stripes (or rows of spots) dorsally (as the common name implies), with the body relatively elongate, tapering posteriorly, and terminating in a pair of long wax filaments.

Ferrisia malvastra (malvastrum mealybug) is a widespread and polyphagous species (Ben-Dov 2005a). Although it is not commonly known to be a significant plant pest, it has been confused with *F. virgata* and prior to recognition of *F. malvastra* as distinct from *F. virgata*, *F. malvastra* may have been misidentified as *F. virgata* (Ben-Dov 2005a). It is considered to be a pest in the US (Miller et al. 2002) and is of concern as a potential agricultural pest in Israel (Ben-Dov 2005b). This species is parthenogenetic and formerly was known as the uniparental form of *F. virgata* and then as *F. consobrina* Williams and Watson (e.g., Williams and Watson 1988 ; Williams and Granara de Willink 1992). Live *Ferrisia malvastra* and *F. virgata* look very similar, but the former has a more rounded body (PJJ, personal observation) and tends to be common on herbs and shrubs (Ben-Dov 2005b). The painting by Mary Foley Benson purported to be an infestation of *F. virgata* on lantana (McKenzie 1967 , color plate X) is probably *F. malvastra* as the collection data match specimens now known to be *F. malvastra* and the accompanying taxonomic drawing resembles *F. malvastra* (PJJ, unpublished).

Phenacoccus tucumanus is known only from South America where it has been found on only a few hosts: several *Citrus* species, an unidentified weed, and Brazilian pepper, *Schinus terebinthifolius* (Anacardiaceae) (Williams and Granara de Willink 1992), a plant native to Brazil and cultivated to a limited extent in this country. Adult females of *Ph. tucumanus*, as observed in this study (Figure 1B), are densely covered with white, powdery wax, with an oval body surrounded by ~18 pairs of short, thick wax filaments (less than a quarter of the length of the body). Although only a few specimens of *Ph. tucumanus* were found, this study confirms that this potential pest is present in Espírito Santo.

Platococcus capixaba is a species recently described (Kondo et al. 2005) and known only from Espírito Santo and São Paulo, Brazil, on *Leandra erinacea* (Melastomataceae) and jaboticaba, *Myrciaria jaboticaba* (Myrtaceae). This record of *Pl. capixaba* on pitanga confirms that this mealybug is likely to be more widespread and common than has previously been realized. On pitanga, *Pl. capixaba* has an appearance similar to that observed on jaboticaba. Nymphs and adult females are light yellow in color or covered with white powdery wax, with two long terminal wax filaments, and the insects form extensive clumps of wax on the undersurface of leaves. Eggs are laid under the wax, which apparently serves as protection for eggs and nymphs. Despite the noticeable infestation on pitanga, sooty mold and leaf drop, which appeared to be associated with this insect on jaboticaba (Kondo et al. 2005), was not apparent on pitanga.

Pseudococcus elisae (banana mealybug) has been recorded previously only from Central and South

America, and Florida, and is known from relatively few host plant species (Ben-Dov 2005a ; Miller et al. 2005b). However, its hosts do include agriculturally important species such as banana (*Musa paradisiaca*) and coffee (*Coffea arabica*) (Williams and Granara de Willink 1992 ; Gimpel and Miller 1996 ; Ben-Dov 2005a), and it is considered to be a pest of black pepper in Brazil as a vector of *Piper yellow mottle virus* (PYMV) (Duarte and Albuquerque 2003), a disease currently not present in Espírito Santo. *Ps. elisae* has been reported from *Coffea* sp. in Trinidad and from *C. arabica* in Colombia (Williams and Granara de Willink 1992) but this is the first record of this species on coffee in Brazil and the first report from *C. canephora*. Adult females of this mealybug are purplish-gray with the body surrounded by long, thin lateral and terminal wax filaments (Miller et al. 2005a).

Although *D. grassii*, *F. malvastra*, *F. virgata*, *Ph. tucumanus*, *Pl. capixaba*, and *Ps. elisae* are not currently known as important pests in Espírito Santo, the records reported here confirm that these potential pests are present in this state. Knowledge of the presence of these mealybugs may enable producers and researchers to consider the presence of such potential pests in production of crops and development of management methods to avoid practices that may favor development of damaging populations of these pests in the future. Cacao and black pepper are economically important crops in Espírito Santo and these records provide confirmation that *F. virgata* and *Ps. elisae*, which are vectors of *Cacao swollen-shoot badnavirus* and *Piper yellow mottle virus*, are present in this state, thus reaffirming the importance of quarantine measures to prevent the entry of these diseases to this area. Since *D. grassii* and *Ps. elisae* have not been noted previously as pests of papaya and *C. canephora*, further research on these species and their effects on these economically important crops is especially warranted.

Acknowledgements

We thank J.A. Ventura and B.E.V. Pacova, INCAPER, for helpful suggestions in reviewing this publication. We also thank T. Kondo, University of California at Davis, for assistance with slide-mounting the mealybug samples. The Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq provided financial support for this work. P.J. Gullan was supported in part by

Hatch funding from the California Agricultural Experiment Station.

References

- Alves F de L. 2003. A cultura do mamão *Carica papaya* no mundo, no Brasil e no Estado do Espírito Santo. In: Martins D dos S, Costa, A de FS da, editors. *A Cultura do Mamoeiro: Tecnologias de Produção*, 1: 13–34. Vitória, ES: INCAPER.
- Anonymous. 2003. *Consulta de Praga*. Ministério da Agricultura, Pecuária e Abastecimento –Coordenação-Geral de Agrotóxicos e Afins/DFIA/SDA. Available from: http://extranet.agricultura.gov.br/agrofit_cons/principal_agrofit_cons (16 June 2005).
- Ben-Dov Y. 2005a. ScaleNet, Pseudococcidae catalogue. <http://www.sel.barc.usda.gov/scalenet/scalenet.htm>. *Dysmicoccus grassii*: <http://www.sel.barc.usda.gov/catalogs/pseudoco/Dysmicoccusgrassii.htm> (29 March 2005). *Dysmicoccus texensis*: <http://www.sel.barc.usda.gov/catalogs/pseudoco/Dysmicoccustexensis.htm> (6 September 2005). *Ferrisia malvastra*: <http://www.sel.barc.usda.gov/catalogs/pseudoco/Ferrisiamalvastra.htm> (22 August 2005). *Ferrisia virgata*: <http://www.sel.barc.usda.gov/catalogs/pseudoco/Ferrisiavirgata.htm> (5 April 2005). *Pseudococcus elisae*: <http://www.sel.barc.usda.gov/catalogs/pseudoco/Pseudococcuselisae.htm> (22 August 2005).
- Ben-Dov Y. 2005b. The malvastrum mealybug *Ferrisia malvastra* (Hemiptera: Coccoidea: Pseudococcidae), distribution, host plants and pest status in Israel. *Phytoparasitica* 33: 154-156.
- Ben-Dov Y, Miller DR, Gibson, GAP. 2005. ScaleNet, Scales in a Country Query Results. Available from: <http://www.sel.barc.usda.gov/scalecgi/region.exe?region=andfamily=Pseudococcidaeandcountry=BRAandgenus=andintro=Aanddetail=Noandsubunit=andregname=Neotropicalandctryname=andaction=Submit+Queryandquerytype=Country+Query> (2 March 2005).
- Bhat AI, Devasahayam S, Sarma YR, Pant RP. 2003. Association of a badnavirus in black pepper (*Piper nigrum* L.) transmitted by mealybug (*Ferrisia virgata*) in India. *Current Science* 84: 1547-1550.
- Culik MP, Gullan PJ. 2005. A new pest of tomato and other records of mealybugs (Hemiptera: Pseudococcidae) from Espírito Santo, Brazil. *Zootaxa* 964: 1-8.
- Culik MP, Martins D dos S, Ventura JA. 2003. *Índice de Artrópodes Pragas do Mamoeiro (Carica papaya L.)*. Vitória, Espírito Santo, Brazil: INCAPER, 48 pp.

- Duarte M de LR, Albuquerque FC de. 2003. Doenças e métodos de controle. In: *Cultivo da Pimenta-do-reino na Região Norte*. Embrapa Amazônia Oriental Sistemas de Produção, 1. Available from: http://www.cpatu.embrapa.br/pimenta_do_reino/paginas/doencasemetodosdecontrole.htm (31 August 2005).
- Gimpel WF, Miller DR. 1996. Systematic analysis of the mealybugs in the *Pseudococcus maritimus* complex (Homoptera: Pseudococcidae). *Contributions on Entomology, International* 2: 1-163.
- Kondo T, Gullan PJ, Ventura JA, Culik MP. 2005. Taxonomy and biology of the mealybug genus *Plotococcus* Miller and Denno (Hemiptera: Pseudococcidae) in Brazil, with descriptions of two new species. *Studies on Neotropical Fauna and Environment* 40: 213-227.
- Martins D dos S, Yamanishi OK and Tatagiba J da S. 2003. Normas Técnicas e Documentos de Acompanhamento da Produção Integrada de Mamão, (Documentos, 120). Vitória, ES: INCAPER, 60 pp.
- Matile-Ferrero D, Williams DJ. 1995. Recent outbreaks of mealybugs on plantain (*Musa* spp.) in Nigeria including a new record for Africa and a description of a new species of *Planococcus* Ferris (Homoptera, Pseudococcidae). *Bulletin de la Société Entomologique de France* 100: 445-449.
- McKenzie HL. 1967. Mealybugs of California with Taxonomy, Biology, and Control of North American Species (Homoptera: Coccoidea: Pseudococcidae). Berkeley: University of California Press, 526 pp.
- Medina JC, Bleinroth EW, Sigrist JMM, Martin ZJ de, Niscida ALAC, Baldini VLS, Leite RSSF, Garcia AEB. 1989. *Mamão: Cultura, Matéria-prima, Processamento e Aspectos Econômicos*, 2nd. ed. Campinas: ITAL, 367 pp.
- Mibey RK. 1997. Sooty moulds. In: Ben-Dov Y, Hodgson CJ, editors. *Soft Scale Insects: Their Biology, Natural Enemies and Control*, 7A: 275-290. Amsterdam and New York: Elsevier.
- Miller GL, Miller DR. 2002. *Dysmicoccus* Ferris and similar genera (Hemiptera: Coccoidea: Pseudococcidae) of the Gulf State Region including a description of a new species and new United States records. *Proceedings of the Entomological Society of Washington* 104: 968-979.
- Miller DR, Miller GL, Watson GW. 2002. Invasive species of mealybugs (Hemiptera: Pseudococcidae) and their threat to U.S. agriculture. *Proceedings of the Entomological Society of Washington* 104: 825-836.
- Miller GL, Miller DR, Limones EM, Gill RJ, Carlson RW. 2005a. The United States National Collection of Scale Insect Photographs. Available from: http://www.sel.barc.usda.gov:591/scale_slide/slide_frame.htm (22 March 2005).
- Miller DR, Miller GL, Hodges GS, Davidson JA. 2005b. Introduced scale insects (Hemiptera: Coccoidea) of the United States and their impact on U.S. agriculture. *Proceedings of the Entomological Society of Washington* 107: 123-158.
- Nafus D, Schreiner I, Moore A, Tudela A. 1999. *Insect pests of Micronesia*. Saipan: Northern Marianas College. Available from: <http://www.crees.org/plantprotection/AubWeb/bugweb/bugroot.htm> (16 June 2005).
- Osborne LS. 2005. Mealybugs. Available from: <http://www.mrec.ifas.ufl.edu/lso/Mealybugs.htm> (1 March 2005).
- Santa-Cecília LVC, Reis PR, Souza JC. 2002. About the nomenclature of coffee mealybug species in Minas Gerais and Espírito Santo States, Brazil. [Sobre a nomenclatura das espécies de cochonilhas-farinhas do cafeeiro nos Estados de Minas Gerais e Espírito Santo]. *Neotropical Entomology* 31: 333-334.
- Silva AGA, Gonçalves CR, Galvão DM, Gonçalves AJL, Gomes J, Silva M do N, Simoni L de. 1968. *Quarto Catálogo dos Insetos que Vivem nas Plantas do Brasil seus Parasitos e Predadores. Parte II – 1º Tomo*. Rio de Janeiro: Ministério da Agricultura, 622p.
- Watson GW, Kubiriba J. 2005. Identification of mealybugs (Hemiptera: Pseudococcidae) on banana and plantain in Africa. *African Entomology* 13: 35-47.
- Williams DJ. 2004. *Mealybugs of Southern Asia*. London: The Natural History Museum, 896 pp.
- Williams DJ, Watson GW. 1988. The Scale Insects of the Tropical South Pacific Region. Part 2: The Mealybugs (Pseudococcidae). London: CAB International Institute of Entomology, 260 pp.
- Williams DJ, Granara de Willink MC. 1992. *Mealybugs of Central and South America*. London: CAB International, 635 pp.