

Adelpha cocala didia (Nymphalidae: Limenitidinae) Feeding on Triumfetta semitriloba (Malvaceae)

Author: Tavares, Wagner De Souza

Source: The Journal of the Lepidopterists' Society, 70(3) : 251-252

Published By: The Lepidopterists' Society

URL: <https://doi.org/10.18473/107.070.0301>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ADELPHA COCALA DIDIA (NYMPHALIDAE: LIMENITIDINAE)
FEEDING ON TRIUMFETTA SEMITRILoba (MALVACEAE)**Additional key words:** Bignoniaceae, butterfly, Papilionoidea, Rhopalocera, Rubiaceae

Adelpha cocala Cramer (Lepidoptera: Nymphalidae: Limenitidinae) has six subspecies (*Adelpha cocala caninia* Fruhstorfer, *Adelpha cocala cocala* Cramer, *Adelpha cocala didia* Fruhstorfer, *Adelpha cocala lorzae* Boisduval, *Adelpha cocala orellanae* Neild, and *Adelpha cocala riola* Fruhstorfer), which are found in Brazil, Bolivia, Colombia, Guatemala, Peru, Suriname, Venezuela, and from Honduras to Panama (Fruhstorfer 1915, Willmott 2003b). The wingspan of these subspecies is around 50 mm (Willmott 2003a). Its larvae feed on *Alseis* spp., *Calycophyllum* spp., *Chimarris* spp., *Chomelia* spp., *Genipa* spp., *Pentagonia* spp., *Psychotria* spp., *Sabicea* spp., *Uncaria* spp., and *Warszewiczia* spp. (Rubiaceae), and *Malania* spp. (Olecaceae) (Beccaloni et al. 2008). *Adelpha cocala didia* occurs in the Brazilian states of Bahia, Espírito Santo, Mato Grosso, Minas Gerais, Santa Catarina, Rio de Janeiro, and São Paulo and in Brasília (Federal District) (Fruhstorfer 1915, Willmott 2003b). The aim of this study was to report a new host plant of *A. cocala didia* in the state of Minas Gerais, Brazil.

Eighty-two first-instar larvae were found defoliating leaves of 18 individuals of *Triumfetta semitriloba* Jacq. (Malvaceae) (around 45 cm in height) on May 2014, in secondary forest, in Viçosa, state of Minas Gerais, Brazil (20° 45' S × 42° 50' W and 655 m above sea level). Some branches were detached from these plants with the larvae and transferred to rearing cages (30 cm width × 30 cm length × 30 cm height) in the Laboratory for Biological Control of Insects (LCBI) of the Federal University of Viçosa (UFV) in Viçosa at 25 ± 1 °C, 12:12 (L:D) h photoperiod and 70 ± 10% R.H., where they remained until adults' emergence. Some females obtained from the collected larvae were sent to the Department of Zoology of the Federal University of Paraná (UFPR) in Curitiba, state of Paraná, Brazil, for identification by O.H.H. Mielke. They have been deposited in the entomological collection of the UFPR.

The number of pupae obtained from the collected larvae and of males and females from these pupae were evaluated. The pupa viability (number of obtained pupae ÷ number of collected larvae) × 100, adult viability (number of obtained adults ÷ number of obtained pupae) × 100, and sex ratio [(number of females ÷ (number of females + number of males))] were calculated.

Seventy-two adults emerged from the 78 obtained pupae—44 females and 28 males (sex ratio= 0.61, pupa viability= 95.12% and adult viability= 92.31%). No parasitoids were observed emerging from the larvae or pupae.

This record adds a new plant genus as host of *Adelpha*. This addition is not surprising because *Adelpha* is diverse and widely distributed across North and South America (Aiello 1984, Willmott 2003b). Previously, this subspecies had been reported in the states of Bahia, Espírito Santo, Mato Grosso, Santa Catarina, Rio de Janeiro, and São Paulo, and in Brasília (Federal District) at 1.344, 402, 1.804, 1.340, 355, 645, and 946 km away from Viçosa, respectively. Larvae of *Adelpha* feed on around 116 species of 42 genera of 22 botanical families (DeVries 1987, Brown Jr. 1992, Diniz & Moraes 1997, Freitas et al. 2001, Freitas 2006). *Adelpha cocala* is known to feed on 11 plant genera in Brazil and other parts of South America (Beccaloni et al. 2008). *Adelpha* was found feeding on Caprifoliaceae Juss. (Dipsacales) (Willmott 2003a) and on Rubiaceae Juss. (Gentianales) (Mullen et al. 2011). In Brazil, *Adelpha malea goyama* Schaus feeds on *Arrabidaea mutabilis* Bur. and K. Schum. (Lamiales: Bignoniaceae) in the state of Paraná (Freitas 2006). *Triumfetta semitriloba* is host of *A. cocala didia* in the Brazilian state of Minas Gerais.

ACKNOWLEDGMENTS

To Dr. Marília Cristina Duarte (Universidade de Mogi das Cruzes, Mogi das Cruzes, São Paulo State, Brazil) for identifying the plant and Dr. Olaf Hermann Hendrik Mielke for identifying the butterfly. To “Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)”, “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES)”, and “Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG)” for financial support.

LITERATURE CITED

- AIELLO, A. 1984. *Adelpha* (Nymphalidae): Deception on the wing. *Psyche* 91: 1–45.
- BECCALONI, G. W., A. L., VILORIA, S. K., HALL & G. S. ROBINSON. 2008. Catalogue of the hostplants of the Neotropical butterflies. *Trop. Lep. Res.* 18: 124–128.
- BROWN JR., K. S. 1992. Borboletas da Serra do Japi: Diversidade, hábitos, recursos alimentares e variação temporal, p.142–187. In Morellato, L. P. C. (ed.), *História natural da Serra do Japi. Ecologia e preservação de uma área florestal no sudeste do Brasil*. Campinas, Editora da Unicamp/Fapesp, 321p.
- DEVRIES, P. J. 1987. *The butterflies of Costa Rica and their natural history*. Princeton University Press, Princeton, New Jersey, xxii + 327p.

- DINIZ, I. R., & H. C. MORAES. 1997. Lepidopteran caterpillar fauna of cerrado host plants. *Biodivers. Conserv.* 6: 817–836.
- FREITAS, A. V. L. 2006. Immature stages of *Adelpha malea goyama* Schaus (Lepidoptera: Nymphalidae, Limenitidinae). *Neotrop. Entomol.* 35: 625–628.
- FREITAS, A. V. L., K. S., BROWN JR., & A. AIELLO. 2001. Biology of *Adelpha mythra* feeding on Asteraceae, a novel plant family for the Neotropical Limenitidinae (Nymphalidae), and new data on *Adelpha* “Species-Group VII”. *J. Lepid. Soc.* 54: 97–100.
- FRUHSTORFER. 1915. Gattung: *Adelpha* Hbn. in Seitz, Gross-Schmett. *Erde* 5: 510–512 (13 September), 513–520 (6 October), 521–528 (25 October), 529–533 (29 November).
- MULLEN, S. P., W. K., SAVAGE, N., WAHLBERG, & K. R. WILLMOTT. 2011. Rapid diversification and not clade age explains high diversity in Neotropical *Adelpha* butterflies. *Proc. R. Soc. B* 278: 1777–1785.
- WILLMOTT, K. R. 2003a. Cladistic analysis of the Neotropical butterfly genus *Adelpha* (Lepidoptera: Nymphalidae), with comments on the subtribal classification of Limenitidini. *Syst. Entomol.* 28: 279–322.
- WILLMOTT, K. R. 2003b. The genus *Adelpha*: Its systematics, biology and biogeography (Lepidoptera: Nymphalidae: Limenitidini). Gainesville, Scientific Publishers. viii + 322 pp.
- WAGNER DE SOUZA TAVARES (corresponding author), *Departamento de Fitotecnia, Universidade Federal de Viçosa, 36570-900, Viçosa, Minas Gerais state, Brazil.* SEBASTIÃO LOURENÇO DE ASSIS JÚNIOR, *Departamento de Engenharia Florestal, Universidade Federal dos Vales do Jequitinhonha e Mucuri, 39100-000, Diamantina, Minas Gerais state, Brazil.* CLAUBERT WAGNER GUIMARÃES DE MENEZES, *Instituto Federal do Norte de Minas Gerais - Campus Januária, 39480-000, Januária, Minas Gerais state, Brazil.* MARCUS ALVARENGA SOARES, *Departamento de Agronomia, Universidade Federal dos Vales do Jequitinhonha e Mucuri, 39100-000, Diamantina, Minas Gerais state, Brazil.* JOSÉ COLA ZANUNCIO, *Departamento de Entomologia/BIOAGRO, Universidade Federal de Viçosa, 36570-900, Viçosa, Minas Gerais state, Brazil; emails: wagnermaias@yahoo.com.br, assisjr_ento@yahoo.com.br, claubertmenezes@yahoo.com.br, marcusasoaes@yahoo.com.br, zanuncio@ufv.br*

Submitted for publication 4 December 2015; revised and accepted 19 March 2016.