

Immature Stages and Natural History of the Neotropical Satyrine *Pareuptychia Ocirrhoe Interjecta* (Nymphalidae: Euptychiina)

Authors: Freitas, André V. L., Barbosa, Eduardo Proença, and Marín, Mario Alejandro

Source: The Journal of the Lepidopterists' Society, 70(4) : 271-276

Published By: The Lepidopterists' Society

URL: <https://doi.org/10.18473/lepi.70i4.a4>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

IMMATURE STAGES AND NATURAL HISTORY OF THE NEOTROPICAL
SATYRINE *PAREUPTYCHIA OCIRRHAE INTERJECTA* (NYMPHALIDAE: EUPTYCHIINA)ANDRÉ V. L. FREITAS^{1,2*}, EDUARDO PROENÇA BARBOSA¹ AND MARIO ALEJANDRO MARÍN¹¹ Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, SP, Brazil² Museu de Zoologia, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, SP, Brazil

* Corresponding author

ABSTRACT. The present paper describes the immature stages of the common Neotropical satyrine butterfly *Pareuptychia ocirrhoe interjecta* (R.F. d'Almeida, 1952). The solitary eggs are white and round, turning black 4 to 6 hours after oviposition. The four solitary larval instars are predominantly green and feed on grasses (Poaceae), including *Setaria* in nature and several other native and introduced species in captivity. The pupa is short and smooth and entirely green. Except for the black eggs, the immature stages are similar to those of other forest species of Euptychiina. The most remarkable and unique characteristic of *P. ocirrhoe interjecta* is the shiny black eggs, a possible synapomorphy for *Pareuptychia* not known in any other Euptychiina.

Additional key words: Atlantic Forest, *Cepheptychia*, *Godartiana*, *Taydebis*, *Zischkaia*

MATERIALS AND METHODS

The Euptychiina (Nymphalidae: Satyrinae) is one of the largest and most diverse butterfly subtribes, contributing a significant portion of Neotropical butterfly diversity (Peña et al. 2010, Marín et al. 2011). With over 400 recognized species (Lamas, 2004), Euptychiina butterflies occur in virtually all habitats and vegetation types from sea level to over 3500 m (DeVries 1987, Marín et al. 2011). However, despite this large species richness, immature stages have been described for a very few species and genera of Euptychiina (DeVries 1987, Murray 2001, Freitas et al. 2016). For the genus *Pareuptychia*, for example, immature stages have never been described in detail, and available information includes only textual descriptions in DeVries (1987) and pictures of larvae and pupae in Janzen and Hallwachs (2015).

Species of *Pareuptychia* are associated with forested habitats across the Neotropics (DeVries 1987), and they may be abundant. The genus is very homogeneous, and its eight described species (Nakahara et al. 2016) are very similar in wing pattern and morphology, suggesting that this is a monophyletic group (except maybe for *Pareuptychia lydia* (Cramer, 1777)). However, because of similarity in wing patterns, species limits and identities are not well established and the genus needs to be studied from both morphological and molecular perspectives (Marín et al. in prep.). In addition, knowledge about the immature stages of *Pareuptychia* could be of help in understanding the taxonomy of this group.

In this context, the present paper offers a detailed description of the immature stages of *Pareuptychia ocirrhoe interjecta* (R.F. d'Almeida, 1952) and compares them with those of other neotropical Euptychiina butterflies.

Study sites. Adults and immatures of *P. ocirrhoe interjecta* were studied in four different localities in São Paulo State, Southeastern Brazil: 1) Reserva Biológica Municipal da Serra do Japi, Jundiá (900–1100 m; 23°13'S, 46°57'W); 2) ARIE Mata de Santa Genebra, Campinas (600–620 m; 22°49'S, 47°6'W); 3) Fazenda Santa Elisa, IAC, Campinas (630–650 m; 22°51'S, 47°5'W); 4) Parque Estadual Xixová-Japuí, São Vicente (20–200 m; 23°59'S, 46°23'W).

Sampling and rearing of immature stages. Fertile eggs were obtained from wild-captured females confined in plastic bags warmed by a 40W bulb and provided with leaves of several species of native and introduced grasses. Larvae of *P. ocirrhoe interjecta* and of the additional species mentioned in the discussion section were reared in plastic containers cleaned daily, with fresh plant material provided every two or three days (following Freitas 2007). Data were recorded on behavior and development time for all stages. Dry head capsules and pupal cases were retained in glass vials. Immature stages were fixed in Kahle-Dietrich solution (Triplehorn & Johnson 2005) when the number of specimens was sufficient. Voucher specimens of the immature stages were deposited in the Museu de Zoologia “Adão José Cardoso” (ZUEC-AVLF), Universidade Estadual de Campinas, Campinas, São Paulo, Brazil.

Morphology. Measurements were taken and general aspects of morphology were observed using a Leica® MZ7.5 stereomicroscope equipped with a micrometric scale. Scanning electron microscopy (SEM) was conducted using a JEOL® JSM-5800 microscope (JEOL Ltd., Japan), and samples were critical-point dried in a Bal-tec® – CPD030 (Leica Microsystems, Germany), attached with double-sided




FIG. 1. Immature stages and adult of *Pareuptychia ocirrhoe interjecta*. **a, b.** egg, white and black; **c, d.** first instar, dorsal and lateral; **e.** light head capsule second instar, lateral; **f, g.** dark head capsule second instar, dorsal and lateral; **h.** third instar, dorsal; **i, j.** fourth (last) instar, lateral and dorsal; **k, l, m.** Pupa, lateral, ventral and dorsal; **n.** Adult male.




FIG. 2. Egg and first instar of *Pareuptychia ocirrhoe interjecta*. **a.** egg; **b.** Head capsule, general frontal view; **c.** Detail of frontal region; **d.** Detail of posterior region: white setae show microsetae and black seta show a pore; **e.** Early first instar, general lateral view.

tape to aluminum stubs, and coated with gold/palladium with a Bal-tec® – SCD050 sputter coater (Leica Microsystems, Germany). Egg size is presented as height and diameter, and head capsule size is the distance between the most external stemmata (as in Freitas 2007). Terminology for early stages descriptions followed García-Barros & Martín (1995) for eggs and Stehr (1987) for larvae and pupae.

RESULTS

Description of immature stages. The following descriptions and measurements are based on material reared from one female from Serra do Japi, Jundiaí, São Paulo (material reared from three other localities gave similar results).

Egg (Figs. 1a,b – 2a). White, round, turning black 4 to 6 hours after oviposition. Height and diameter 0.94 – 0.96 mm; duration 5 – 6 days (n = 8).

First instar (Figs. 1c,d – 2b-e). Head capsule width 0.64 – 0.70 mm (mean = 0.66 mm; SD = 0.026 mm; n = 5); head scoli 0.08 – 0.10 mm (mean = 0.09 mm; SD = 0.008 mm; n = 5). Head black, with enlarged chalazae, bearing a pair of short scoli on vertex, each with two long narrow black setae. Third stemma larger than other stemmata. Body light green, with white longitudinal stripes, turning dark green after first meal; caudal filaments very short. Legs and prolegs light green. Setae light green from T1 to A7, black from A8 to A10, all dorsal and subdorsal clubbed at tip (Fig. 2e). Maximum length 5 mm. Duration 6 – 7 days (n = 8).

Second instar (Figs. 1e,f,g). Head capsule width 0.90 – 1.00 mm (mean = 0.94 mm; SD = 0.052 mm; n = 5); head scoli 0.40 – 0.46 mm (mean = 0.43 mm; SD = 0.022 mm; n = 5). Head green with two short dark brown scoli on vertex (Fig. 1e,g); in some individuals head is mostly dark brown (Figs. 1f,g). Body green, striped longitudinally with white; caudal filaments short. Legs and prolegs light green. Maximum length 9 mm. Duration 4 – 8 days (n = 8).

Third instar (Fig. 1h). Head capsule width 1.38 – 1.46 mm (mean = 1.45 mm; SD = 0.056 mm; n = 5); head scoli 0.60 – 0.66 mm (mean = 0.62 mm; SD = 0.024 mm; n = 5). Head green, with two diverging very short scoli on vertex; these are reddish with a posterior dark brown line extending to head capsule. Body light green with a conspicuous dark green dorsal stripe and several additional longitudinal markings; caudal filaments short. Legs and prolegs light green. Maximum length 15 mm. Duration 7 – 9 days (n = 6).

Fourth (last) instar (Figs. 1i,j). Head capsule width 2.18 – 2.20 mm (mean = 2.20 mm; SD = 0.016 mm; n = 4); head scoli 0.74 – 0.82 mm (mean = 0.78 mm; SD = 0.034 mm; n = 4). Head green, with two diverging short scoli with reddish apex on vertex. Body light green with a conspicuous dark green dorsal stripe and several fine longitudinal stripes; caudal filaments short and reddish on apex. Legs and prolegs green. Maximum length 25 mm. Duration 7 – 10 days (n = 6).

Pupa (Figs. 1k,l,m). Short and smooth; entirely green; short rounded ocular caps; cremaster light green; dorsal abdomen smooth without projections. Total length 10.0 – 10.5 mm (n = 5). Duration 9 – 14 days (n = 4).

Behavior and natural history. *P. ocirrhoe interjecta* (Fig. 1n) is common in several environments in the Atlantic Forest from sea level to 1000 m altitude. Oviposition behavior was not observed in the field, but two solitary eggs, two larvae and one pupa were collected in the field in an unidentified native species of *Setaria* (Poaceae) in a shaded narrow trail inside the forest. Eggs were laid singly in the laboratory and larvae

easily accepted the same hostplant mentioned above. Larvae were solitary in all instars and did not exhibit cannibalistic behavior (several larvae of different instars were reared together in small pots). Adults were easily observed along forest edges and in clearings flying low in the understory, rarely rising above 2 m above the ground. Flight was erratic and fast and, when disturbed, butterflies performed unpredictable aerial maneuvers with alternating flashes of white and dark, which makes them difficult to capture in flight. Adults feed on fermenting fallen fruits, feces and several other decaying substances, being never observed visiting flowers. However, there are reports of adults feeding on extrafloral nectar (Barbosa 2013; AVLf pers. obs.).

DISCUSSION

The immature stages of *P. ocirrhoe interjecta* are quite morphologically simple and similar to those of several other species of Neotropical Euptychiina: larvae lack body scoli, present short head horns and caudal filaments and pupae are short and smooth. The entirely green last instar, although not rare in Euptychiina, is apparently uncommon; most known last instars of small forest Euptychiina are predominantly brown or have color tones that make them cryptic on the background of dead leaves and stems (Singer et al. 1983, DeVries 1987, Murray 2001, Kaminski & Freitas 2008, Janzen & Hallwachs 2015). Species with similar green last instars and/or pupae includes other species of *Pareuptychia* (AVLF unpublished; Janzen & Hallwachs 2015), *Cepheuptychia cephus* (Fabricius, 1775), *Taydebis peculiaris* (A. Butler, 1874), *Zischkaia pacarus* (Godart, [1824]), *Godartiana muscosa* (A. Butler, 1870) (Figure 3) and *Forsterinaria pronophila* (Freitas & Peña 2006). Although *Cepheuptychia* and *Taydebis* are genera related to *Pareuptychia* in part of the “*Pareuptychia* clade” (Peña et al. 2010), the similarity in color pattern is not related to phylogenetic relationships. For example, other forest species belonging to the “*Pareuptychia* clade” present predominantly non-green last instars, including *Chloreuptychia arnaca* (Fabricius, 1776), *Megeuptychia antonoe* (Cramer, 1775) (Janzen & Hallwachs 2015), *Splendeuptychia furina* (Hewitson, 1862) and *Splendeuptychia doxes* (Godart, [1824]) (Freitas, unpublished). The greenish last instars of *Satyrotaygetis satyrina* (H. Bates, 1865) are quite distinct from those of *Pareuptychia*, especially in the longer, strongly diverging head horns.

The most remarkable and unique characteristic of the immature stages of *P. ocirrhoe interjecta* is the shiny black eggs. This trait, present also in other species of *Pareuptychia* (DeVries 1987, AVLf unpublished), is unique among all known Euptychiina (however, color




FIG. 3. Immature stages of Euptychiina (all from Brazil). **a, b.** *Cepheuptychia cephus*, fourth (last) instar and pupa (both lateral), Upper Juruá River, Marechal Thaumaturgo, Acre; **c, d.** *Taydebis peculiaris*, fourth (last) instar and pupa (both lateral), Santa Virgínia, São Luís do Paraitinga, São Paulo; **e, f.** *Zischkaia pacarus*, third instar, dorsal and lateral, Morro Grande, Cotia, São Paulo; **g, h.** *Godartiana muscosa*, fourth (last) instar and pupa (both lateral), Serra do Japi, Jundiaí, SP.

change in fertile eggs is not unique in *Pareuptychia*, and has been reported in *Calisto* Hübner, [1823] (Sourakov 1996)). Although the possible adaptive significance of this striking egg color is unknown, the similarity of these black eggs to the parasitized eggs of other species of Euptychiina is evident (Freitas et al. pers. obs.). Since it is widely known that visual cues are important for ovipositing butterflies (e.g. Rausher 1979, Williams & Gilbert 1981, Shapiro 1981, Freitas & Oliveira 1996, Sendoya et al. 2009), this opens the possibility that such cues are equally important for ovipositing females of egg parasitoids, which are known to show a low preference by black eggs (Lobdell et al. 2005). In this context, the black eggs of *Pareuptychia* could mimic parasitized eggs, thus reducing parasitoid oviposition and decreasing egg parasitism. Field and laboratory studies, however, are needed to test this hypothesis.

ACKNOWLEDGEMENTS

Thanks go to Ronaldo B. Francini and Keith S. Brown for their help with some pictures of the Figure 3. To Keith Willmott and Andrei Sourakov for their valuable suggestions in the submitted manuscript. AVLF acknowledges support from FAPESP (Biota-Fapesp grants 2011/50225-3, 2012/50260-6 and 2013/50297-0) and from the Brazilian Research Council - CNPq (fellowship 302585/2011-7 and 303834/2015-3). MAM acknowledges FAPESP under grant (2014/16481-0) for a graduate fellowship and Colciencias under grant (No. 1118-521-28378). EPB acknowledges FAPESP under grant (2012/03750-8) for a graduate fellowship. This publication is part of the RedeLep 'Rede Nacional de Pesquisa e Conservação de Lepidópteros' SISBIOTA-Brasil/CNPq (563332/2010-7) and from the National Science Foundation (DEB-1256742).

LITERATURE CITED

- BARBOSA, E. P. 2013. The fruit-feeding butterfly *Pareuptychia ocirrhoe* (Nymphalidae: Satyrinae) feeding on extrafloral nectaries. *J. Lepid. Soc.* 67: 64–65.
- DEVRIES, P. J. 1987. The butterflies of Costa Rica and their natural history. Papilionidae, Pieridae, Nymphalidae. Princeton, Princeton University Press, 327p.
- FREITAS, A. V. L. 2007. A new species of *Moneuptychia* Forster (Lepidoptera: Satyrinae, Euptychiina) from the Highlands of Southeastern Brazil. *Neotrop. Entomol.* 36: 919–925.
- FREITAS, A. V. L. & P. S. OLIVEIRA. 1996. Ants as selective agents on herbivore biology: effects on the behaviour of a non-myrmecophilous butterfly. *J. Animal Ecol.* 65: 205–210.
- FREITAS, A. V. L. & C. PEÑA. 2006. Description of genus *Guaianaza* for "*Euptychia*" *pronophila* (Lepidoptera: Nymphalidae: Satyrinae) with a description of the immature stages. *Zootaxa* 1163: 49–59.
- FREITAS, A. V. L., J. Y. O. CARREIRA, J. P. SANTOS & E. P. BARBOSA. 2016. Immature stages and natural history of two species of *Forsterinaria* from southeastern Brazil (Lepidoptera: Nymphalidae). *Trop. Lep. Res.* 26: 13–18.
- GARCÍA-BARROS, E. & J. MARTÍN. 1995. The eggs of European satyrine butterflies (Nymphalidae): external morphology and its use in systematics. *Zool. J. Linn. Soc. London* 115: 73–115.
- JANZEN, D. H. & W. HALLWACHS. 2015. Dynamic database for an inventory of the macrocaterpillar fauna, and its food plants and parasitoids, of Área de Conservación Guanacaste (ACG), northwestern Costa Rica. <http://janzen.sas.upenn.edu>. Accessed 01 March 2016.
- KAMINSKI, L. A. & A. V. L. FREITAS. 2008. Immature stages of the butterfly *Magneuptychia libye* (L.) (Lepidoptera: Nymphalidae, Satyrinae). *Neotrop. Entomol.* 37: 169–172.
- LOBDELL, C. E., T-H YONG & M. P. HOFFMANN. 2005. Host color preferences and short-range searching behavior of the egg parasitoid *Trichogramma ostrinae*. *Entomol. Experiment. Appl.* 116: 127–134.
- MARÍN, M. A., C. PEÑA, A. V. L. FREITAS, N. WAHLBERG & S. I. URIBE. 2011. From the phylogeny of the Satyrinae butterflies to the systematics of Euptychiina (Lepidoptera: Nymphalidae): history, progress and prospects. *Neotrop. Entomol.* 40: 1–13.
- MURRAY, D. L. 2001. Immature stages and biology of *Taygetis* Hübner (Lepidoptera: Nymphalidae). *Proc. Entomol. Soc. Wash.* 103: 932–945.
- NAKAHARA, S., M. A. MARÍN & A. F. E. NEILD. 2016. A new species of *Pareuptychia* Forster, 1964 (Lepidoptera, Nymphalidae, Satyrinae) from the northwestern Amazon basin with characterisation of two potential synapomorphies for the genus. *Trop. Lepid. Res.* 26: 6–12.
- PEÑA, C., S. NYLIN, A. V. L. FREITAS & N. WAHLBERG. 2010. Biogeographic history of the butterfly subtribe Euptychiina (Lepidoptera, Nymphalidae, Satyrinae). *Zool. Script.* 39: 243–258.
- RAUSHER, M. D. 1979. Egg recognition: its advantage to a butterfly. *Animal Behav.* 27: 1034–1040.
- SENDOYA, S. F., A. V. L. FREITAS & P. S. OLIVEIRA. 2009. Egg-Laying Butterflies Distinguish Predaceous Ants by Sight. *Am. Nat.* 174: 134–140.
- SHAPIRO, A. M. 1981. The pierid red-egg syndrome. *Am. Nat.* 117: 276–294.
- SINGER, M. C., P. J. DEVRIES & P. R. EHRLICH. 1983. The *Cissia confusa* species-group in Costa Rica and Trinidad (Lepidoptera: Satyrinae). *Zool. J. Linn. Soc. London* 79: 101–119.
- SOURAKOV, A. 1996. Notes on the genus *Calisto*, with descriptions of the immature stages (Part I) (Lepidoptera: Nymphalidae: Satyrinae). *Tropical Lepid.* 7(1: Part 2): 91–112.
- STEHF, F. W. 1987. Order Lepidoptera. In Stehr FW, editor. Immature insects. Vol. 1. Dubuque (IA): Kendall-Hunt Publishing Company. p. 288–305.
- TRIPLEHORN, C. A. & N. F. JOHNSON. 2005. Borror and Delong's introduction to the study of the insects. 7th Edition. Belmont, C A. Thomson Brooks/Cole. 864p.
- WILLIAMS, K. S. & L. E. GILBERT. 1981. Insects as selective agents on plant vegetative morphology: egg mimicry reduces egg laying by butterflies. *Science* 212: 467–469.

Submitted for publication 10 March 2016; revised and accepted 4 April 2016.