

Advantage or Disadvantage: Is Asexual Reproduction Beneficial to Survival of the Tunicate, *Polyandrocarpa misakiensis*

Authors: Kawamura, Kazuo, and Fujiwara, Shigeki

Source: Zoological Science, 17(3) : 281-291

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/jsz.17.281>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

REVIEW

Advantage or Disadvantage: Is Asexual Reproduction Beneficial to Survival of the Tunicate, *Polyandrocarpa misakiensis*?

Kazuo Kawamura* and Shigeki Fujiwara

Laboratory of Cellular and Molecular Biotechnology, Faculty of Science, Kochi University, Japan

ABSTRACT—It has been believed that clonal propagation by asexual reproduction has serious disadvantages for long-term survival, because asexual reproduction seems not to remove harmful mutations, it seems not to give rise to genetic variations upon which evolution depends and it seems not to reset cell aging. In this article, we re-consider those arguments, by reviewing asexual reproduction of the tunicate, *Polyandrocarpa misakiensis*. Tracer experiments of bud formation and growth using morphological and chimeric phenotypes showed that the parental epithelial tissues surrounding the bud primordium do not enter the growing bud. It is possible, therefore, to assume that budding involves the purge of a large number of parental somatic cells and tissues. Unlike sexuals, asexuals do not carry out meiotic recombination nor gene shuffling that are two major sources of genetic variation, but we can show that in *P. misakiensis* at least two genes have significant redundancy and genetic variation even in a clonal colony. Telomerase expressed in germlines is thought to reset the molecular clock executed by telomere shortening. In our *Polyandrocarpa* cDNA projects, four out of about 2,000 cDNAs examined were matched with retroviral reverse transcriptase that is the catalytic subunit of telomerase, suggesting that telomerase might work in asexual reproduction. In *P. misakiensis*, dedifferentiation system is used to make new asexual generations. TC14 lectin plays an important role in the maintenance of multipotent but differentiated state of the formative tissue. It is antagonized by tunicate serine protease (TRAMP) that has striking mitogenic and dedifferentiation-inducing activities on the multipotent cells. This system would serve to delay aging of somatic cells. In conclusion, empirical arguments that asexual reproduction is disadvantageous to long-term life do not appear to be tenable to budding of *P. misakiensis*.

INTRODUCTION

The capacity to repair missing parts of the body is a general characteristic shared by every organism, although the extent of repair varies among species. The regenerative potential of some marine and freshwater invertebrates is so remarkable that a piece of tissues can reconstruct the whole body (Morgan, 1901). Asexual reproduction takes advantage of this potential for propagating individuals (Brien, 1968). It can be found in most phyla in the animal kingdom including Polifera, Cnidaria, Plathelminthes, Annelida, Bryozoa, Echinodermata, Enteropneusta and Chordata.

Asexual reproduction accompanies neither meiotic recombination nor shuffling of male and female genomes, as is usual with sexual reproduction (Barton and Charlesworth, 1998). It is carried out by multipotent somatic cells (cf., Nakauchi, 1982) instead of single germ cells, indicating that the colonial population shares genomic constitution. In the sense that clonal indi-

viduals are given off, asexual reproduction has resemblance to parthenogenesis in rotifers (Wurdak and Gilbert, 1977), aphids (Normark, 2000) and others. Actually, evolutionary geneticists regard both asexual reproduction and parthenogenesis as synonym (e.g., Wuethrich, 1998). Nonetheless, they are discernible from each other, because parthenogenesis begins with female (sexual) gametes, in which there is a good chance giving rise to genetic recombination and, consequently, genetic variation during oogenesis (Kabay and Gilbert, 1977). In this paper, the term, asexual reproduction, is limited to the narrow definition by which a new individual comes from somatic cells. For broader definition, if necessary, clonal reproduction will be used to refer to the natural creation of clonal individuals.

It has been generally believed that clonal reproduction has disadvantage to long-term survival. This is because clonal reproduction seems not to remove harmful mutations, it seems not to give rise genetic variation and it seems not to reset cell aging. In this paper, we would like to re-consider those classical problems that asexual reproduction addresses, based on our recent findings in the budding tunicate, *Polyandrocarpa misakiensis*.

* Corresponding author: Tel. +81-88-844-8313;
FAX. +81-88-844-8313.
E-mail: kazuk@cc.kochi-u.ac.jp

Purge of deleterious genes

Many evolutionary geneticists argue that the fitness (viability and reproductive success) of sexual reproduction is superior to that of asexual one (Barton and Charlesworth, 1998). They insist that sex serves to assemble beneficial variations and so it creates a well-adapted lineage in the face of a rapidly changing environment (e.g., Ishii *et al.*, 1989). Although genetic mutations are obviously the source of evolution, most mutations affecting fitness appear to be harmful rather than beneficial (Keightley *et al.*, 1998), leading to the extinction of offspring (Fig. 1A). One of the most important merits of sexual reproduction may be, therefore, to eliminate deleterious mutations rather than to accumulate beneficial mutations (Leigh, 1973). Now, we have plenty of hypotheses to explain how sex purges genetic scapegoats (Drake, 1991), although we do not know yet about the accurate rate at which sex removes harmful mutations. Gametogenesis may offer an opportunity to self-diagnose the integrity of genome, by sorting-out of mutations during meiosis (Fig. 1B). Then, harmful genes will be thrown away together with the carrier (gamete) in the trash.

On the other hand, clonal reproduction seems not to have such an opportunity of sorting-out of harmful mutations. The offspring inherit all of their bad genes and may pick up another through a new mutation (Wuethrich, 1998). In this way, mutations continue to be accumulated in both individuals and in the population (Sniegowski *et al.*, 1997; Taddei *et al.*, 1997). Those notions mentioned above are invariably correct in parthenogenesis where genetic clones are produced by female populations. In the rotifer, *Asplanchna sieboldi*, diploid females reproduce parthenogenetically via mitotic oogenesis (Gilbert, 1976). This way of reproduction does not give

any chances to throw harmful genes away, as a full set of genetic components should be transferred through female generations. Depending on nutritional conditions, some of the females produce eggs undergoing meiotic oogenesis (Kabay and Gilbert, 1997). The haploid eggs develop parthenogenetically into males instead of females. However, these males can scarcely transfer their selected genome to the offspring by fertilization, as the females produce diploid eggs. The accumulation of harmful mutations has a risk leading to the extinction of those populations and, finally, species. As discussed below, asexual reproduction, another mode of clonal reproduction, would not necessarily come into the same consequence as parthenogenesis.

Can asexual reproduction remove harmful mutations?

Tunicates belonging to the phylum Chordata are phylogenetically the highest organism that can reproduce asexually. Asexual animals of *P. misakiensis* were first collected in 1970 (Watanabe and Tokioka, 1972), and have ever since been cultured for 30 years in Japanese marine laboratories, including ours. Their mode of propagation is referred to as paleal budding (cf., Nakauchi, 1982). A paleal bud grows out from the parent body wall that consists of the outer and inner epithelia and mesenchymal cells intervening between them (Fig. 2) (Kawamura and Watanabe, 1982a,b; Kawamura and Nakauchi, 1984, 1986).

It is noteworthy that the bud begins with heterogeneous cell population instead of a single cell with which sexual reproduction starts. When harmful mutations occur spontaneously in some of those somatic cells, deficient cells may possibly be extinguished as a result of competition with remaining normal cells (Fig. 2A-a). In case they are alive but




Fig. 1. Fate of harmful mutations in sexual reproduction. (A) In sexual reproduction, when a zygote (top) has harmful mutations (red rod) all descendants have the same deficiencies (middle, bottom). (B) Harmful mutations (red rods) in germ lines (top) may be segregated during meiosis (middle), providing a chance to eliminate those mutations in the next generation (bottom).

have less proliferative activity, the probability that they spread to the clonal population becomes very low (Fig. 2A-b). In either cases, harmful mutations would not bring about serious consequence in the clonal life of budding animals. We wonder if deficient cells have similar or somewhat higher proliferative activity than normal cells (Fig. 2A-c). The former (deficient cells) may be substituted gradually for the latter,

thus some individuals of the next blastogenic generation being occupied by the offspring of the deficient cells (cf., Figs. 2B and 2C). The spreading (or the block of spreading) of harmful mutations in asexual reproduction depends largely on the amount of parental tissues that participate in bud formation.

In botryllid and polystyelid tunicates, both buds and adult animals are connected with the extra-individual vascular sys-




Fig. 2. Fate of harmful mutations in asexual reproduction. (A) In *P. misakiensis*, a bud consists of heterogeneous cell populations, the outer epidermal and inner endodermal epithelia and intervening mesenchymal cells. Deficient cells (red boxes) may be extinguished as a result of competition with normal cells (a). If deficient cells are alive but less proliferative than normal cells, they do not have serious effect on the life of clonal individuals (b). When mutant cells have the same or somewhat higher proliferative activity than normal cells, they can occupy a definite area of the animal (c). (B, C) In some cases mutations will be inherited to the next blastogenic generation (right), and in the other cases they will not be inherited (left).




Fig. 3. Bud growth in *P. stolonifera*. (A) Adult animal with buds (stolons), ventral view. The test vessel system develops around the basal margin of the body. a, ample; e, endostyle; h, heart; i, intestine; o, orifice; p, pyloric caecum; s, stolon; sti, stigmata; sto, stomach. Bar, 1 mm. (B) Daily growth of a bud. Vascular ampullae are omitted, but orifices (a1, a2, b1, b2) are plotted. As noted by a2 and b2, the parental epidermal tissue does not move toward the growing bud. (Reproduced from Kawamura and Watanabe, 1981).

tem by orifices (Newberry, 1965; Mukai *et al.*, 1978) (Fig. 3A). In *Botryllus*, *Botrylloides* and *Sympylegma*, the orifices are located at definite areas on the zooidal ventral surface. In *Polyandrocarpa*, on the other hand, they are distributed very abundantly around the basal margin where the bud primordium appears (Fig. 3A), making it easy to trace the epithelial tissue movement during bud growth. Strictly, the orifice is a good landmark of the epidermis but not the inner epithelium. However, many literatures have shown that both epithelia behave synchronously during budding (Berrill, 1941; Izzard, 1973; Kawamura and Nakauchi, 1986). In *P. stolonifera*, an adult animal of at most 5 mm in length produces very elongated buds (stolon) of more than 10 mm in length (Kawamura and Watanabe, 1981). Figure 3B shows the redrawing of their observation of bud growth with reference to orifices. The bud primordium was about 0.3–0.4 mm in width and 0.2 mm in height along the basal margin of the parent body wall. It had two orifices (a, b) at first that were soon divided, respectively, into two sub-orifices (a1, a2, b1, b2). Interestingly, neither a2 nor b2 entered the growing stolon. The orifices increased in distance exclusively between a1 and a2 or between b1 and b2. It is noteworthy that after the bud primordium stage parental epithelial tissues seem not to participate in the stolonial outgrowth.

This empirical observation mentioned above has been supported by chimera experiments (Kawamura and Watanabe, 1984). Two strains of *P. misakiensis*, *white spot* and *spotless*, are discernible from each other by the presence or absence of a white circular spot on the dorsal surface. The color phenotypes are determined by epithelial components of animals (Kawamura and Watanabe, 1984; Ishii *et al.*, 1993). Buds of the *spotless* strain was amputated and grafted into the body wall of the host adult animal of the *white spot* strain. The grafts expressed a *spotless* phenotype that was the same as the donor strain, even after they grew more than twice as large as the original size (Kawamura and Watanabe, 1984). This result is consistent with the notion that bud growth does not need the supply of parent (host) epithelial tissues.

Both results of *P. stolonifera* and *P. misakiensis* strongly suggest that epithelial components of a new blastogenic generation are all derived from a relatively small number of cells of the bud primordium. In other words, budding involves the removal of a large number of somatic cells and tissue of the parent animal. We would like to emphasize, therefore, that asexuals as well as sexuals have a good chance to purge harmful mutations.

Genetic variation in sexual reproduction

Genetic variation is undoubtedly the motive force of biological evolution. It can be promoted by chromosomal recombination (Barton and Charlesworth, 1998), shuffling of male and female genomes (Maynard Smith, 1978), neutral or beneficial mutations by nucleotide substitution (Kimura, 1967; Johnson, 1999), and genomic change of larger scale such as duplication or deletion (Ohno, 1970; Hughes, 1994; Wolfe and Shields, 1997; Holland *et al.*, 1994). Sexual reproduction has

the privilege of executing both the recombination and shuffling of genomes. It has been believed that those recombination and shuffling are beneficial by allowing favorable alleles to come together (Fisher, 1930). At the same time, however, it is also possible that favorable sets of genes having been accumulated through natural selection are broken up by genetic recombination (Barton and Charlesworth, 1998). The compromise is that recombination can be selectively advantageous if different gene combinations are favored in different generations and in different circumstances (Maynard Smith, 1978). In any case, sex offers some efficient methods for genetic variations without harmful mutations. Clonal reproduction either by parthenogenesis or asexual reproduction does not have such convenient methods, which is one of major reasons why clonal reproduction is thought to be disadvantageous to long-term survival (Barton and Charlesworth, 1998).

How fast are genetic variations fixed in the genome? The mutation rates seem to be determined by a balance between natural selection favoring lower mutation rates and opposing selective forces favoring higher mutation rates (Dawson, 1998). They vary widely among different species (Drake, 1991). In *Escherichia coli*, the rate has been estimated as 1.7×10^{-4} /haploid genome/generation (Kibota and Lynch, 1996). In eukaryotes, it is 0.84 in *Daphnia* (Deng and Lynch, 1997), 0.3–0.4 in *Drosophila melanogaster* (Mukai *et al.*, 1972; Keightley, 1994), about 0.1 in inbred population of mice (Caballero and Keightley, 1998), and about 5 in human (Kondrashov and Crow, 1993).

Does clonal population give rise to genetic variation?

Some aphids in Australia exhibit a complete absence of sexual reproduction. These wild-living parthenogenetic lineages have genetic variations in microsatellites and a few other inheritable components (Wilson *et al.*, 1999). A clone of laboratory-maintained parthenogenetic aphids was examined genetically over 32 generations (Lushai *et al.*, 1998). A putative germline mutation was noted once and somatic mutations were noted four times. Mitotic unequal crossing over seems to occur in X chromosome (Mandrioli *et al.*, 1999). A greenbug in the United States reproduces primarily by apomictic parthenogenesis, which is interrupted by a periodic sexual cycle. Shufra *et al.* (1997) have shown that the intergenic spacer of the rDNA is stable within parthenogenetic clones and that periodic sexual reproduction is a primary mechanism for the generation and maintenance of genetic variability.

In *P. misakiensis*, two examples of genetic variation have been known so far. One is tunicate C-type lectins of 14 kDa (TC14) (Suzuki *et al.*, 1990). C-type lectins are calcium-dependent carbohydrate-recognition proteins (Drickamer, 1993). They have a common sequence motif of 115 to 130 amino acid (aa) residues. TC14 consisting of 125 aa contains only the carbohydrate recognition domain that binds to D-galactose (Suzuki *et al.*, 1990) and D-fucose (Poget *et al.*, 1999). A cDNA encoding another type of TC14 has been isolated (Shimada *et al.*, 1995), named TC14-2 in relation to the original one that was renamed TC14-1. Two additional isoforms

Sniegowski (1997) predicts that optimal mutation rates will evolve only in asexual populations. In *Polyandrocarpa*, as mentioned earlier, buds are endowed with a small number of parental epithelial cells during their growth (cf., Figs. 2,3). If only normal cells are transferred to the bud from the parent that is a carrier of deficient genes, the mutation will be extinguished in the next asexual animal (Fig. 2A). On the other hand, if some of those cells transferred have a deficient gene, the mutation will be emphasized in the subsequent asexual generations, thus fixing genetic polymorphism very promptly in a clonal population. Therefore, our conclusion is that asexual

A

TC14-1 MVFTAILVAVIACGIPTSYA MDYEILFSDETMNYADAGTYCQSRGMALVSSAMRDSTMVKAILAFTEVKGHD
TC14-2 MVFTAILVAVIACGIPTSYA VDYDILFSDETMNYADAGAYCQSRGMVLVSSAMRDSTMVKAILDFTKDRRYN
TC14-3 MVFTAILVAVIACGIPASYA DNYEILMSGKAMTYSDAETYCESIGMTVAAKAIRDSTTVKAIVDFAAVTGKS
TC14-4 MIHGAILIAFLACGIPASYA GNYEILIADRSMNANYAKAFCDCKGMRLAAASLRDADTSQDILAYTASSGY

*.: ***:*.:*****:** :*:***:..*:*:* :*:.. ** :.: :*:** :*:**

TC14-1 YWVGADNLQDGAYNFLWNDGVSLPTSDSLWSPNEPSNPQSQWLVCQIWSKYNLLDDVGC GGARRVICEKELDD
TC14-2 YWVGADNLEK GAYNFLWNDGVSLPTSDSLWSPDEPSNPQSSSELVCQIWNRYNLFDDTGCGNAKRVICEKELDD
TC14-3 YWVGADNKELGGYNFLWNDGVSLPDSDDLWAPGEPSNPAGKNLCVQLWNKYGLLDDTGCGQKTKNPICEKSLTK
TC14-4 YWVGADNLQNGGYDFLWSDGVALPASSNLWAPKEPSNPQSSQLCVQLWNKYNMLDDTSCGDAKRVICEKEVK-
***** :*.:***:**:*:*:*:*:* ***** . :*****:*.:***:***:*** :.: *****:

B

P-serpin 1

P-serpin 2

P-serpin 3

P-serpin 4

P-serpin 1 AAT CCT GAA CAT --- --- --- --- --- CTG AAA CAT CTG A CTT GTG GTT GTG ACG
P-serpin 2 AAT CCT GAA CAT --- --- --- --- --- CTG AAA CAT CTG A CTT GTG --- --- ACG
P-serpin 3 AAT CCT GAA CAT CTT GAA CAT CTG AAA CAT CTG AAA CAT CTG A CTT GTG GTT GTG ACG
P-serpin 4 AAT CCT GAA CAT CTT GAA CAT --- --- --- CTG AAA CAT CTG A CTT GTG --- --- ACG

Downloaded From: <https://complete.bioone.org/journals/Zoological-Science> on 09 Jun 2025
Terms of Use: <https://complete.bioone.org/terms-of-use>

reproduction can give rise to genetic variation.

Is cell renewal the privileged phenomenon for sexual reproduction?

Unlike immortal germ cells, normal somatic cells have a finite life span. Nonetheless, nuclear transplantation experiments have shown that somatic cell nuclei are capable of resetting the developmental clock and resuming ontogeny after introduced into enucleated oocytes of frogs (King and Briggs, 1956; Gurdon, 1962), sheeps (Wilmut *et al.*, 1997) and mice (Wakayama *et al.*, 1998). This reversible aging of somatic cell nuclei has recently been accounted for in relation to the structure of chromosomal extremities (telomeres). Cell division accompanies the progressive shortening of telomeres (Harley *et al.*, 1990; Hastie *et al.*, 1990). It has been first discovered in *Tetrahymena* that the chromosomal integrity is maintained by telomerase, a ribonucleoprotein that is capable of synthesizing telomeric repeats (Greider and Blackburn, 1985, 1987). Telomerase is a kind of terminal transferase (Morin, 1989) with the activity of reverse transcriptase (Lingner *et al.*, 1997). It is expressed strongly in germline of *Xenopus* (Mantell and Greider, 1994) and human (Wright *et al.*, 1996), in immortal and cancer cells (Kim *et al.*, 1994) and in stem cells such as hematopoietic progenitors (Chiu *et al.*, 1996). On the other hand, it appears to be stringently repressed in normal human somatic cells (Kim *et al.*, 1994). Telomerase-deficient mouse cells showed chromosomal abnormalities including end-to-end fusion (Blasco *et al.*, 1997). When the catalytic subunit of human telomerase is expressed in telomerase-negative normal cells, the telomerase-expressing clones can extend the length of telomeric DNA and exceed their life-span (Bodnar *et al.*, 1998, Vaziri and Benchimol, 1998). These results provide evidence for the telomere hypothesis that telomere shortening is the molecular clock that triggers replicative senescence (Harley, 1991).

Sexual reproduction appears to reset telomere shortening at each generation, as telomerase is expressed in germline. It is quite a mystery if telomerase also operates in asexual reproduction, but our recent EST (expressed sequence tag) data suggested that it might be the case. Out of about 2,000 ESTs in *P. misakiensis* (Kawamura *et al.*, 1998 and our unpublished data), four matched to retrotransposon and retroviral reverse transcriptase with very high similarity (Table 1), although full length of cDNAs remains to be analyzed. As already mentioned, telomerase has the enzyme activity of reverse transcriptase related to retrotransposon and retroviral reverse transcriptase (Cech *et al.*, 1997).

Another mechanism is known to influence cell aging in asexual reproduction of *P. misakiensis*. We would like first to refer to a dedifferentiation system for asexual reproduction (Fig. 5A). In botryllid and polystyelid tunicates, a bud consists of outer and inner epithelia (Kawamura and Watanabe, 1982a, b), of which the inner epithelium evaginates or invaginates to form various organs such as the brain, pharynx and gut (Kawamura and Nakauchi, 1984; Kawamura and Sugino, 1999). It is evident, therefore, that the inner, atrial epithelium

of the bud is developmentally multipotent (Kawamura and Nakauchi, 1984; Fujiwara and Kawamura, 1992). In *P. misakiensis*, the inner, multipotent epithelium expresses several differentiation markers such as pigment granules and alkaline phosphatase in the cytoplasm or on the cell surface (Fujiwara and Kawamura, 1992, Kawamura and Fujiwara, 1994). The cell has a prolonged doubling time of more than 170 hrs (Kawamura *et al.*, 1988). At stages of morphogenesis, on the other hand, cells of organ placodes lose the differentiation markers and enter rapid cell cycling of about 13 hrs, indicating that transdifferentiation should occur (Fujiwara and Kawamura, 1992; Kawamura and Fujiwara, 1994).

In hydras (Bosch and David, 1991) and planarians (Baguña *et al.*, 1989; Shibata *et al.*, 1999), stem cell systems are used for asexual reproduction and regeneration (Fig. 5A). They are very sensitive to ionizing radiation (Fradlin *et al.*, 1978; Baguña *et al.*, 1989). In *P. misakiensis*, on the other hand, irradiated colonies resume bud formation and bud development in temporal accordance with the restoration of mitotic activity even after irradiation at 80 Gy (Kawamura *et al.*, 1995). This remarkable resistance to ionizing radiation affords another evidence for dedifferentiation system in asexual reproduction of *Polyandrocarpa*.

Several factors have been identified to regulate dedifferentiation and cell division of multipotent cells in *P. misakiensis*. One is TC14 lectins. Mature proteins are secreted into the hemolymph during budding and cause mesenchymal cell aggregation to form epithelial tissues (Kawamura *et al.*, 1991). As already mentioned, our published and unpublished studies revealed four kinds of isoforms (TC14-1, TC14-2, TC14-3, TC14-4) (Suzuki *et al.*, 1990; Shimada *et al.*, 1995). They form homo- or heterodimers (Poget *et al.*, 1999). Heterodimers of TC14-2 and TC14-3 isolated from living animals exhibited a remarkable effect on multipotent cells *in vitro* (Matsumoto *et al.*, in preparation). When cells growing at log phase were treated with TC lectins, they became swollen, ceased to divide and expressed the inner epithelium-specific antigen. This activity of the lectins was blocked by D-galactose. By preparing recombinant proteins, Matsumoto *et al.* (in preparation) found that only TC14-3 caused cell cycle arrest and induced cell differentiation. These results have shown that TC14-3 acts as autocrine factor to give replicative senescence to the multipotent epithelium. Interestingly, C-type lectins that TC14s belong to have the three-dimensional structure related to angiogenesis inhibitor endostatin, although almost no sequence similarity has been found (Hohenester *et al.*, 1998).

A few factors that counteract with TC lectins have been known so far in *P. misakiensis*. Both retinoic acid (RA) and proteases are able to induce *in vivo* the secondary bud axis when applied to growing buds (Hara *et al.*, 1992; Kawamura and Watanabe, 1987; Kawamura *et al.*, 1993). Although RA did not directly cause *in vitro* dedifferentiation of the atrial epithelium, the conditioned medium of RA-treated mesenchymal cells caused dedifferentiation of explants of the atrial epithelium (Kawamura *et al.*, in preparation). The conditioned medium showed the increased activity of trypsin-like serine

Table 1. ESTs coding for possible nuclear proteins in *P. misakiensis*.

EST number	Putative identification	Frequencies
Pm13161	ATP-dependent rRNA helicase SPB4	5
Pm12654	brain-specific homeobox/POU domain protein	1
Pm11972	balbiani ring protein 3 precursor	1
Pm12031	chromosome assembly protein XCAP-C	1
Pm01109	cleavage and polyadenylation specificity factor	3
Pm00951	CREB-binding protein	1
Pm01535	DNA-directed RNA polymerase II	2
Pm20820	DNA ligase 1	1
Pm11266	DNA-repair protein	1
Pm13516	EBNA 2 nuclear protein	1
Pm12401	14-3-3 protein	2
Pm01014	GTP-binding nuclear protein RAN/TC4	4
Pm00312	histone H1	3
Pm20502	homeobox protein OM (1D)	1
Pm11435	homeobox protein OTX 1	1
Pm01065	immediate-early protein	1
Pm00672	nitrogen regulatory protein	1
Pm01020	20kd nuclear CAP binding protein	1
Pm00535	nuclear localization sequence binding protein	3
Pm00458	nuclear transition protein 2	1
Pm01574	nucleolar transcription factor 1	1
Pm10208	nucleophosmin	2
Pm10439	nucleoprotein TPR	1
Pm11970	nucleoporin	2
Pm01023	octapeptide-repeat protein T2	1
Pm13356	oocyte zinc finger protein	2
Pm01082	pre-mRNA processing protein PRP39	1
Pm00612	protamine	2
Pm01216	protein HGV2	1
Pm13452	protein kinase CLK2	1
Pm20574	proto-oncogene DBL precursor	2
Pm10306	putative 90.2 KD zinc finger protein	1
Pm00549	putative tumor suppressor luca 15	1
Pm11068	retrovirus-related pol polyprotein from transposon	4
Pm12155	RNA-directed DNA polymerase	1
Pm00554	RNA polymerase b subunit	2
Pm12132	shuttle craft protein	1
Pm12713	60 KD TAT interactive protein	1
Pm10159	small nuclear ribonucleoprotein SM D2	2
Pm10494	sperm histone	1
Pm21031	splicing factor SC35	1
Pm10164	suppressor of zeste protein	1
Pm20704	thyroid receptor interacting protein	1
Pm20842	transposon TX1 hypothetical 149 KD protein	1
Pm00774	ubiquitin carboxyl-terminal hydrolase	1
Pm00863	yeast nuclear protein	1
Pm00864	zinc-binding protein A33	1
Total		72

protease (Kawamura and Fujiwara, 1995). RA is one of endogenous retinoids in *P. misakiensis* (Kawamura *et al.*, 1993). In developing buds, the enzymatic activity of possible retinoic acid synthase is expressed in the epidermis around the dedifferentiating tissue (Kawamura *et al.*, 1993). Both retinoic acid receptor (Hisata *et al.*, 1998) and retinoid X receptor (Kamimura *et al.*, 2000) are induced to express in mesenchymal cells by RA. These results suggested that RA is the primary signal to trigger bud development. RA-treated mesenchymal cells might secrete the secondary signal to

induce dedifferentiation of the atrial epithelium.

Recently, cDNA fragments have been isolated from RA-treated mesenchymal cells by differential display technique (Ohashi *et al.*, 1999). One of them had sequence similarity to trypsin. A full-length cDNA had a serine protease domain at the C-terminus of deduced amino acid sequence (Fig. 5B). At the N-terminus, on the other hand, multiple functional domains were located. The boundary between N- and C-termini had the recognition site cut by plasmin, plasminogen activator and other serine proteases, suggesting that the protein attains to




Fig. 5. Dedifferentiation systems of multipotent cells in *P. misakiensis*. (A) Comparison of a dedifferentiation system with a stem cell system. (B) Primary structure of TRAMP protein involved in dedifferentiation. 1, signal peptide; 2, 7, Scavenger receptor; 3, 4, 6, Low density lipoprotein receptor; 5, kringle domain-like hairpin loop; 8, serine protease. (C) Circuit of differentiation and dedifferentiation of multipotent cells regulated by TC lectin and TRAMP in culture.

functional maturity after enzymatic modification. Thus, this multi-functional protein is called tunicate retinoic acid-inducible modular protease (TRAMP) (Ohashi *et al.*, 1999).

Recombinant protein of the TRAMP catalytic domain has promoted cell growth and cell motility in a dose-dependent manner (Ohashi *et al.*, 1999). Cells became smaller and lost

cytoplasmic granules, suggesting that dedifferentiation might occur. TRAMP showed trypsin-like enzyme activity, while authentic trypsin did not show any mitogenic activity (Ohashi *et al.*, 1999). This result suggested that the sequence responsible for the mitogenic activity should be located outside the conserved region of serine proteases.

Urokinase-type plasminogen activator (uPA) is the well-known vertebrate protease that has the mitogenic activity (Kirchheimer *et al.*, 1988). Human uPA consists of 411 amino acids except for signal peptide. Like TRAMP, it is a modular protease with EGF-like domains and kringle motifs at the N-terminus and with a catalytic domain at the C-terminus (Leslie *et al.*, 1990). Many investigators agree with the notion that the mitogenic activity of uPA is independent of the enzymatic activity (Kirchheimer *et al.*, 1987). Glycosylated EGF and/or kringle domains of uPA at N-terminus have the mitogenic effect on melanoma cells (Koopman *et al.*, 1998) and osteosarcoma cells (Kirchheimer *et al.*, 1987). In these cases, the growth signal of uPA is mediated by high affinity uPA receptors (Rabbani *et al.*, 1992). On the other hand, a N-terminal polypeptide is insufficient for the growth of normal fibroblasts (De Petro *et al.*, 1994), and a full length of uPA is required by the vascular smooth muscle cells (Kanse *et al.*, 1997). Here, uPA seems to use a signal transduction system other than the high affinity receptor. A similar system may work on tunicate cells, as TRAMP recombinant proteins lacking the N-terminal region still have growth-promoting activity.

In the dedifferentiation system of *P. misakiensis*, the multipotent epithelium is regulated to differentiate by counteracting factors such as TC14 lectins and serine protease (Fig. 5C). In the quiescent state or in the presence of TC14s, multipotent cells have the doubling time of more than 170 hr, at least 13-fold longer than that of dedifferentiated cells (Kawamura *et al.*, 1988). We assume that this slow cell cycling of the multipotent epithelium is practically effective in suppressing cell aging.

Conclusion

Unlike empirical hypotheses, we could not find any serious demerits of asexual reproduction for survival in *P. misakiensis*. Asexual reproduction involves the removal of a large number of somatic cells in the process of budding. Clonal animals budded have recognizable genetic variations including gene redundancy and polymorphism. A few polypeptides can regulate differentiation states of the multipotent cells, which would serve to retard their aging. It is interesting to ask how many years hereafter the clones of *P. misakiensis* can continue their asexual lives. We are also interested in the difference in genomic composition between the subclone of Usa and that of Shimoda that have been isolated from each other for just 20 years.

ACKNOWLEDGMENTS

We would like to thank Dr. T. Yubisui for encouraging us during the course of study. Thanks are also due to J. Matsumoto and Y.

Kariya for the permission of use of a part of their unpublished data. This work was supported in part by Grant-in-Aid to KK (11480221) and SF (11780535) from the Ministry of Education, Sports, Culture and Science of Japan.

REFERENCES

- Baguña J, Saló E, Auladell C (1989) Regeneration and pattern formation in planarians III. Evidence that neoblasts are totipotent stem cells and the source of blastema cells. *Development* 107: 77–86
- Barton NH, Charlesworth B (1998) Why sex and recombination? *Science* 281: 1986–1990
- Berrill NJ (1941) The development of the bud in *Botryllus*. *Biol Bull* 80: 169–184
- Blasco MA, Lee HW, Hande MP, Samper E, Landorp PM, DePinho RA, Greider CW (1997) Telomere shortening and tumor formation by mouse cells lacking telomerase RNA. *Cell* 91: 25–34
- Bosch TC, David CN (1991) Decision making in interstitial stem cells of *Hydra*. *In Vivo* 5: 515–520
- Bodnar AG, Kim NW, Effros RB, Chiu CP (1996) Mechanism of telomerase induction during T cell activation. *Exp Cell Res* 228: 58–64
- Bodnar AG, Ouellette M, Frolkis M, Holt SE, Chiu C-P, Morin GB, Harley CB, Shay JW, Lichtsteiner S, Wright WE (1998) Extension of life-span by introduction of telomerase into normal human cells. *Science* 279: 349–352
- Brien P (1968) Blastogenesis and morphogenesis. In "Advances in Morphogenesis Vol 7" Ed by M Abercrombie, J Brachet and TJ King, Academic Press, New York, pp 151–203
- Caballero A, Keightley PD (1998) Influences on genome-wide deleterious mutation rates in inbred populations of *Drosophila* and mice. *Genetica* 102–103: 229–239
- Cech TR, Nakamura TM, Lingner J (1997) Telomerase is a true reverse transcriptase. A review. *Biochemistry* 62: 1202–1205
- Chiu CP, Dragowska W, Kim NW, Vaziri H, Yui J, Thomas TE, Harley CB, Lansdorp PM (1996) Differential expression of telomerase activity in hematopoietic progenitors from adult human bone marrow. *Stem Cells* 14: 239–248
- Dawson KJ (1998) Evolutionarily stable mutation rate. *J Theor Biol* 194: 143–157
- Deng HW, Lynch M (1997) Inbreeding depression and inferred deleterious mutation parameters in *Daphnia*. *Genetics* 147: 147–155
- De Petro G, Copeta A, Barlati S (1994) Urokinase-type and tissue-type plasminogen activators as growth factors of human fibroblasts. *Exp Cell Res* 213: 286–294
- Drake JW (1991) A constant rate of spontaneous mutation in DNA-based microbes. *Proc Natl Acad Sci USA* 88: 7160–7164
- Drickamer K (1993) Ca²⁺-dependent carbohydrate-recognition domains in animal proteins. *Curr Opin Struct Biol* 3: 393–400
- Fisher RA (1930) The genetical theory of natural selection. Oxford Univ Press, Oxford
- Fradlin M, Kakis H, Campbell RD (1978) Effect of gamma irradiation of hydra: elimination of interstitial cells from viable hydra. *Radiat Res* 76: 187–197
- Fujiwara S, Kawamura K (1992) Ascidian budding as a transdifferentiation-like system: multipotent epithelium is not undifferentiated. *Develop Growth Differ* 34: 463–472
- Gilbert JJ (1976) Selective cannibalism in the rotifer *Asplanchna sieboldi*: contact recognition of morphotype and clone. *Proc Natl Acad Sci USA* 73: 3233–3237
- Greene LJ, Bartelt DC (1969) The structure of the bovine pancreatic secretory trypsin inhibitor—Kazal' inhibitor. II The order of the tryptic peptides. *J Biol Chem* 244: 2646–2657
- Greider CW, Blackburn EH (1985) Identification of a specific telom-

- ere terminal transferase activity in *Tetrahymena* extracts. *Cell* 43: 405–413
- Greider CW, Blackburn EH (1987) The telomere terminal transferase of *Tetrahymena* is a ribonucleoprotein enzyme with two kinds of primer specificity. *Cell* 51: 887–898
- Gurdon JB (1962) The developmental capacity of nuclei taken from intestinal epithelial cells of feeding tadpoles. *J Embryol Exp Morphol* 10: 622–640
- Hara K, Fujiwara S, Kawamura K (1992) Retinoic acid can induce a secondary axis in developing buds of a colonial ascidian, *Polyandrocarpa misakiensis*. *Develop Growth Differ* 34: 437–445
- Harley CB (1991) Telomere loss: mitotic clock or genetic time bomb? *Mutat Res* 256: 271–282
- Harley CB, Futcher AB, Greider CW (1990) Telomeres shorten during aging of human fibroblasts. *Nature* 345: 458–460
- Hastie ND, Dempster M, Dunlop MG, Thompson AM, Green, DK, Allshire RC (1990) Telomere reduction in human colorectal carcinoma and with aging. *Nature* 346: 866–868
- Hisata K, Fujiwara S, Tsuchida Y, Ohashi M, Kawamura K (1998) Expression and function of a retinoic acid receptor in budding ascidians. *Dev Genes Evol* 208: 537–546
- Hohenester E, Sasaki T, Olsen BR, Timpl R (1998) Crystal structure of the angiogenesis inhibitor endostatin at 1.5 angstrom resolution. *EMBO J* 17: 1656–1664
- Holland PWH, Garcia-Fernández J, Williams NA, Sidow A (1994) Gene duplication and the origins of vertebrate development. *Development (Suppl)* 125–133
- Hughes AL (1994) The evolution of functionally novel proteins after gene duplication. *Proc Roy Soc Lond Ser B* 256: 119–124
- Ishii K, Matsuda H, Iwasa Y, Sasaki A (1989) Evolutionarily stable mutation rate in a periodically changing environment. *Genetics* 121: 163–174
- Ishii T, Saito Y, Taneda Y (1993) Histological differences between the color patterns of two strains of the compound ascidian *Polyandrocarpa misakiensis*. *Zool Sci* 10: 977–982
- Izzard CS (1973) Development of polarity and bilateral asymmetry in the palaeal bud of *Botryllus schlosseri* (Pallas). *J Morphol* 139: 1–26
- Johnson T (1999) Beneficial mutations, hitchhiking and the evolution of mutation rates in sexual populations. *Genetics* 151: 1621–1631
- Kaback DB (1995) Yeast genome structure. In “The Yeast Vol 6” Eds AE Wheals, Rose AH, JS Harrison, Academic Press, London, pp 179–222
- Kabay ME, Gilbert JJ (1977) Polymorphism and reproductive mode in the rotifer, *Asplanchna sieboldi*: relationship between meiotic oogenesis and shape of body-wall outgrowths. *J Exp Zool* 201: 21–27
- Kamimura M, Fujiwara S, Kawamura K, Yubisui T (2000) Functional retinoid receptors in budding ascidians. *Develop Growth Differ* (in press)
- Kanse SM, Benzakour O, Kanthou C, Kost C, Lijnen R, Preissner KT (1997) Induction of vascular SMC proliferation by urokinase indicates a novel mechanism of action in vasoproliferative disorders. *Arter Throm Vas Biol* 17: 2848–2854
- Kawamura K, Fujiwara S (1994) Transdifferentiation of pigmented multipotent epithelium during morphallactic development of budding tunicates. *Int J Dev Biol* 38: 369–377
- Kawamura K, Fujiwara S (1995) Cellular and molecular characterization of transdifferentiation in the process of morphallaxis of budding tunicates. *Semin Cell Biol* 6: 117–126
- Kawamura K, Fujiwara S, Sugino YM (1991) Budding-specific lectin induced in epithelial cells is an extracellular matrix component for stem cell aggregation in tunicates. *Development* 113: 995–1005
- Kawamura K, Hashimoto, K, Nakauchi M (1995) Development of irradiated tunicate buds: Is cell division cycle required for morphallaxis? *Develop Growth Differ* 37: 487–496
- Kawamura K, Hara K, Fujiwara S (1993) Developmental role of endogenous retinoids in the determination of morphallactic field in budding tunicates. *Development* 117: 835–845
- Kawamura K, Hayata D, Fujiwara S, Yubisui T (1998) Serine protease inhibitors expressed in the process of budding of tunicates as revealed by EST analysis. *J Biochem* 124: 1004–1012
- Kawamura K, Nakauchi M (1984) Changes of the atrial epithelium during the development of anteroposterior organization in palaeal buds of the polystyelid ascidian, *Polyandrocarpa misakiensis*. *Mem Fac Sci Kochi Univ Ser D* 5: 15–35
- Kawamura K, Nakauchi M (1986) Development of spatial organization in palaeal buds of the compound ascidian, *Symplesma reptans*. *Biol Bull* 171: 520–537
- Kawamura K, Ohkawa S, Michibata H, Nakauchi M (1988) Autoradiographic studies on cell population kinetics in epithelial and haematopoietic stem cell lines in the process of palaeal budding of ascidians. *Mem Fac Sci Kochi Univ Ser D* 9: 1–12
- Kawamura K, Sugino YM (1999) Cell adhesion in the process of asexual reproduction of tunicates. *Micro Res Tech* 40: 269–278
- Kawamura K, Watanabe H (1981) Studies of Japanese compound styelid ascidians. III. A new, possibly asexual *Polyandrocarpa* from Shimoda Bay. *Publ Seto Mar Biol Lab* 26: 425–436
- Kawamura K, Watanabe H (1982a) Pattern development in palaeal buds of botryllid ascidians: Relation between parent and its buds in their body axes. *J Morphol* 173: 293–304
- Kawamura K, Watanabe H (1982b) Localized morphogenetic activity in the palaeal bud of a polystyelid ascidian, *Polyandrocarpa misakiensis*. I. A histological study. *Mem Fac Sci Kochi Univ Ser D* 3: 55–69
- Kawamura K, Watanabe H (1984) The mechanism of anteroposterior cell determination in ascidian palaeal buds: A gap of positional values triggers posterior formation. *Roux's Arch Dev Biol* 193: 24–35
- Kawamura K, Watanabe H (1987) Possible role of the epidermis and mesenchyme in the determination of morphogenesis domain during palaeal budding in ascidians. *Mem Fac Sci Kochi Univ Ser D* 8: 15–27
- Kazal LA, Spicer DS, Brahinsky RA (1948) Isolation of a crystalline trypsin inhibitor-anticoagulant protein from the pancreas. *J Am Chem* 70: 304–340
- Keightley PD (1994) The distribution of mutation effects on viability in *Drosophila*. *Genetics* 138: 1315–1322
- Keightley PD, Caballero A, Garcia-Dorado A (1998) Population genetics: Surviving under mutation pressure. *Curr Biol* 8: 235–237
- Kibota TT, Lynch M (1996) Estimate of the genomic mutation rate deleterious to overall fitness in *E. coli*. *Nature* 381: 694–696
- Kim NW, Piatyszek MA, Prowse KR, Harley CB, West MD, Ho PL, Coviello GM, Wright WE, Weinrich SL, Shay JW (1994) Specific association of human telomerase activity with immortal cells and cancer. *Science* 266: 2011–2015
- Kimura M (1967) On the evolutionary adjustment of spontaneous mutation rates *Genet Res* 9: 23–34
- King TJ, Briggs R (1956) Serial transplantation of embryonic nuclei. *Cold Spring Harbor Symp Quant Biol* 21: 271–289
- Kirchheimer JC, Wojta J, Christ G, Binder BR (1987) Proliferation of a human epidermal tumor cell line stimulated by urokinase. *FASEB J* 1: 125–128
- Kirchheimer JC, Wojta J, Christ G, Binder BR (1988) Mitogenic effect of urokinase on malignant and unaffected adjacent human renal cells. *Carcinogenesis* 9: 2121–2123.
- Kondrashov AS, Crow JF (1993) A molecular approach to estimating the human deleterious mutation rate. *Hum Mutat* 2: 229–234
- Koopman JL, Slomp J, de Bart AC, Quax PH, Verheijen JH (1998) Mitogenic effects of urokinase on melanoma cells are independent of high affinity binding to the urokinase receptor. *J Biol Chem*

- 273: 33267–33272
- Leigh EG (1973) The evolution of mutation rates. *Genetics* 73: 1–18
- Leslie ND, Kessler CA, Bell SM, Degen JL (1990) The chicken urokinase-type plasminogen activator gene. *J Biol Chem* 265: 1339–1344
- Lingner J, Hughes TR, Shevchenko A, Mann M, Lundblad V, Cech TR (1997) Reverse transcriptase motifs in the catalytic subunit of telomerase. *Science* 276: 561–567
- Lushai G, De Barro PJ, David O, Sherratt TN, Maclean N (1998) Genetic variation within a parthenogenetic lineage. *Insect Mol Biol* 7: 337–344
- Mandrioli M, Manicardi GC, Bizzaro D, Bianchi U (1999) NOR heteromorphism within a parthenogenetic lineage of the aphid *Megoura viciae*. *Chromosome Res* 7: 157–162
- Mantell LL, Greider CW (1994) Telomerase activity in germline and embryonic cells of *Xenopus*. *EMBO J* 13: 3211–3217
- Maynard Smith J (1978) *The evolution of sex*. Cambridge Univ Press, Cambridge
- Morgan TH (1901) *Regeneration*. MacMillan Company, London
- Morin GB (1989) The human telomere terminal transferase enzyme is a ribonucleoprotein that synthesizes TTAGGG repeats. *Cell* 59: 521–529
- Mukai H, Sugimoto K, Taneda Y (1978) Comparative studies on the circulatory system of the compound ascidians, *Botryllus*, *Botrylloides* and *Symplegma*. *J Morphol* 157: 49–78
- Mukai T, Chigusa SI, Mettler LE, Grow JF (1972) Mutation rate and dominance of genes affecting viability in *Drosophila melanogaster*. *Genetics* 72: 333–355
- Nakauchi M (1982) Asexual development of ascidians: its biological significance, diversity, and morphogenesis. *Amer Zool* 22: 753–763
- Newberry AT (1965) Vascular structures associated with budding in the polystyelid ascidian *Metandrocarpa taylori*. *Ann Soc Roy Zool Berg* 95: 57–74
- Normark BB (2000) Molecular systematics and evolution of the aphid family Lachnidae. *Mol Phylogenet Evo* 14: 1331–140
- Ohashi M, Kawamura K, Fujii N, Yubisui T, Fujiwara S (1999) A retinoic acid-inducible modular protease in budding ascidians. *Develop Biol* 214: 38–45
- Ohno S (1970) *Evolution by gene duplication*. Springer-Verlag, Heidelberg
- Poget SF, Legge GB, Proctor MR, Jonathan P, Butler G, Bycroft M, Williams RL (1999) The structure of tunicate C-type lectin from *Polyandrocarpa misakiensis* complexed with D-galactose. *J Mol Biol* 290: 867–879
- Rabbani SA, Mazar AP, Bernier SM, Haq M, Bolivar I, Henkin J, Goltzman D (1992) Structural requirements for the growth factor activity of the amino-terminal domain of urokinase. *J Biol Chem* 267: 14151–14156
- Shibata N, Umesono Y, Orii H, Sakurai T, Watanabe K, Agata K (1999) Expression of vasa (vas)-related genes in germline cells and totipotent somatic stem cells of planarians. *Develop Biol* 206: 73–87
- Shimada M, Fujiwara S, Kawamura K (1995) Expression of genes for two C-type lectins during budding of the ascidian *Polyandrocarpa misakiensis*. *Roux's Arch Dev Biol* 204: 406–411
- Shufra KA, Peters CD, Webster JA (1997) Generation of clonal diversity by sexual reproduction in the greenbug, *Schizaphis graminum*. *Insect Mol Biol* 6: 203–209
- Sniegowski PD (1997) Evolution: Setting the mutation rate. *Curr Biol* 7: 487–488
- Sniegowski PD, Gerrish PJ, Lenski RE (1997) Evolution of high mutation rates in experimental populations of *E. coli*. *Nature* 387: 703–705
- Suzuki T, Takagi T, Furukohri T, Kawamura K, Nakauchi M (1990) A calcium-dependent galactose-binding lectin from the tunicate *Polyandrocarpa misakiensis*. *J Biol Chem* 265: 1274–1281
- Taddei F, Radman M, Maynard-Smith J, Toupance B, Gouyon PH, Godelle B (1997) Role of mutator alleles in adaptive evolution. *Nature* 387: 700–702
- Wakayama T, Perry AC, Zuccotti M, Johnson KR, Yanagimachi R (1999) Full-term development of mice from enucleated oocytes injected with cumulus cell nuclei. *Nature* 394: 369–374
- Watanabe H, Tokioka T (1972) Two new species and one possibly new race of social styelids from Sagami Bay, with remarks on their life history, especially the mode of budding. *Publ Seto Mar Biol Lab* 19: 327–345
- Wilmot I, Schnieke AE, McWhir J, Kind AJ, Campbell KH (1997) Viable offspring derived from fetal and adult mammalian cells. *Nature* 385: 810–813
- Wilson AC, Sunnucks P, Hales DF (1999) Microevolution, low clonal diversity and genetic affinities of parthenogenetic sitobion aphids in New Zealand. *Mol Ecol* 8: 1655–1666
- Wolfe KH, Shields DC (1997) Molecular evidence for an ancient duplication of the entire yeast genome. *Nature* 387: 708–713
- Wright WE, Piatyszek MA, Rainey WE, Byrd W, Shay JW (1996) Telomerase activity in human germline and embryonic tissues and cells. *Dev Genet* 18: 173–179
- Wuethrich B (1998) Why sex? Putting theory to the test. *Science* 281: 1980–1982
- Wurdak ES, Gilbert JJ (1976) Polymorphism in the rotifer *Asplanchna sieboldi*. Fine structure of saccate, cruciform and campanulate females. *Cell Tissue Res* 169: 435–448

(Received February 21, 2000 / Invited Review)