

Region-Specific Loss of Two-Headed Ciliary Dyneins in Ascidian Endostyle

Authors: Konno, Alu, and Inaba, Kazuo

Source: Zoological Science, 37(6): 512-518

Published By: Zoological Society of Japan

URL: https://doi.org/10.2108/zs200095

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Region-specific Loss of Two-headed Ciliary Dyneins in Ascidian Endostyle

Alu Konno^{1,2} and Kazuo Inaba^{2*}

¹Department of Virology and Parasitology, Hamamatsu University School of Medicine, 1-20-1 Handayama, Higashi-ku, Hamamatsu, Shizuoka 431-3192, Japan ²Shimoda Marine Research Center, University of Tsukuba, Shimoda 5-10-1, Shizuoka 415-0025, Japan

A mucous secreting organ in ascidians, the endostyle, consists of several epithelial zones with different ciliary length, density, and beating direction. Here we found by transmission electron microscopy that long cilia in endostyle zone 1 showed 9+2 axonemal structures but completely lacked the outer arm dynein. In contrast, cilia in other zones bore both outer and inner dynein arms. Western blotting and immunofluorescence microscopy further revealed that zone 1 appeared to lack not only outer arm dynein but also two-headed inner arm dynein. These results suggest a mechanism for a region-specific gene suppression that causes the limited loss of two-headed axonemal dyneins in the endostyle epithelium. The loss of these dyneins in zone 1 is considered to contribute to the generation of undulating ciliary movement that is essential for a unique circuit of mucus flow in the endostyle.

Key words: cilia, flagella, dynein, ciliogenesis, Ciona, endostyle

INTRODUCTION

Cilia and flagella are microtubule-based organelles with conserved internal 9+2 cytoskeletal structures, called the axonemes (Gibbons, 1981; Inaba, 2003; Konno et al., 2012). Motile cilia and flagella drive the generation of extracellular fluid flow, such as that in cell locomotion, epithelial mucociliary clearance, and vertebrate left-right symmetry breaking event. The motility is generated by two projections on the nine outer doublet microtubules, the outer and inner arm dyneins, which are responsible for the sliding of adjacent doublet microtubules in the axonemes. The outer and inner arm dyneins mainly play roles in the increase of beat frequency and in the formation and propagation of basic ciliary or flagellar bend, respectively. Loss of dyneins causes human diseases affecting multiple organs, called primary ciliary dyskinesia (PCD), Kartagener syndrome or immotile ciliary syndrome (Afzelius, 1976; Satir and Christensen, 2008).

Metazoan cilia and flagella show structural diversity. The 9+2 motile cilia are considered to be ancient forms of immotile cilia, which might have lost the structures for motility during ciliary diversifications (Mitchell, 2007; Inaba, 2015). For example, sea urchin swimming blastula possess long cilia on the animal pole, called the apical tuft. These cilia show less motility and appear specialized for mechanical reception with apparently normal 9+2 structures (Jin et al., 2013). In ascidian larvae, the structures of cilia show more diversity than those of sea urchins. They have not only completely motile cilia in the neural tube but also many diverse forms of immotile cilia in the epithelia and sensory vesicle (Konno et al., 2010). Completely immotile cilia lack the dynein arms or the central two singlet microtubules as well in some cases. These cilia are specialized for cellular antennas that sense extracellular stimuli and convert them into an intracellular signaling pathway, as seen in primary cilia or sensory cilia (Marshall and Nonaka, 2006).

The endostyle of adult ascidians is considered to be an ancient organ homologous to the vertebrate thyroid and plays a critical role in the transport of mucus to the branchial basket (Fujita and Nanba, 1971; Thorndyke, 1978; Holley, 1986; Burighel and Cloney, 1997; Sasaki et al., 2003). The ciliary types and their distribution in the endostyle epithelium are well organized in order to drive fluid flow for efficient transport of secreted mucus to the branchial basket (Fujita and Nanba, 1971; Pennachetti, 1984; Holley, 1986; Burighel and Cloney, 1997; Petersen, 2007). Here, we show the structural diversity of ciliary axonemes in endostyle of the ascidian *Ciona intestinalis*. We found that the zone 1 of endostyle epithelia with extremely long cilia lacked both outer and inner arm two-headed dyneins. To our knowledge, this is the first work that reports such cilia in nature.

MATERIALS AND METHODS

Preparation and observation of endostyle

Adults of *Ciona intestinalis* (recently renamed as *Ciona intestinalis* type A or *Ciona robusta*) were collected at Onagawa Bay near the Education and Research Center of Marine Bio-Resources, Tohoku University, and in part supplied by the Department of Zoology, Kyoto University and by Misaki Marine Biological Station, University of Tokyo through National Bio-Resource Project (NBRP), Japan. A closely related species, *Ciona savignyi*, was collected in Otsuchi Bay and used for supplementary use for observation of the endostyle structure. Endostyles were cut out from the

^{*} Corresponding author. E-mail: kinaba@shimoda.tsukuba.ac.jp doi:10.2108/zs200095

branchial basket and transferred to a petri dish filled with filtered seawater. They were further trimmed and divided into small pieces for observation under a differential interference contrast microscope (Olympus BX53). Images and movies were recorded using a DF74 (Olympus) or D3 high-speed (DITECT) camera with a 10× or 100× objective for zone 1 or 5 cilia, respectively, and analyzed using ImageJ.

Transmission electron microscopy (TEM)

The endostyles were dissected out and fixed with the fixative (2.5% glutaraldehyde, 0.45 M sucrose, 0.1 M sodium cacodylate, pH 7.4) for 2 h at 4° C. The subsequent steps were conducted according to the method described previously (Konno et al., 2010).

Analysis of dynein arm structures

To obtain the average images of dyneins on each doublet microtubule, TEM images of cross-sectioned cilia were corrected and transformed by rotation according to the method described by Gadelha et al. (2006).

Preparation of endostyle cilia

Excised endostyles were rinsed several times with filtered sea water and transferred to an ice-cold hypertonic solution containing 1.5 M NaCl, 1 mM MgCl₂, 0.5 mM EGTA, 20 mM Tris-HCl, pH 8.0 (Konno et al., 2015). They were kept for 2 min under gentle agitation for deciliation. After the tissues were removed, the suspension was briefly centrifuged at 500 g for 3 min to precipitate the tissue debris. The supernatant containing endostyle cilia was used for immunostaining. For biochemical analysis, the supernatant was further centrifuged at 8000 g for 10 min at 4° C to recover cilia as a pellet. The pellet was rinsed twice with filtered sea water and used for SDS-PAGE and subsequent immunoblotting.

Preparation of sperm flagella

Sperm were collected from the sperm duct of an adult by dissection. The flagella were isolated by homogenization and centrifugation, according to the method described previously (Padma et al., 2003).

Antibodies

Primary antibodies used in this study were those against outer arm IC2 and IC3 (Hozumi et al., 2006; Padma et al., 2001); outer arm LC1 (LRR) and LC2 (Tctex2) (Hozumi et al., 2006; Padma et al., 2001); PKA-R (Hozumi et al., 2008); calaxin (Mizuno et al., 2009); Ap58 (Ogawa and Inaba, 2006); inner arm IC116 (Hozumi et al., 2008); RSP3 (Padma et al., 2003); HSP40 (Satouh et al., 2005); LRR37 (Padma et al., 2001); MORN40 (Satouh et al., 2005); PF16 (Ci-Spag6; Satouh and Inaba, 2009); inner arm dynein IC105 (ortholog of Chlamydomonas f/l1 dynein IC97; Wirschell et al., 2009). All antibodies were prepared according to the method of Padma et al. (2003). Anti-acetylated α-tubulin (Lys40) rabbit monoclonal antibody (RM318) was purchased from Invitrogen Corp. Anticentrin mouse monoclonal antibody (clone 20H5) was purchased from Sigma-Aldrich Co., Ltd. For secondary antibody, goat antimouse IgG (H+L)-HRP (Invitrogen Corp.) was used for western blotting. Alexa 488-conjugated anti-rabbit IgG antibody and Alexa 546-anti-mouse IgG antibody were purchased from Molecular Probes and used as secondary antibodies for immunofluorescent labeling (1:1000 dilution).

Electrophoresis and immunoblotting

SDS-PAGE was performed according to the method described by Laemmli (1970) with 10% polyacrylamide as the separating gel. Proteins were transferred onto a polyvinylidene fluoride membrane and subjected to western blotting according to the method previously described (Padma et al., 2003). The blots were developed using the ECL[®] (enhanced chemiluminescence)-plus detection system (Amersham Biosciences Inc.).

Immunofluorescence microscopy

Cilia detached from endostyle epithelia were attached on a poly-L-lysine-coated glass slide and fixed with cold methanol at -30° C for 10 min. After air-drying, the samples were rehydrated by PBS and incubated in 10% goat serum in PBS in a moist chamber for 1 hr. Immunostaining was performed using a previously described method (Hozumi et al., 2008) and observed using an Olympus BX53 fluorescent microscope with a $\times 20$ objective.

RESULTS

The endostyle is a gutter-shaped organ running along the whole ventral side of a branchial sac and is composed of nine parallelly running regions or "zones". All regions but zones 7 and 9 are ciliated. The ciliary length, density, or beating directions are different among zones (Fig. 1A). Among them, zone 1 runs in the midline and forms a tuft of quite long cilia with the lengths reaching over 500 μ m (448 \pm $68 \,\mu m$, n = 18). Typical cilia of zone 1 in the tuft showed very slow undulating or irregular movement with small amplitudes (less than 10 μ m), with beat frequency of 0.86 \pm 0.38 Hz (n = 12) (Fig. 1B; see Supplementary Movie S1). Atypical movements with larger amplitudes and shorter wavelengths were occasionally observed in the ciliary tuft when the endostyle was flattened and zone 1 cilia were spread laterally between the glass slide and the coverslip (see Supplementary Movie S2).

On the other hand, the cilia in other zones showed typical ciliary movements with high beat frequency. For example, the middle region of an endostyle, zone 5, possessed short cilia that actively beat (see Supplementary Movie S3). The ciliary length and the beat frequency of zone 5 cilia were estimated as $8.7 \pm 0.8 \ \mu m \ (n = 16)$ and $6.1 \pm 1.5 \ Hz \ (n = 12)$, respectively. Although several TEM observations have been done in the endostyle (Burighel and Cloney, 1997), the ciliary ultrastructure has not been reported in detail.

To determine the ciliary structures in different zones of the endostyle in more detail, we examined the structures of zone 1 cilia and compared them with those in zone 5, which showed typical ciliary beating and was the best region to easily prepare TEM samples for analysis. Cilia from both zones were multiciliated (Fig. 1C and 1D). Close observations of the axonemal structures revealed that cilia of zone 1 had 9+2 structures without the outer dynein arms (Fig. 1E), whereas those in zone 5 showed normal motile 9+2 structures with both outer and inner dynein arms (Fig. 1F).

To confirm biochemically if the cilia of zone 1 lacked the outer arm dyneins and other axonemal components, we carried out western blotting of whole endostyle cilia with antibodies against the subunits of outer arm dynein, as well as those against other axonemal components (Fig. 2). We compared the blot patterns of endostyle cilia with those of sperm flagella, which have complete sets of outer and inner arm dyneins (Padma et al., 2001, 2003; Hozumi et al., 2006, 2008; Satouh et al., 2005; Satouh and Inaba, 2009). We expected that the immunoblots reflected the components of zone 1 cilia because their long cilia occupied a substantial portion of whole endostyle cilia. In fact, the ratio of the number of zone 1 long cilia to the sum of short cilia in other zones would be 8:100, based on the schematic distribution of cilia in an endostyle (Burighel and Cloney, 1997). If we suppose



Fig. 1. Cilia in the zone 1 of an endostyle lacking outer arm dyneins. **(A)** Left, a schematic drawing of adult *C. intestinalis*. The body axes are shown; ventral (V), dorsal (D), anterior (A), posterior (P). Traditionally a branchial side is treated as ventral. Endostyle runs along the entire ventral midline of the branchial sac. Bc, branchial sac; En, endostyle. Right, a schematic drawing of a cross section of the endostyle. It is composed of nine distinct zones. **(B)** Movements of typical cilia in zone 1. Five sequential images with 50 msec intervals are overwritten. Scale bar, 50 μ m. **(C)** An electron micrograph of zone 1. Lower magnification. Scale bar, 500 nm. **(D)** An electron micrograph of zone 5. Lower magnification. Scale bar, 500 nm. **(F)** A cross section of long cilia found in zone 1, showing 9+2 structure without outer dynein arms. Scale bar, 100 nm. **(F)** A cross section of short cilia found in zone 5, showing 9+2 structure with both dynein arms. Scale bar, 100 nm.

that the ciliary length in zone 1 is 500 μ m and that in other zones is 10 μ m, the ratio of ciliary amount would be 4000:1000. This demonstrates that zone 1 cilia contribute to the total amount of ciliary protein in the endostyle significantly enough to detect the lack of an axonemal protein by western blotting.

Regarding the components of outer arm dynein, two intermediate chains (IC2 and IC3) and two light chains (a leucine-rich repeat light chain, LC1, and a Tctex2-related light chain, LC2; Inaba, 2011) showed significant decreases in endostyle cilia, compared to those in sperm flagella (Fig. 2). Interestingly, LC1 exceptionally showed signal with similar levels between endostyle cilia and sperm flagella. In addition, three proteins that are known to be associated with



Fig. 2. Western blot analysis of axonemal components in sperm flagella and endostyle cilia. Coomassie Brilliant Blue R250 stained tubulin bands are shown as a loading control. The proteins were detected in endostyle cilia (left lanes) and sperm flagella (right lanes) using primary antibodies as follows: outer arm dynein intermediate chains IC2 and IC3; outer arm dynein light chains LC1 and LC2; the regulatory subunit of cAMP-dependent protein kinase (PKA-R); outer arm dynein associated calcium sensor calaxin; outer arm dynein associated protein Ap58; intermediate chains of two-headed inner arm dynein IC105 and IC116; radial spoke protein RSP3, heat shock protein HSP40, leucine-rich repeat protein LRR37; MORN repeat protein MORN40; central apparatus protein PF16 (Ci-Spag6). En, endostyle cilia; Sp, sperm flagella.

the outer arm dynein were also tested: i.e., the regulatory subunit of type II cAMP-dependent protein kinase (PKA-R; Nomura et al., 1999; Itoh et al., 2000; Hozumi et al., 2008), calaxin (Mizuno et al., 2009, 2012), and Ap58 (Ogawa and Inaba, 2006). All of these proteins showed no or only weak signals in endostyle cilia (Fig. 2).

Next, we examined two components of a two-headed inner arm dynein (f/l1 dynein), IC105 (ortholog of Chlamydomonas f/l1 dynein IC97; Wirschell et al., 2009) and IC116 (ortholog of Chlamydomonas IC140; Hozumi et al., 2008); four components of radial spokes, RSP3 (Padma et al., 2003), HSP40 (Satouh et al., 2005), LRR37 (Padma et al., 2003), and MORN40 (Satouh et al., 2005), and a central apparatus component, PF16 (Satouh and Inaba, 2009) (Fig. 2). All four components of radial spokes showed similar signals in western blotting between endostyle cilia and sperm flagella. However, the signal of PF16 was significantly decreased in endostyle cilia. We particularly focused on the fact that two components of f/l1 dynein, IC105 and IC116, were included in the endostyle cilia at significantly lower levels. This suggests that f/l1 dynein, as well as the outer arm dynein, is missing in endostyle zone 1.

To examine the structure of the inner arm dyneins in zone 1, we further analyzed TEM images of dynein by averaging (Gardner et al., 1994; Gadelha et al., 2006). The averaged images clearly showed the lack of outer arm dynein in zone 1 but also the difference in the structure of inner arm dyneins (Fig. 3A and 3B). In zone 5 cilia, the inner arm dynein was observed as two projections toward the adjacent B-tubule Fig. 3C), as reported previously (Kamiya et al., 1992). The axoneme of zone 1 cilia lacked the outer part of the projections (Fig. 3D; dotted circle in Fig. 3C), which cor-



Fig. 3. Nine-fold averaged images of axonemal structures in zones 1 and 5. The images were obtained by rotational averaging of two high-contrast, cross-sectioned TEM images of zone 1 and zone 5 cilia. **(A)** An averaged image of zone 5 cilia. **(B)** An averaged image of zone 1 cilia. **(C)** Magnified four images of a doublet microtubule in zone 5. **(D)** Magnified four images of a doublet microtubule in zone 1. The zone 1 cilia lack the outer dynein arm (arrow) and the outer part of inner dynein arm (arrowhead). A dashed circle in **(C)** (left panel) indicates the outer part of inner dynein arm in zone 5 cilia that is missing in zone 1. The asterisk in **(D)** (left panel) indicates the electron-dense structure in the inside lumen of A-tubule.

responds to the two-headed f/l1 dynein (Kamiya et al., 1992). This result supports the data of western blotting that suggest the absence of f/l1 in zone 1 cilia (Fig. 2). In addition, an electron dense structure was seen inside the lumen of the A-tubule in zone 1 cilia (white asterisk in Fig. 3D). Immuno-fluorescence microscopy further showed that an intermediate chain of f/l1 inner arm dynein, IC116, was only detected in short cilia but not in long cilia (Fig. 4A). Outer arm dynein IC2 was not also detected in long cilia, supporting the idea that both outer and inner arm two-headed dyneins are missing in the zone 1 long cilia (Fig. 4B). In contrast, centrin, a component of some single-headed inner arm dyneins, was detected in both long and short cilia from the endostyle (Fig. 4C).

DISCUSSION

The most intriguing finding in this study is that zone 1 cilia lack both outer and inner arm two-headed dyneins. This unique property in the axonemal structure is thought to be linked to the production of undulating motility by these long cilia. The outer arm dynein is known to contribute to the increase of beat frequency (Gibbons and Gibbons, 1979; Brokaw and Kamiya, 1987), whereas the f/l1 inner arm dynein is essential for beating in the axonemes lacking outer arm dynein or single-headed inner arm dyneins (Kamiya and Yagi, 2014). In *Chlamydomonas*, double mutants lacking two of three types of dyneins, i.e., outer arm, inner arm, and single-headed inner arm dyneins, are completely non-motile



Fig. 4. Immunofluorescence microscopy of isolated cilia. **(A)** Double staining by antibodies against acetylated α -tubulin and twoheaded inner arm dynein IC116. **(B)** Double staining by antibodies against acetylated α -tubulin and outer arm dynein IC2. **(C)** Double staining by antibodies against acetylated α -tubulin and centrin, a component of single-headed inner arm dyneins. White and gray arrowheads show the long cilia stained by acetylated α -tubulin antibody and the short cilia stained by antibodies against dynein components, respectively. Scale bar, 50 µm.

under physiological conditions (Kamiya, 1995; Kamiya and Yagi, 2014).

However, zone 1 cilia in *Ciona* endostyle show undulating motility, though they lack both outer and inner arm twoheaded dyneins and appear to possess only single-headed inner arm dyneins. It is known that apparent non-motile double mutants in *Chlamydomonas* become motile under non-physiological conditions, such as high concentrations of salt or organic compounds (Yagi and Kamiya, 2000). The motility of zone 1 cilia would therefore be induced by some non-physiological factors, most likely the mechanical constraints by tight association in the tuft of zone 1 cilia. This might be supported by the fact that a part of cilia in the tuft change their motility when they are mechanically spread (Supplementary Movie S2).

Endostyle secretes the mucus that is essential for filter-

ing the sea water and trapping particles such as plankton. The mucus is transported from the endostyle through branchial epithelium to the dorsal lamina (Burighel and Cloney, 1997). The ciliary activity of endostyle epithelia is considered to be important not only for sending mucus out to the branchial basket but also for the formation of the flow circuit for mucus (Holley, 1986). The zone 1 epithelium spreads out long cilia along the midline. However, the cilia show less motility with occasional undulating movement and appear not to exert active fluid flow (Supplementary Movie S1). This poor motility of zone 1 cilia appears to make sense, because the cilia appear to create a fence that separates the mucus flow pathway in two ways along either side of the branchial basket (see Fig. 1B). The occasional undulation of the cilia may facilitate the mucus transport without disturbing the flow along the long cilia.

The radial spokes and central apparatus are involved in the activation of a subset of dynein to generate planar ciliary waves (Porter and Sale, 2000). Lack of these structures results in helical movements, as seen in the flagella of eel sperm (Gibbons et al., 1985) and Asian horseshoe crab (Ishijima et al., 1988). It is interesting that zone 1 cilia still keep the radial spokes and central apparatus, despite the fact that they only show occasional undulating movement. Further waveform analysis of zone 1 cilia might give a clue to know how single-headed inner arm dyneins are regulated by radial spokes and central apparatus and drive ciliary motility.

We found a structure in the inside lumen of the A-tubule in zone 1 cilia (Fig. 3). This structure appears to be one of the lumen proteins of doublet microtubules, called MIPs. A recent study suggested that MIPs directly affect the stability of doublet microtubules (Ichikawa et al., 2019). Although it is unknown why the structure is present in zone 1 cilia but not distinctly in zone 5 cilia, it is possible that the structure might stabilize the long axonemes or reinforce the mechanical weakness of doublet microtubules due to the lack of twoheaded dyneins.

The loss of both the outer arm dynein and the twoheaded inner arm dynein (IADs) in zone 1 cilia suggests that a common component is involved in the assembly of both dyneins. Dynein subunits, as well as other axonemal components, are assembled in the cytoplasm and transported into cilia by intraflagellar transport (Fowkes and Mitchell, 1998; Inaba and Mizuno, 2016). Several factors for cytoplasmic assembly of axonemal dyneins have been reported so far, including DNAAF1 (LRRC50/ODA7) (Duquesnoy et al., 2009), DNAAF2 (Ktu/PF13) (Omran et al., 2008) and DNAAF3 (PF22) (Loges et al., 2009; Mitchison et al., 2012), DYX1C1/DNAAF4 (Tarkar et al., 2013; Yamamoto et al., 2017), DNAAF5 (HEATR2) (Horani et al., 2012), LRRC6 and its associated assembly factor ZMYND10 (Zariwala et al., 2013), and SPAG1 (Knowles et al., 2010). In Chlamydomonas, two assembly factors for IADs are known. MOT48 (IDA10) contains a PIH domain and is involved in the assembly of IAD species b, c and d (Yamamoto et al., 2010). The other is the MIA complex, which is located at the distal part of IAD f/ I1 and is involved in the assembly of this dynein (Yamamoto et al., 2013, 2017). However, none of these factors has been shown to be specifically involved in the assembly of twoheaded dyneins. Thus, studies on the mechanism of how two-headed dyneins are lost in endostyle zone 1 would give a key clue for finding novel cytoplasmic assembly factors for two-headed axonemal dynein and for determining the mechanism of the region-specific gene regulation of these factors.

ACKNOWLEDGMENTS

We thank Dr. Ryuji Yanase for his help in TEM imaging. We also thank all staff members of the Education and Research Center of Marine Bio-Resources, Tohoku University, International Coastal Research Center, The Atmosphere and Ocean Research Institute (University of Tokyo), and the National BioResource Project for supplying *C. intestinalis* and *C. savignyi*. This work was supported in part by Grants-in-Aid Nos. 22370023 and 25291069 for Scientific Research (B) to KI from the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT).

COMPETING INTERESTS

The authors declare they have no competing interests.

AUTHOR CONTRIBUTIONS

All authors designed the research; AK performed electron microscopy and western blot analysis; KI performed video observation, immunofluorescent staining, TEM image analysis and western blotting; All authors performed data analysis and wrote the manuscript.

SUPPLEMENTARY MATERIALS

Supplementary material for this article is available online. (URL: https://doi.org/10.2108/zs200095)

Supplementary Movie S1. Motility of zone 1 long cilia. The movie plays at real speed. Scale bar, 100 μ m.

Supplementary Movie S2. Some cilia in zone 1 showing movement with larger amplitudes and shorter wavelengths. The movie plays at a speed of $\times 0.6$. Scale bar, 50 µm.

Supplementary Movie S3. Motility of zone 5 cilia. The movie plays at real speed. Scale bar, 20 $\mu m.$

REFERENCES

- Afzelius BA (1976) A human syndrome caused by immotile cilia. Science 193: 317–319
- Brokaw CJ, Kamiya R (1987) Bending patterns of *Chlamydomonas* flagella: IV. Mutants with defects in inner and outer dynein arms indicate differences in dynein arm function. Cell Motil Cytoskeleton 8: 68–75
- Burighel P, Cloney RA (1997) Urochordata: Ascidiacea. In "Microscopic Anatomy of Invertebrates. Vol. 15. Hemichordata, Chaetognatha, and the Invertebrate Chordates" Ed by FW Harrison and EE Ruppert, Wiley-Liss, New York, pp 221–347
- Duquesnoy P, Escudier E, Vincensini L, Freshour J, Bridoux AM, Coste A, et al. (2009) Loss-of-function mutations in the human ortholog of *Chlamydomonas reinhardtii* ODA7 disrupt dynein arm assembly and cause primary ciliary dyskinesia. Am J Hum Genet 85: 890–896
- Fowkes ME, Mitchell DR (1998) The role of preassembled cytoplasmic complexes in assembly of flagellar dynein subunits. Mol Biol Cell 9: 2337–2347
- Fujita H, Nanba H (1971) Fine structure and its functional properties of the endostyle of ascidians, *Ciona intestinalis*. A part of phylogenetic studies of the thyroid gland. Z Zellforsch 121: 455– 469
- Gadelha C, Wickstead B, McKean PG, Gull K (2006) Basal body and flagellum mutants reveal a rotational constraint of the central pair microtubules in the axonemes of Trypanosomes. J Cell Sci 119: 2405–2413
- Gardner LC, O'Toole E, Perrone CA, Giddings T, Porter ME (1994)

Components of a "dynein regulatory complex" are located at the junction between the radial spokes and the dynein arms in *Chlamydomonas* flagella. J Cell Biol 127: 1311–1325

- Gibbons BH, Gibbons IR (1979) Relationship between the latent adenosine triphosphatase state of dynein 1 and its ability to recombine functionally with KCI-extracted sea urchin sperm flagella. J Biol Chem 254: 197–201
- Gibbons BH, Baccetti B, Gibbons IR (1985) Live and reactivated motility in the 9+0 flagellum of Anguilla sperm. Cell Motil 5: 333-350
- Gibbons IR (1981) Cilia and flagella of eukaryotes. J Cell Biol 91: 107s-124s
- Habermacher G, Sale WS (1997) Regulation of flagellar dynein by phosphorylation of a 138-kD inner arm dynein intermediate chain. J Cell Biol 136: 167–176
- Holley MC (1986) Cell shape, spatial patterns of cilia, and mucusnet construction in the ascidian endostyle. Tissue Cell 18: 667–684
- Horani A, Druley TE, Zariwala MA, Patel AC, Levinson BT, Van Arendonk LG, et al. (2012) Whole-exome capture and sequencing identifies HEATR2 mutation as a cause of primary ciliary dyskinesia. Am J Hum Genet 91: 685–963
- Hozumi A, Satouh Y, Makino Y, Toda T, Ide H, Ogawa K, et al. (2006) Molecular characterization of *Ciona* sperm outer arm dynein reveals multiple components related to outer arm docking complex protein 2. Cell Motil Cytoskeleton 63: 591–603
- Hozumi A, Padma P, Toda T, Ide H, Inaba K (2008) Molecular characterization of axonemal proteins and signaling molecules responsible for chemoattractant-induced sperm activation in *Ciona intestinalis*. Cell Motil Cytoskeleton 65: 249–267
- Ichikawa M, Khalifa AAZ, Kubo S, Dai D, Basu K, Maghrebi MAF, et al. (2019) Tubulin lattice in cilia is in a stressed form regulated by microtubule inner proteins. Proc Natl Acad Sci USA 116: 19930–19938
- Inaba K (2003) Molecular architecture of the sperm flagella: molecules for motility and signaling. Zool Sci 20: 1043–1056
- Inaba K (2011) Sperm flagella: comparative and phylogenetic perspectives of protein components. Mol Hum Reprod 17: 524– 538
- Inaba K (2015) Calcium sensors of ciliary outer arm dynein: functions and phylogenetic considerations for eukaryotic evolution. Cilia 4: 6
- Inaba K, Mizuno K (2016) Sperm dysfunction and ciliopathy. Reprod Med Biol 15: 77–94
- Ishijima S, Sekiguchi K, Hiramoto Y (1988) Comparative study of the beat patterns of American and Asian horseshoe crab sperm: evidence for a role of the central pair complex in forming planar waveforms in flagella. Cell Motil Cytoskeleton 9: 264–270
- Itoh A, Inaba K, Ohtake H, Fujinoki M, Morisawa M (2003) Characterization of a cAMP-dependent protein kinase catalytic subunit from rainbow trout spermatozoa. Biochem Biophys Res Commun 305: 855–861
- Jin Y, Yaguchi Y, Shiba K, Yamada L, Yaguchi J, Shibata D, et al. (2013) Glutathione transferase theta in apical ciliary tuft regulates mechanical reception and swimming behavior of sea urchin embryos. Cytoskeleton 70: 453–470
- Kamiya R (1995) Exploring the function of inner and outer dynein arms with *Chlamydomonas* mutants. Cell Motil Cytoskeleton 32: 98–102
- Kamiya R, Yagi T (2014) Functional diversity of axonemal dyneins as assessed by in vitro and in vivo motility assays of *Chlamydomonas* mutants. Zool Sci 31: 633–644
- Kamiya R, Kurimoto E, Muto E (1992) Two types of *Chlamydomonas* flagellar mutants missing different components of inner-arm dynein. J Cell Biol 112: 441–447
- King SJ, Dutcher SK (1997) Phosphoregulation of an inner dynein

arm complex in *Chlamydomonas reinhardtii* is altered in phototactic mutant strains. J Cell Biol 136: 177–191

- Knowles MR, Ostrowski LE, Loges NT, Hurd T, Leigh MW, Huang L, et al. (2013) Mutations in SPAG1 cause primary ciliary dyskinesia associated with defective outer and inner dynein arms. Am J Hum Genet 93: 711–720
- Konno A, Kaizu M, Hotta K, Horie T, Sasakura Y, Ikeo K, et al. (2010) Distribution and structural diversity of cilia in tadpole larvae of the ascidian *Ciona intestinalis*. Dev Biol 337: 42–62
- Konno A, Setou M, Ikegami K (2012) Ciliary and flagellar structure and function-their regulations by posttranslational modifications of axonemal tubulin. Int Rev Cell Mol Biol 294: 133–170
- Konno A, Shiba K, Cai C, Inaba K (2015) Branchial cilia and sperm flagella recruit distinct axonemal components. PLoS One 10: e0126005
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature 227: 680-685
- Loges NT, Olbrich H, Becker-Heck A, Häffner K, Heer A, Reinhard C, et al. (2009) Deletions and point mutations of LRRC50 cause primary ciliary dyskinesia due to dynein arm defects. Am J Hum Genet 85: 883–889
- Marshall WF, Nonaka S (2006) Cilia: tuning in to the cell's antenna. Curr Biol 16: R604–614
- Mitchell DR (2007) The evolution of eukaryotic cilia and flagella as motile and sensory organelles. Adv Exp Med Biol 607: 130– 140
- Mitchison HM, Schmidts M, Loges NT, Freshour J, Dritsoula A, Hirst RA, et al. (2012) Mutations in axonemal dynein assembly factor DNAAF3 cause primary ciliary dyskinesia. Nat Genet 44: 381–389
- Mizuno K, Padma P, Konno A, Satouh Y, Ogawa K, Inaba K (2009) A novel neuronal calcium sensor family protein, calaxin, is a potential Ca²⁺-dependent regulator for the outer arm dynein of metazoan cilia and flagella. Biol Cell 101: 91–103
- Mizuno K, Shiba K, Okai M, Takahashi Y, Shitaka Y, Oiwa K, et al. (2012) Calaxin drives sperm chemotaxis by Ca²⁺-mediated direct modulation of a dynein motor. Proc Natl Acad Sci USA 109: 20497–20502
- Nomura M, Inaba K, Morisawa M (2000) Cyclic AMP- and calmodulin-dependent phosphorylation of 21 kDa and 26 kDa proteins in axoneme is a prerequisite for SAAF-induced motile activation in ascidian spermatozoa. Develop Growth Differ 42: 129– 138
- Ogawa K, Inaba K (2006) Ap58: a novel in situ outer dynein armbinding protein. Biochem Biophys Res Commun 343: 385–930
- Omran H, Kobayashi D, Olbrich H, Tsukahara T, Loges NT, Hagiwara H, et al. (2008) Ktu/PF13 is required for cytoplasmic pre-assembly of axonemal dyneins. Nature 456: 611–616
- Padma P, Hozumi A, Ogawa K, Inaba K (2001) Molecular cloning and characterization of a thioredoxin/nucleoside diphosphate kinase related dynein intermediate chain from the ascidian, *Ciona intestinalis*. Gene 275: 177–183
- Padma P, Satouh Y, Wakabayashi K, Hozumi A, Ushimaru Y, Kamiya R, et al. (2003) Identification of a novel leucine-rich repeat protein as a component of flagellar radial spoke in the ascidian Ciona intestinalis. Mol Biol Cell 14: 774–785
- Pennachetti CA (1984) Functional morphology of the branchial basket of Ascidia paratropa (Tunicata, Ascidiacea). Zoomorphology 104: 216–222
- Petersen JK (2007) Ascidian suspension feeding. J Exp Mar Biol Ecol 342: 127–137
- Porter ME, Sale WS (2000) The 9 + 2 axoneme anchors multiple inner arm dyneins and a network of kinases and phosphatases that control motility. J Cell Biol 151: F37–F42
- Sasaki A, Miyamoto Y, Satou Y, Satoh N, Ogasawara M (2003) Novel endostyle-specific genes in the ascidian *Ciona*

intestinalis. Zool Sci 20: 1025-1030

- Satir P, Christensen ST (2008) Structure and function of mammalian cilia. Histochem Cell Biol 129: 687–693
- Satouh Y, Inaba K (2009) Proteomic characterization of sperm radial spokes identifies a novel spoke protein with an ubiquitin domain. FEBS Lett 583: 2201–2207
- Satouh Y, Padma P, Toda T, Satoh N, Ide H, Inaba K (2005) Molecular characterization of radial spoke subcomplex containing radial spoke protein 3 and heat shock protein 40 in sperm flagella of the ascidian *Ciona intestinalis*. Mol Biol Cell 16: 626– 636
- Tarkar A, Loges NT, Slagle CE, Francis R, Dougherty GW, Tamayo JV, et al. (2013) DYX1C1 is required for axonemal dynein assembly and ciliary motility. Nat Genet 45: 995–1003
- Thorndyke MC (1978) Evidence for a 'mammalian' thyroglobulin in endostyle of the ascidian *Styela clava*. Nature 271: 61–62
- Wirschell M, Hendrickson T, Sale WS (2007) Keeping an eye on I1: 11 dynein as a model for flagellar dynein assembly and regulation. Cell Motil Cytoskeleton 64: 569–579
- Wirschell M, Yang C, Yang P, Fox L, Yanagisawa H, Kamiya R, et al. (2009) IC97 is a novel intermediate chain of I1 dynein that interacts with tubulin and regulates interdoublet sliding. Mol Biol Cell 20: 3044–3054

- Yagi T, Kamiya R (2000) Vigorous beating of *Chlamydomonas* axonemes lacking central pair/radial spoke structures in the presence of salts and organic compounds. Cell Motil Cytoskeleton 46: 190–199
- Yamamoto R, Hirono M, Kamiya R (2010) Discrete PIH proteins function in the cytoplasmic preassembly of different subsets of axonemal dyneins. J Cell Biol 190: 65–71
- Yamamoto R, Song K, Yanagisawa HA, Fox L, Yagi T, Wirschell M, et al. (2013) The MIA complex is a conserved and novel dynein regulator essential for normal ciliary motility. J Cell Biol 201: 263–278
- Yamamoto R, Obbineni JM, Alford LM, Ide T, Owa M, Hwang J, et al. (2017) *Chlamydomonas* DYX1C1/PF23 is essential for axonemal assembly and proper morphology of inner dynein arms. PLoS Genet 13: e1006996
- Zariwala MA, Gee HY, Kurkowiak M, Al-Mutairi DA, Leigh MW, Hurd TW, et al. (2013) ZMYND10 is mutated in primary ciliary dyskinesia and interacts with LRRC6. Am J Hum Genet 93: 336– 345

(Received June 11, 2020 / Accepted August 9, 2020 / Published online October 29, 2020)