

Guiding Light: Mechanisms and Adjustments of Environmental Light Interpretation with Insights from *Platynereis dumerilii* and Other Selected Examples

Authors: Wulf, Paul O., Häfker, N. Sören, Hofmann, Kaelin, and Tessmar-Raible, Kristin

Source: Zoological Science, 42(1) : 52-59

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zs240099>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

[REVIEW]

Guiding Light: Mechanisms and Adjustments of Environmental Light Interpretation With Insights From *Platynereis dumerilii* and Other Selected Examples

Paul O. Wulf^{1,2,5}, N. Sören Häfker³, Kaelin Hofmann^{1,5}, and
Kristin Tessmar-Raible^{1,3,4,5*}

¹Max Perutz Labs, University of Vienna, 1030 Vienna, Austria

²Vienna BioCenter PhD Program, Doctoral School of the University of Vienna,
Medical University of Vienna, 1030 Vienna, Austria

³Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research,
27570 Bremerhaven, Germany

⁴Institute for Chemistry and Biology of the Marine Environment (ICBM),
School of Mathematics and Science, Carl von Ossietzky
Universität Oldenburg, 26129 Oldenburg, Germany

⁵Department of Neuroscience and Developmental Biology,
Faculty of Life Science, 1030 Vienna, Austria

Animals possess many light-sensitive molecules. They exist as dedicated photoreceptors, or as byproducts of biochemical reactions. Their numbers are often high even in species that live in environments that humans would consider dark, as well as in species that are considered comparably simple (e.g., worms, cnidarians).

But why are there so many photoreceptors? We provide some considerations on this question. Light conveys a significant amount of information to animals, through complex spectral and intensity changes, often specific to the spatial and temporal ecological niches a species inhabits. We discuss that the large number of opsins and cryptochromes, often also present outside the eyes and partially co-expressed, represent adaptation mechanisms to the highly complex light environment within a given niche. While theoretical, it is a plausible hypothesis given that most experimentally tested opsins and cryptochromes have been shown to be functional photoreceptors. The example of lunar and solar timing of the marine annelid *Platynereis dumerilii* provides insight on how animals use the biochemical and cellular properties of different photoreceptors to decode solar versus lunar light, and their different adaptations in *Drosophila melanogaster*. We suggest that the future understanding of biological processes will strongly benefit from comparative lab and field work on the same species, and provide a first example for such work in *P. dumerilii*. Finally, we point out that work on animal light detection systems and their adaptability is crucial to understand the impact of anthropogenic changes on species and ecosystems.

Key words: light receptors, opsin, cryptochrome, lunar timing, solar timing, *Platynereis dumerilii*

INTRODUCTION: THE HIGH COMPLEXITY OF LIGHT INFORMATION

Regular changes to light intensity and spectrum have occurred ever since the dawn of life. These changes harbor critical (and often regular) information about an organism's location and the state of the environment (e.g., extent of vegetation, weather conditions, passing of potential prey or predators) and time (from tidal to annual). Light of biological

origin (bioluminescence and fluorescence), serves as communication signals (e.g., Sarfati et al., 2021, 2023; Hensley et al., 2023), but can also be used to optimize physiological processes by serving as a wavelength converter (Bollati et al., 2022). Given that light can thus provide plenty of information, whose correct interpretation can be highly advantageous, this provides a likely explanation for the presence of the large number and diversity of animal light receptors. Light can be measured by light-sensitive processes (i.e., light-sensitive biochemical reactions, such as vitamin D synthesis or the production of urocanic acid in mammalian skin) or by dedicated receptors (i.e., opsins and cryptochromes);

* Corresponding author. E-mail: kristin.tessmar@mfp.ac.at
doi:10.2108/zs240099

both types of processes have been shown to impact animal physiology and behavior (reviewed in Mat et al., 2024). Given the multitude of processes that can be affected by light, but should be affected only by the “right” light changes, it also implies that visual and non-visual photoreceptors have to be properly adapted to the specific spatial and temporal ecological light niches a species inhabits (Carothers and Jakšić, 1984; Hut et al., 2012; Menchetti et al., 2019; Häfker et al., 2022b).

One particularly complex cue highly influential for animal behavior and physiology consists of the changes in light spectrum and intensity over time (Fig. 1). Probably most obvious and best known to humans are the daily and annual (i.e., daylength in the temperate zones, monsoon/non-monsoon in the subtropical/tropical zones) changes of the solar cycle (Fig. 1B–E). Many animals use this information for daily and annual timing, but although much focus had been on light intensity and photoperiod (i.e., daylength), evidence for the importance of relative spectral changes for precise timing is increasing (Walmsley et al., 2015; Veedin Rajan et al., 2021). The latter requires an ability of the light sensory systems to decode such relative spectral shifts. The exact spectral shifts depend on the respective ecological niches and hence the decoding is expected to be optimized by species-specific adaption mechanisms. Another impor-

tant, yet often less regarded, temporal cue is moonlight (Fig. 1A, C–E). Its duration and timing relative to the daily solar light cycle provides animals with information on the month, but also time of day relative to the month (Fig. 1C–E). When taking dusk and dawn into consideration, this results in highly complex, seasonally changing light spectra and intensity changes (Fig. 1) (Nordtug and Melø, 1988; Veedin Rajan et al., 2021), again also ecological-niche dependent, because the light conditions themselves will be influenced by plant or algal growth and the animals' behaviors. In the marine environment, the aspect of temporal timing cues gets even more complicated by the in-parallel-occurring tidal water depths changes (Fig. 2). While some coastal regions, such as the Island of Ischia, Italy in the Mediterranean experience water depth changes only in the range of 10–50 cm and are hence within the range of waves, other places, such as Roscoff, France on the Atlantic Coast of Britany experience tidal changes in the range of several meters, while others, like Osaka, Japan facing the Seto Inland Sea are in between (Fig. 2A–C). Whether or not there are several meters of water above organisms has again significant consequences for the light intensity and spectrum that can reach them from both sun- and moonlight due to the light scattering and absorbance properties of the water, which will also differ depending on turbidity from organic and inor-




Fig. 1. Complex natural light cues with much temporal information for animals. (A, B) schematized lunar and solar cycle. For (B), the yellow/gray pie chart pieces indicate day and night length, respectively. (C–E) Typical solar/lunar light regimes at different marine locations. Lunar cycle across the annual equinoxes and solstices are shown. White: lunar light, yellow: solar light, orange astronomical dusk/dawn periods, black: no natural sun or moonlight visible at location. Figure from C. Amort under CC-BY license.




Fig. 2. Influence of light spectrum and intensity by tides. **(A–C)** Tidal changes at marine locations known to harbor animals with complex light-dependent chronobiology. **(D, E)** Light changes depending on water depth, redrawn from Mat et al. (2024). Figure from C. Amort under CC-BY license.

ganic material (Fig. 2D, E) (Speight and Henderson, 2010; Depauw et al., 2012; Veedin Rajan et al., 2021).

DECODING THE LIGHT INFORMATION

How can organisms decode the relevant light information from the complex spatially and temporally changing signals that reach them? As mentioned above, the number of animal genes encoding proteins that belong to the known photoreceptor classes of opsins and cryptochromes is – at first glance – unexpectedly high and diverse (Mat et al., 2024). The frequently occurring argument that likely many of those may not function as actual light receptors is ad hoc difficult to prove or disprove. While loss of sensitivity to photons has certainly happened under some circumstances, such as the mechanosensory function of opsins in the Johnston organ of the fly antenna (Senthilan et al., 2012; Katana et al., 2019), even in this case it likely represents an evolutionary derivation rather than the original functionality of the opsin, as comparative functional analyses in annelids suggest (Revilla et al., 2021). A more extended discussion on this topic is present elsewhere (Mat et al., 2024). Interestingly, it is at present even unclear if a non-photon-dependent function is evolutionarily ancient for the cryptochrome type-II class, best studied in mammals, where they function as light-independent transcriptional repressors (Michael et al., 2017). In contrast to their mammalian light-independent

function, the teleost CRYII-class member, cry1a, is light-sensitive (Tamai et al., 2007), suggesting specific mammalian adaptations, possibly during the nocturnal bottleneck period Cretaceous–Paleogene extinction event (Gerkema et al., 2013). This is consistent with the idea that cryptochromes are originally light-sensitive, given their evolutionary close relatedness to light-dependent DNA repair enzymes, the photolyases (Gehring and Rosbash, 2003; Oliveri et al., 2014). In summary, it can be stated that the evidence for a generalization of presumptive non-photon-dependent functions of opsins is low. Thus, the many animal photoreceptors likely represent adaptations to the species-specific light niches. These adaptations can be on several levels, most obvious is the maximal spectral sensitivity and the spectral sensitivity range, which cannot be predicted from the amino acid sequence and structure. Noteworthy examples for this are the two ciliary-type opsins in the marine bristle worm *Platynereis dumerilii*, both close in sequence, but differing significantly in the spectral sensitivity properties (activating light range: U-VA/deep violet for c-opsin1, blue/green for c-opsin2; Tsukamoto et al., 2017; Ayers et al., 2018; Veedin Rajan et al., 2021).

Bistability versus monostability of opsins, and differing downstream signaling cascades provide additional features that can be used by organisms to optimize their photoreceptors' functions to a given environment. It is important to note

that bistability, the ability of opsins to change the state of its co-factor (typically between 11-cis and all-trans retinal) back and forth by the absorption of different wavelengths makes them “evaluators” of the respective environmental spectra, because the speed and level of re-activation depend on the available wavelength (Zhong et al., 2012; Tsukamoto, 2014). In contrast, monostable opsins need a catalyzing photoisomerase to return retinal to the (typically) 11-cis state and hence depend on the kinetics of this enzyme for their re-activation (Zhong et al., 2012; Tsukamoto, 2014). From the signaling perspective, a single monostable opsin acts more like a switch, rather than the more constant signaling bistable opsins can perform. Again, examples of likely specific environmental adaptations exist. In medaka fish, the opsins *tmt-opsin1b* and *tmt-opsin2* are co-expressed in several brain regions, in addition to another opsin family, the *val-opsins* (Fischer et al., 2013). While *tmt-opsin1* is bistable, *tmt-opsin2* is monostable (Sakai et al., 2015), likely contributing to differential environmental signal interpretation. Of note, the signaling properties of medaka *val* opsins are still unclear. Truly understanding the combined functions of such non-visual opsins in an environmental context will certainly be challenging future endeavors, as mutants in just two of these opsins exhibit age- and temperature-dependent synergistic effects (Fontinha et al., 2021; Zekoll et al., 2021).

Yet, it is important to tackle these challenges, because such co-expression of photoreceptors, both opsins and cryptochromes, may turn out to be the rule rather than the exception in invertebrates and vertebrates. A notable example in an environmental context is the regulation of growth and feeding in echinoderms (Liu et al., 2025), where larvae can regulate such intricate processes as the pyloric (lower stomach) opening in response to light (Yaguchi and Yaguchi, 2021), likely tuning digestive processes to time and position in the water.

Similarly, the study of eyes in the diversity of animal species has revealed a complexity of photoreceptors, their wavelength, and downstream signaling that show how different from “conventional” molecular text book model systems other animals can function. The rhabdomeric photoreceptors of the terrestrial slug *Limax* co-express at least three functional opsins and one retinochrome with different cellular signaling features (Matsuo et al., 2023). Similarly, single cell sequencing revealed the co-expression of several *opsins* and the light-receptive cryptochrome *l-cry* in the rhabdomeric eyes of *Platynereis*, at least partially co-expressed in the same cells. While several of them function in the differentiated photoreceptors, one of them, *c-opsin1*, is present in the neurogenic precursor cells in the eyes prior to photoreceptor differentiation. In *c-opsin1* loss-of-function mutants, combined single-cell RNA sequencing and labeling of actively dividing cells by EdU suggest that *c-opsin1* regulates divisions and differentiation of these neural stem cells in the eye, separate from its function in the differentiated ciliary photoreceptors in the worm’s median brain (Milivojev et al., 2024 preprint). This leads to the intricate question of why these bristle worms would adjust their eye growth based on light conditions? We hypothesize that this may allow them to adjust to the different depths the species has been reported to inhabit, given that *Platynereis* has been documented for coastal habitats across a range of

depths from 1–5 m, but also at 10 m and below (Gambi et al., 1992; Somaschini et al., 1998; Galparsoro et al., 2012; Özpolat et al., 2021). Individuals settling at these different depths’ light conditions will significantly differ (Fig. 2, also see discussion in Häfker et al., 2024). Another reason for the likely light-dependent adjustment of eye growth could be the seasonal changes in the light conditions these worms also experience (Veedin Rajan et al., 2021). In either case, the actual meaning of larger eyes for behavior and/or physiology of the animals remains a yet unresolved puzzle.

Another example of the astonishing complexity and sophisticated environmental adaptation to light are the various opsins present in sea urchins (Raible et al., 2006), which appear to render the adult animals into whole visual systems, i.e., the animal functions as an eye (Kirwan et al., 2024 preprint). This again raises many fascinating questions, especially in the context of the different environments sea urchins inhabit.

Most of the above-mentioned light-sensory systems likely represent adaptation to light environments under daylight. However, lunar light also provides a significant amount of information to animals, which can be used for orientation (Foster et al., 2019) or timing (Kronfeld-Schor et al., 2013; Last et al., 2016; Andreatta and Tessmar-Raible, 2020). At least under some instances, this requires animals to distinguish lunar and solar light and – in order to phase synchronize their inner timing systems – even moon phase.

How animals adapted to decode moon phase: the *Platynereis* example

The marine bristle worm *Platynereis dumerilii* exhibits a reproductive strategy tightly linked to the lunar cycle. By timing their sexual maturation and synchronized swarming events through an endogenous oscillator, aka inner calendar or circalunar clock, which is entrained by nocturnal light of full moon duration, these worms enhance fertilization success. This synchronization ensures simultaneous maturation, a strategy also observed in many other externally fertilizing marine species, but even increasingly documented for terrestrial animals (reviewed in Ritter and Tessmar-Raible, 2024).

Accurate interpretation of moonlight against other light stimuli is crucial for synchronizing the circalunar clock among individuals. The cryptochrome L-Cry plays a key role in this differentiation. Loss-of-function mutants of *l-cry* adjust their circalunar clocks more quickly to shifts in moon phase under artificial light, but do not show significant disadvantages under naturalistic light stimuli (Poehn et al., 2022). This suggests that correctly identifying the full moon requires molecular collaboration between L-Cry and at least one other photoreceptor.

To correctly identify the full moon, photoreceptors must detect both the intensity and the duration of the moonlight stimulus. Intensity alone is insufficient because – dependent on weather and other environmental conditions – different moon phases can produce similar light levels, making it difficult to distinguish the full moon based solely on brightness. Likewise, duration of a light stimulus alone cannot be relied upon because sunlight would be confusing to animals (Fig. 1C–E). Therefore, combining information about both the intensity and duration of moonlight allows for a determina-

tion of moon phases, including the full moon.

Biochemically, L-Cry forms a homodimer that predominantly resides in the nucleus under dark conditions (Poehn et al., 2022; Vu et al., 2023). This dimer state serves as a prerequisite for its ability to distinguish between the vastly different light intensities of moonlight and sunlight, which amount to 10^5 – 10^6 orders of magnitude difference (Posch et al., 2010; Poehn and Tessmar-Raible, 2022). Each monomer within the dimer contains the cofactor flavin adenine dinucleotide (FAD), which is sensitive to light. When exposed to light mimicking moonlight intensities and spectra for an extended period of several hours, only one of the two FAD molecules within the dimer becomes photoreduced, supposedly resulting in an asymmetric dimer: one L-Cry in the dimer has its FAD photoreduced. Protein structural changes cause the oxidized FAD in the second L-Cry of the dimer to require more energy for the photoreduction process. This results in a scenario under which the two FADs in the L-Cry dimer have different sensitivities to light. Notably, even prolonged moonlight exposure of up to 12 hours does not lead to the reduction of this second FAD (Poehn et al., 2022).

In contrast, under the much more intense illumination of sunlight, both FADs in the dimer are rapidly photoreduced within minutes. Sunlight provides enough photons to overcome the lower quantum efficiency of the second L-Cry-bound FAD within minutes (Fig. 3A) (Poehn et al., 2022). This full activation induces a conformational change in L-Cry, leading to the dissociation of the dimer into monomers (Vu et al., 2023). The monomers are then exported from the nucleus, where they can still be detected for 30–60 min. Subsequently they are degraded in the cytoplasm (Poehn et al., 2022; Vu et al., 2023). However, analyses of L-Cry protein levels under continuous sunlight exposure reveal that apparently this is certainly not capturing the entire complexity of the process, because L-Cry can be re-detected after 8 hrs, but the exact level may depend itself on the phase of the circalunar clock (Zurl et al., 2022; Coric, Orel, and Tessmar-Raible, unpublished).

The differential sensitivity of the two L-Crys within a dimer allows L-Cry to act both as a highly sensitive moonlight detector and as a sunlight sensor, but importantly, due to its kinetics under moonlight, it provides the animals with information on moon phase, which is crucial for setting their inner calendars to the same phase across the population. Moonlight needs several hours to photoreduce the first FAD in the L-Cry dimer, a duration of moonlight that in the natural habitat of *Platynereis dumerilii* in the Bay of Naples is only reached by the times around full moon.

While the inner monthly oscillator thus requires a detection of moon phase, classical work has shown for species as diverse as humans and algae that light intensity also impacts the period length of the ~24-hr circadian oscillators (Aschoff, 1960, 1965). This means that the exposure to moonlight could also influence the daily timing. Comparative work on the bristle worm *Platynereis* versus *Drosophila* suggests that such influences can be tuned depending on the temporal niches of the species. *Platynereis*, as nocturnal animal, uses the exposure to moonlight (together with influences by its circalunar oscillator) to change the period length of its plastic ~24-hr circadian-circalunidian oscillator to tune the onset of the population-synchronized swarming to the hour just before the rise of the moon (Zurl et al., 2022). *Drosophila melanogaster* as a crepuscular animal should avoid shifting its daily timing in response to moonlight, as dusk and dawn times are independent of this. Consistently, whereas *Platynereis* uses its moonlight detection system to tune its ~24-hr clock, *Drosophila* uses such a system to prevent large effects of moonlight. While both animals use their cry-I type system in combination with rhabdomeric opsins, the systems function mechanistically differently (Zurl et al., 2022). In contrast to the L-Cry of *Platynereis*, the fly's ortholog dCry does not respond to moonlight as a purified protein. The nevertheless high sensitivity suggested by functional genetic analyses (Vinayak et al., 2013) may thus be a property enhanced by endogenous cellular processes.

Both *Platynereis* and *Drosophila* can then use a combi-




Fig. 3. Concepts for decoding complex natural light information by different photoreceptors. **(A)** Model of an L-Cry dimer, which shows differential quantum efficiency depending on the states of its FAD co-factors. In dark state, both FADs are oxidized and photon capture results in photoreduction of the first FAD with high quantum efficiency, which can be achieved by extended moonlight illumination. The FAD photoreduction results in a presumptive structural change of the L-Cry protein dimer. A subsequent photoreduction of the second FAD is now much less quantum efficient and can only be reached by the high energy of sunlight. Hypothesis and figure modified from Poehn et al. (2022). **(B)** Hypothesis about how the combination of photoreceptors with different light sensitivities and/or biochemical light activation properties can in its combinatoric decode complex natural light information. Figure from C. Amort under CC-BY license.

natorial code of photoreceptors to detect moonlight due to the differential response properties of their photoreceptors to sun- versus moonlight (Fig. 3B). Such combinatorics allow for the identification of the type of natural light, but not the duration of moon phase, which for *Drosophila* (or other diurnal animals) is, however, likely not relevant, as they mainly aim to protect their endogenous circadian oscillators from being impacted by this type of dim nocturnal light irrespective of the specific moon phase. This comparison illustrates how similar proteins can diverge in function based on the ecological requirements of different species.

ANIMAL ADAPTATIONS TO THEIR NATURAL LIGHT NICHE IN A HUMAN-LIT WORLD

The intricate and specifically tuned light sensory mechanisms present in animals are nowadays challenged by the changes humans are making to the light environment. For scientists, this is relevant from at least two different perspectives: first, there are the direct environmental interferences that humans do by artificial illumination, typically referred to as artificial light at night (ALAN) (Davies et al., 2014; Kronfeld-Schor et al., 2017; Walker II et al., 2019). Humans directly (by artificial light sources) and indirectly (e.g., by sky glow) light up the night and illuminate places, such as the oceans' dark/twilight zones that naturally have no or much less light. On the other end, solar panels and other devices are put in places on land and water and block out light that

would normally shine on the organisms underneath. There are even discussions about artificially reducing the amount of light that reaches the earth's surface to reduce global warming. While the latter is targeting the sun, it will certainly also affect lunar light. Furthermore, natural light intensity and spectrum can be affected by snow and ice loss, increase in algal growth or vegetation, etc. (e.g., Häfker et al., 2022a, b). Given the complexity and importance that natural light has for animals it can certainly be stated that the effects of such geoengineering proposals and interventions on complex environmental balances can at present not be realistically estimated and would be likely to unhinge long-evolved systems we may not have even discovered yet. Thus, research into the functions and adaptations of light sensing systems in animals is highly relevant not just for the understanding of biological mechanisms, but to also educate the general public and political decision makers about the consequences human interference with natural sun- and moonlight can have.

Second, when performing lab experiments, we as scientists typically work with animals raised and tested under lab conditions, i.e., artificial illumination. While the lights may appear similarly white to us, their spectrum and intensity differs significantly from natural light (e.g., Dekens et al., 2017), with consequences for the results of experiments.

This calls for a consequent reporting of the experimentally used light conditions, but also for comparative analyses




Fig. 4. Comparative diel and lunar transcriptomic analyses of worms under lab vs. field conditions. (A–C) Experimental setup in the field. The worms were kept in UV-transparent plexiglass containers at Haidhof, Austria, an area without artificial light sources. Tanks contained a filtering and temperature control system (visible in [A]), to ensure water quality and minimize the impact of temperature. For sampling, worms were allowed to pre-settle on plexiglass squares with glued glass slides that were easily harvested at time of sampling (visible in [C]). (D) Temporal sampling scheme. Worms were sampled every 4 hours over a 24 h period in four biological replicates (4–5 heads pooled) and analyzed by Illumina-sequencing. (E, F) Initial Principal Component Analysis (PCA) of RNA-Seq data for wildtype and *l-cry* mutants ($-/-$) samples across lab and field conditions. Each biological replicate is shown as an individual data point, color-coded according to the corresponding moon phase and the environment where sampled. Details on the experimental set-up are available from the authors upon request.

of experiments under lab versus more natural (light) conditions.

Driven by these considerations, we designed tanks for outside maintenance of *Platynereis dumerilii* worms (Fig. 4A–C), maintained them under these conditions for several months and subsequently sampled their heads for transcriptomic analyses over the 24 h cycle and at two different lunar phases (new moon: NM and full moon: FM, Fig. 4D). The tanks were designed such that the temperature was regulated and thus constant over the course of the experiment, to focus on the effects of illumination.

The same sampling scheme was in parallel mimicked in the lab, allowing us to assess the specific effects of natural versus artificial light conditions. The amount of data resulting from this approach was large and will require more long-term dedicated analyses beyond the scope of this article. However, initial principal component analyses (PCAs) for the data structure already reveal two interesting aspects. First, for wild-type animals, the difference between lab versus field tank animals is significant for transcript level, exceeding the difference between moon phases and time of day (Fig. 4E). Second, this is different for mutants of the light-interpreter *l-cry*. These mutant worms show fewer differences between lab and field tanks, being overall closer to lab counterparts than are the lab and field wild-type worms (Fig. 4E, F). While PCAs represent differences caused by several variables, these plots are consistent with our interpretation of the lab experiments under artificial versus naturalistic light that *l-cry* mutants are less able to discriminate between the different types of light (Poehn et al., 2022; Zurl et al., 2022).

ACKNOWLEDGMENTS

We are grateful to Andrij Belokurov, Margaryta Borysova, and Netsanet Getachew for routine worm cultures and genotyping support, as well as all members of the Tessmar-Raible and Raible labs for continuous constructive discussions. We further thank the team of the Research Station Haidhof, where field tank experiments were conducted, as well as Dunja Rokvic, Marta Beauchamp, Petra Schaffer, Pedro O. Brum, and Netsanet Getachew for help with sampling and sample preparations. This work was supported by Helmholtz Society, Distinguished Professorship by the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, H2020 European Research Council, ERC Grant Agreement #819952, Austrian Science Funds (FWF) SFB F78 and HFSP RGP021/2024 to KT-R. None of the funding bodies were involved in the design of the study, the collection, analysis, and interpretation of data, or in writing the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

Conceptualization: POW, NSH, KT-R. Methodology: NSH, KH, KT-R. Investigation: POW, NSH, KH, KT-R. Funding acquisition: KT-R. Project administration: KT-R. Supervision: NSH, KT-R. Writing – original draft: POW, KT-R. Writing – review and editing: POW, NSH, KH, KT-R.

REFERENCES

Andreatta G, Tessmar-Raible K (2020) The still dark side of the moon: Molecular mechanisms of lunar-controlled rhythms and clocks. *J Mol Biol* 432: 3525–3546

- Aschoff J (1960) Exogenous and endogenous components in circadian rhythms. *Cold Spring Harb Symp Quant Biol* 25: 11–28
- Aschoff J (1965) Circadian rhythms in man. *Science* 148: 1427–1432
- Ayers T, Tsukamoto H, Gühmann M, Tessmar-Raible K (2018) A G_o-type opsin mediates the shadow reflex in the annelid *Platynereis dumerilii*. *BMC Biol* 16: 41
- Bollati E, Lyndby NH, D'Angelo C, Köhl M, Wiedenmann J, Wangpraseurt D (2022) Green fluorescent protein-like pigments optimise the internal light environment in symbiotic reef-building corals. *eLife* 11: e73521
- Carothers JH, Jakšić FM (1984) Time as a niche difference: the role of interference competition. *Oikos* 42: 403–406
- Davies TW, Duffy JP, Bennie J, Gaston KJ (2014) The nature, extent, and ecological implications of marine light pollution. *Front Ecol Environ* 12: 347–355
- Dekens MP, Foulkes NS, Tessmar-Raible K (2017) Instrument design and protocol for the study of light controlled processes in aquatic organisms, and its application to examine the effect of infrared light on zebrafish. *PLOS ONE* 12: e0172038
- Depauw FA, Rogato A, Ribera d'Alcalá M, Falcatore A (2012) Exploring the molecular basis of responses to light in marine diatoms. *J Exp Bot* 63: 1575–1591
- Fischer RM, Fontinha BM, Kirchmaier S, Steger J, Bloch S, Inoue D, et al. (2013) Co-expression of VAL- and TMT-opsins uncovers ancient photosensory interneurons and motoneurons in the vertebrate brain. *PLoS Biol* 11: e1001585
- Fontinha BM, Zekoll T, Al-Rawi M, Gallach M, Reithofer F, Barker AJ, et al. (2021) TMT-Opsins differentially modulate medaka brain function in a context-dependent manner. *PLoS Biol* 19: e3001012
- Foster JJ, Kirwan JD, el Jundi B, Smolka J, Khaldy L, Baird E, et al. (2019) Orienting to polarized light at night – matching lunar sky-light to performance in a nocturnal beetle. *J Exp Biol* 222: jeb188532
- Galparsoro I, Borja Á, Germán Rodríguez J, Muxika I, Pascual M, Legorburu I (2012) 35 - Rocky reef and sedimentary habitats within the continental shelf of the Southeastern Bay of Biscay. In “Seafloor Geomorphology as Benthic Habitat” Ed by PT Harris, EK Baker, Elsevier, London, pp 493–507
- Gambi MC, Lorenti M, Russo GF, Scipione MB, Zupo V (1992) Depth and seasonal distribution of some groups of the vagile fauna of the *Posidonia oceanica* leaf stratum: Structural and trophic analyses. *Mar Ecol* 13: 17–39
- Gehring W, Rosbash M (2003) The coevolution of blue-light photoreception and circadian rhythms. *J Mol Evol* 57: S286–S289
- Gerkema MP, Davies WI, Foster RG, Menaker M, Hut RA (2013) The nocturnal bottleneck and the evolution of activity patterns in mammals. *Proc Royal Soc B* 280: 20130508
- Häfker NS, Andreatta G, Manzotti A, Falcatore A, Raible F, Tessmar-Raible K (2022a) Rhythms and clocks in marine organisms. *Annu Rev Mar Sci* 15: 509–538
- Häfker NS, Connan-McGinty S, Hobbs L, McKee D, Cohen JH, Last KS (2022b) Animal behavior is central in shaping the realized diel light niche. *Commun Biol* 5: 562
- Häfker NS, Holcik L, Mat AM, Coric A, Vadiwala K, Beets I, et al. (2024) Molecular circadian rhythms are robust in marine annelids lacking rhythmic behavior. *PLoS Biol* 22: e3002572
- Hensley NM, Rivers TJ, Gerrish GA, Saha R, Oakley TH (2023) Collective synchrony of mating signals modulated by ecological cues and social signals in bioluminescent sea fireflies. *Proc Royal Soc B* 290: 20232311
- Hut RA, Kronfeld-Schor N, van der Vinne V, De la Iglesia H (2012) In search of a temporal niche: Environmental factors. In “The Neurobiology of Circadian Timing. Progress in Brain Research, Vol. 199” Ed by A Kalsbeek, M Mrosovsky, T Roenneberg, RG Foster, Elsevier, London, pp 281–304

- Katana R, Guan C, Zanini D, Larsen ME, Giraldo D, Geurten BRH, et al. (2019) Chromophore-independent roles of opsin apoproteins in *Drosophila* mechanoreceptors. *Curr Biol* 29: 2961–2969.e4
- Kirwan JD, Li T, Ullrich-Lüter J, La Camera G, Nilsson D-E, Arnone MI (2024) The sea urchin *Paracentrotus lividus* orients to visual stimuli. Preprint at <https://www.biorxiv.org/content/10.1101/2024.01.05.574409v1>
- Kronfeld-Schor N, Dominoni D, de la Iglesia H, Levy O, Herzog ED, Dayan T, et al. (2013) Chronobiology by moonlight. *Proc R Soc B* 280: 20123088
- Kronfeld-Schor N, Visser ME, Salis L, van Gils JA (2017) Chronobiology of interspecific interactions in a changing world. *Phil Trans R Soc B* 372: 20160248
- Last Kim S, Hobbs L, Berge J, Brierley Andrew S, Cottier F (2016) Moonlight drives ocean-scale mass vertical migration of zooplankton during the Arctic winter. *Curr Biol* 26: 244–251
- Liu L, Cong X, Wang X, Chen M (2025) Feeding and growth are regulated by light and opsins-mediated phototransduction in sea cucumber *Apostichopus japonicus*. *Aquaculture* 594: 741410
- Mat A, Vu HH, Wolf E, Tessmar-Raible K (2024) All light, everywhere? Photoreceptors at nonconventional sites. *Physiology* 39: 30–43
- Matsuo R, Koyanagi M, Sugihara T, Shirata T, Nagata T, Inoue K, et al. (2023) Functional characterization of four opsins and two G alpha subtypes co-expressed in the molluscan rhabdomeric photoreceptor. *BMC Biol* 21: 291
- Menchetti M, Guéguen M, Talavera G (2019) Spatio-temporal ecological niche modelling of multigenerational insect migrations. *Proc R Soc B* 286: 20191583
- Michael AK, Fribourgh JL, Van Gelder RN, Partch CL (2017) Animal cryptochromes: Divergent roles in light perception, circadian timekeeping and beyond. *Photochem Photobiol* 93: 128–140
- Milivojevic N, Velastegui Gamboa CL, Andreatta G, Raible F, Tessmar-Raible K (2024) A c-opsin functions in a ciliary-marginal zone-like stem cell region of an invertebrate camera-type eye. Preprint at <https://www.biorxiv.org/content/10.1101/2024.08.19.608633v1>
- Nordtug T, Melø TB (1988) Diurnal variations in natural light conditions at summer time in arctic and subarctic areas in relation to light detection in insects. *Ecography* 11: 202–209
- Oliveri P, Fortunato AE, Petrone L, Ishikawa-Fujiwara T, Kobayashi Y, Todo T, et al. (2014) The cryptochrome/photolyase family in aquatic organisms. *Mar Genom* 14: 23–37
- Özpolat BD, Randel N, Williams EA, Bezares-Calderón LA, Andreatta G, Balavoine G, et al. (2021) The Nereid on the rise: Platynereis as a model system. *EvoDevo* 12: 10
- Poehn B, Tessmar-Raible K (2022) Assessment of circalunar (~monthly) rhythms. In “Circadian Clocks” Ed by T Hirota, M Hatori, S Panda, Springer US, New York, NY, pp 217–230
- Poehn B, Krishnan S, Zurl M, Coric A, Rokvic D, Häfker NS, et al. (2022) A cryptochrome adopts distinct moon- and sunlight states and functions as sun- versus moonlight interpreter in monthly oscillator entrainment. *Nat Commun* 13: 5220
- Posch T, Freyhoff A, Uhlmann T (2010) Das Ende der Nacht: Die globale Lichtverschmutzung und ihre Folgen, 1st Ed. Wiley VCH, Weinheim
- Raible F, Tessmar-Raible K, Arboleda E, Kaller T, Bork P, Arendt D, et al. (2006) Opsins and clusters of sensory G-protein-coupled receptors in the sea urchin genome. *Dev Biol* 300: 461–475
- Revilla IDR, Rajan VB, Waldherr M, Prohaczka G, Musset H, Orel L, et al. (2021) Characterization of cephalic and non-cephalic sensory cell types provides insight into joint photo- and mechanoreceptor evolution. *eLife* 10: e66144
- Ritter A, Tessmar-Raible K (2024) Time me by the moon : The evolution and function of lunar timing systems. *EMBO Rep* 25: 3169–3176
- Sakai K, Yamashita T, Imamoto Y, Shichida Y (2015) Diversity of active states in TMT opsins. *PLOS ONE* 10: e0141238
- Sarfati R, Hayes JC, Peleg O (2021) Self-organization in natural swarms of *Photinus carolinus* synchronous fireflies. *Sci Adv* 7: eabg9259
- Sarfati R, Joshi K, Martin O, Hayes JC, Iyer-Biswas S, Peleg O (2023) Emergent periodicity in the collective synchronous flashing of fireflies. *eLife* 12: e78908
- Senthilan PR, Piepenbrock D, Ovezmyradov G, Nadrowski B, Bechstedt S, Pauls S, et al. (2012) *Drosophila* auditory organ genes and genetic hearing defects. *Cell* 150: 1042–1054
- Somaschini A, Martini N, Gravina MF, Belluscio A, Corsi F, Ardizzone GD (1998) Characterization and cartography of some Mediterranean soft-bottom benthic communities (Ligurian Sea, Italy). *Sci Mar* 62: 27–36
- Speight M, Henderson (2010) Marine Ecology: Concepts and Applications. Wiley-Blackwell, West Sussex
- Tamai TK, Young LC, Whitmore D (2007) Light signaling to the zebrafish circadian clock by cryptochrome 1a. *Proc Natl Acad Sci U S A* 104: 14712–14717
- Tsukamoto H (2014) Diversity and functional properties of bistable photopigments. In “Evolution of Visual and Non-visual Pigments” Ed by DM Hunt, MW Hankins, SP Collin, NJ Marshall. Springer US, Boston, MA, pp 219–239
- Tsukamoto H, Chen IS, Kubo Y, Furutani Y (2017) A ciliary opsin in the brain of a marine annelid zooplankton is ultraviolet-sensitive, and the sensitivity is tuned by a single amino acid residue. *J Biol Chem* 292: 12971–12980
- Veedin Rajan VB, Häfker NS, Arboleda E, Poehn B, Gossenreiter T, Gerrard EJ, et al. (2021) Seasonally relevant UVA light alters neurohormone amounts and behavior via a ciliary opsin in a marine mass spawning annelid. *Nat Ecol Evol* 5: 204–218
- Vinayak P, Coupar J, Hughes SE, Fozdar P, Kilby J, Garren E, et al. (2013) Exquisite light sensitivity of *Drosophila melanogaster* cryptochrome. *PLoS Genet* 9: e1003615
- Vu HH, Behrmann H, Hanic M, Jeyasankar G, Krishnan S, Dannecker D, et al. (2023) A marine cryptochrome with an inverse photo-oligomerization mechanism. *Nat Commun* 14: 6918
- Walker II WH, Meléndez-Fernández OH, Nelson RJ, Reiter RJ (2019) Global climate change and invariable photoperiods: A mismatch that jeopardizes animal fitness. *Ecol Evol* 9: 10044–10054
- Walmsley L, Hanna L, Moulard J, Martial F, West A, Smedley AR, et al. (2015) Colour as a signal for entraining the mammalian circadian clock. *PLoS Biol* 13: e1002127
- Yaguchi J, Yaguchi S (2021) Sea urchin larvae utilize light for regulating the pyloric opening. *BMC Biol* 19: 64
- Zekoll T, Waldherr M, Tessmar-Raible K (2021) Characterization of *tmt-opsin2* in medaka fish provides insight into the interplay of light and temperature for behavioral regulation. *Front Physiol* 12: 726941
- Zhong M, Kawaguchi R, Kassai M, Sun H (2012) Retina, retinol, retinal and the natural history of vitamin A as a light sensor. *Nutrients* 4: 2069–2096
- Zurl M, Poehn B, Rieger D, Krishnan S, Rokvic D, Veedin Rajan VB, et al. (2022) Two light sensors decode moonlight versus sunlight to adjust a plastic circadian/circalunidian clock to moon phase. *Proc Natl Acad Sci U S A* 119: e2115725119

(Received November 10, 2024 / Accepted December 1, 2024 / Published online January 30, 2025)