

cDNA Cloning and Expression of the *Xenopus* Homologue of the Neural Adhesion Molecule, Contactin (F3/F11)

Authors: Nagata, Saburo, Fujita, Naoko, Takeuchi, Kosei, and Watanabe, Kazutada

Source: Zoological Science, 13(6) : 813-820

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.13.813>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

cDNA Cloning and Expression of the *Xenopus* Homologue of the Neural Adhesion Molecule, Contactin (F3/F11)

Saburo Nagata¹, Naoko Fujita¹, Kosei Takeuchi^{2,3},
and Kazutada Watanabe²

¹*Department of Chemical and Biological Sciences, Faculty of Science, Japan Women's University, Mejirodai 2-8-1, Bunkyo-ku, Tokyo 112, Japan*

²*Department of Experimental Biology, Tokyo Metropolitan Institute of Gerontology, Sakaecho 35-2, Itabashiku, Tokyo 173, Japan*

ABSTRACT—Contactin (F3/F11) is an immunoglobulin superfamily cell surface glycoprotein predominantly expressed in the nervous system. To examine the structure and tissue distribution of *Xenopus* contactin, a cDNA clone was isolated based on the amino acid sequences conserved among chicken and mammalian contactin proteins. The conceptual translate of the cDNA consists of 1005 amino acid residues that have 70% identity to those of chicken and mammalian contactin. Northern blot hybridization using a labeled cDNA fragment revealed specific expression of 6.5 kb mRNA in the brain. Monoclonal and polyclonal antibodies prepared to the recombinant *Xenopus* contactin peptides detected a single 135 kD band on Western blots of the brain and spinal cord extracts. Differential extraction and phosphatidylinositol-specific phospholipase C (PI-PLC) digestion experiments showed that the immunoreactive 135 kD proteins bind, at least in part, to the membrane by GPI anchor. On brain tissue sections, strong contactin immunoreactivities were detected on nerve fibers of a subset of cerebral and cerebellar neurons. These results suggest that the basic structure and tissue distribution of *Xenopus* contactin are similar to those in other vertebrates.

INTRODUCTION

Morphogenesis of the nervous system involves sequential cellular interaction that regulates cell differentiation, tissue organization, and formation of the highly stereotyped neuronal network. Recent extensive studies on molecular mechanisms of such cellular interaction have pointed out critical roles of several groups of membrane glycoproteins. The immunoglobulin superfamily is one of the major groups of such membrane glycoproteins. A growing number of immunoglobulin superfamily proteins have been identified in the nervous system of diverse vertebrate and invertebrate species, suggesting conserved structural and functional properties of this superfamily (see reviews, Hynes and Lander, 1992; Sonderegger and Rathjen, 1992; Vaughan and Bjorkman, 1996).

Contactin (also called F3 in mammals and F11 in chickens) is a member of the immunoglobulin superfamily predominantly expressed in the nervous system of chickens (Ranscht, 1988; Brummendorf *et al.*, 1989) and mammals (Gennarini *et al.*, 1989; Reid *et al.*, 1994; Hosoya *et al.*, 1995; Watanabe *et al.*, 1995). The molecules are 135 kD

glycoproteins that belong to the immunoglobulin/fibronectin type III subfamily, having 6 immunoglobulin C2-like domains and 4 fibronectin type III-like domains (Rathjen and Jessel, 1991; Brummendorf and Rathjen, 1993; Vaughn and Bjorkman, 1996). Although chicken contactin was first reported to be a transmembrane protein with a cytoplasmic domain (Ranscht, 1988), the later studies have shown that mammalian and chicken contactin bind to the membrane by a C-terminal glycosylphosphatidylinositol (GPI) anchor. A soluble form of contactin was also reported to occur in the brain and cerebrospinal fluid (Gennarini *et al.*, 1989; Durbec *et al.*, 1992; Moss and White, 1992).

Contactin mediates cell-to-cell adhesion and neurite extension in cultured neural cells by homophilic interaction (Gennarini *et al.*, 1991). In addition, it heterophilically interacts with various membrane glycoproteins such as Ng-CAM (Brummendorf *et al.*, 1993), Nr-CAM/Bravo (Morales *et al.*, 1993) and receptor tyrosine phosphatase β (Peles *et al.*, 1995), which are expressed on neurons or neuroglial cells. These heterophilic interactions also cause neural cell adhesion and neurite outgrowth in vitro. In addition, contactin mediates molecular recognition of extracellular matrix glycoproteins such as restrictin/janusin and tenascin, which leads to adhesion of neural cells or repulsion of growing neurites (Rathjen *et al.*, 1991; Norenberg *et al.*, 1992; Pesheva *et al.*, 1993; Zisch *et al.*, 1992; Vaughan *et al.*, 1994). These homophilic and

³ Present address: Graduate School of Biological Sciences, Nara Institute of Science and Technology, Takayamacho 8916-5, Ikoma 630-01, Japan.

heterophilic interactions with various molecules are likely to represent functional diversity of the domains of this molecule. In mouse and rat brains, contactin proteins distribute preferentially on the axonal membrane and the amounts of its expression markedly increase after birth (Gennarini *et al.*, 1990; Faivre-Sarrailh *et al.*, 1992; Yoshihara *et al.*, 1995; Watanabe *et al.*, 1995). All these findings seem to imply that contactin may have major roles in cell-to-cell and cell-to-extracellular matrix interactions of post-mitotic neurons which regulate neurite outgrowth, axonal path finding, fasciculation, myelination and synaptogenesis. The detailed molecular mechanisms and the physiological roles of such contactin's actions are, however, remained to be determined in the developing nervous system.

We have introduced the South African clawed frog *Xenopus laevis* as a model to study roles of contactin in the developing nervous system, since this animal provides advantageous experimental approaches to investigate gene expression and cellular interactions in the vertebrate embryonic development. Thus, the pattern and the inductive tissue interactions in early neurogenesis were described in detail, and many genes, some of which are vertebrate homologue of the *Drosophila* neurogenic genes, are defined to regulate these processes (Lee *et al.*, 1995). We report here cloning of the cDNA encoding the *Xenopus* homologue of contactin and expression of its mRNA and proteins in brains of young adult frogs. Besides a well-known immunoglobulin superfamily member, NCAM (Kintner, 1988; Kreig *et al.*, 1989; Tonissen and Kreig, 1993), this is, to our knowledge, the first study to report the structure and tissue distribution of the neural adhesion molecule of this superfamily in amphibia.

MATERIALS AND METHODS

Animals

Six to 10 month-old J strain *Xenopus laevis* (Tochinai and Katagiri, 1975) were used for cDNA library construction and Southern blot analysis. In the other experiments outbred or J strain frogs at ages of 6-12 months were used.

Isolation and characterization of the contactin cDNA

To amplify cDNA fragments encoding the *Xenopus* homologue of contactin by PCR, a set of degenerate oligonucleotide primers was designed from the amino acid sequences conserved among chicken (Brümmendorf *et al.*, 1989), mouse (Gennarini *et al.*, 1989), rat (Hosoya *et al.*, 1995), bovine (Watanabe *et al.*, 1995) and human (Reid *et al.*, 1994) contactin proteins: (5')-TGGGTNGARCAYAT-HAATGA-(3') for amino acid residues 323-329 of chicken contactin, WVEHND and (5')-CKRAAYTCRTAYTCCATCCA-(3') for amino acid residues 664-670, WMEYEFR. The total RNA was isolated from frog brains as described (Chomczynski and Sacchi, 1987), and the template cDNA was synthesized using AMV reverse transcriptase according to the manufacturer's instruction (Takara shuzo Co., Ohtu, Japan). Using these template and primers, a DNA fragment of about 1 kb, which was the size expected from the contactin cDNA sequences, was amplified and cloned in pBluescript. The partial sequence of this fragment showed about 70% nucleotide identity to the chicken sequence. The cloned fragment was labeled with digoxigenin through PCR amplification (Boehringer Mannheim, Mannheim, Germany) and used as the screening probe. Poly(A)⁺ RNA was isolated from frog

brains using the FastTrack mRNA isolation kit (Invitrogen, San Diego, CA), and an oligo (dT)-primed cDNA library in the λ ZapII vector (Stratagene, La Jolla, CA) was constructed. After screening approximately 2×10^6 recombinant plaques, 21 hybridizing clones were isolated and converted from recombinant phages to plasmids. One of the clones containing an entire long open reading frame was used for further characterization. Nucleotide sequences were determined by the dideoxy chain termination method using fluorescent primers and an automated DNA sequencer (Pharmacia Biotech, Uppsala, Sweden).

Southern and Northern blotting

Southern blot analysis was performed according to the previously described method (Kreig *et al.*, 1989). The genomic DNA was isolated from peripheral blood cells, digested with restriction endonucleases, fractionated by electrophoresis on a 0.8% agarose gel and transferred to the nitrocellulose membrane. The probes used were ³²P-labeled cDNA fragment a and b shown in Fig. 1A. For Northern blot analysis, total RNA preparations from various frog tissues were fractionated on a 1.2% agarose gel containing formaldehyde. The blots on nitrocellulose membranes were hybridized with the labeled cDNA fragment. The hybridized Southern and Northern blots were autoradiographed using X-ray film for several days at -70°C.

Antibodies and Western blotting

The cDNA fragments c and d were amplified by PCR and cloned into the bacterial expression vectors, pET-3b (Novagen Inc., Madison, WI) and pGEX-3X (Pharmacia Biotech), respectively. The transformed bacteria were induced to produce the recombinant peptides according to the instruction of the vector providers. On SDS-polyacrylamide gel electrophoresis (SDS-PAGE) of the bacterial lysate, the recombinant peptide of amino acid residues 18-292 (peptide I) of the *Xenopus* contactin homologue (cDNA fragment c in Fig. 1A) was identified as a 33 kD band and the GST-fusion peptide containing amino acid residues 319-665 (peptide II) as a 66 kD band. To purify recombinant peptides, the bacterial inclusion bodies were isolated (Sambrook *et al.*, 1989), fractionated by SDS-PAGE and the peptides were eluted from the gel by using the Electroeluter (BioRad Laboratories, Hercules, CA).

New Zealand white rabbits were injected subcutaneously with 500 μ g of purified GST-peptide II in the complete Freund adjuvant, followed by 2 monthly injections in the incomplete adjuvant. Antibody production was monitored by Western blotting and the high titer antisera were harvested. The rabbit antiserum to the recombinant rat contactin (F3) peptide was produced similarly (Takeuchi *et al.*, unpublished). The sera were affinity-purified using an antigen-bound nitrocellulose membrane as an adsorbent according to the described method (Harlow and Lane, 1988). Hybridoma cell lines secreting monoclonal antibodies to the recombinant peptide I (1H1; subclass IgG1(κ)) and peptide II (5C9; IgG1(κ)) were produced according to the established method (Harlow and Lane, 1988). Hybridomas were injected into the Balb/c mice and the tumor ascites containing monoclonal antibodies were harvested.

Western blotting was performed as described previously (Watanabe *et al.*, 1995) and the blots were developed using ECL Western Blotting Detection Reagent (Amersham Life Science, Inc., Little Chalfont, England).

Brain membrane fractions and PI-PLC digestion

Preparation and PI-PLC digestion of the brain membrane fraction were performed essentially as described previously (Gennarini *et al.*, 1989). In brief, the membrane fraction was resuspended at a concentration of 5 mg protein/ml in 50 mM Tris-HCl containing protease inhibitors. One hundred μ l of the membrane fraction were incubated with 0.5 U of PI-PLC from *B. cereus* (Boehringer Mannheim) for 40 min at 37°C, freeze-and-thawed once, and then reincubated with the fresh enzyme. The controls were incubated in the buffer with or without

0.5% NP-40. After incubation, the samples were centrifuged at 140 k ×g for 90 min to separate the pelleted membrane and the supernatant, and the membrane was washed twice with the buffer. Contactin in the membrane and the supernatant was detected by Western blotting.

Immunocytochemistry

Frogs were anesthetized in 0.2% MS-222 (ethyl 3-aminobenzoate methane sulfonic acid; Aldrich Chem. Co., Milwaukee, WI), and perfused with PBS and then with 4% paraformaldehyde in PBS. After perfusion, the brains were dissected and immersed in the fixative overnight. They were washed with PBS, immersed in 30% sucrose, and embedded in the OCT compound (Miles Inc., Elkhart, IL). Frozen sections of about 10 μm thick were cut using a cryostat. The sections on glass slides were successively incubated with the affinity-purified rabbit antiserum (dilution 1:2000) and Vectastain ABC reagents (Vector Lab., Burlingame, CA). Control sections were incubated with a normal rabbit serum or the antiserum preabsorbed with either the recombinant GST-contactin peptide II or GST.

RESULTS

Structure of *Xenopus contactin*

The isolated cDNA clone is about 6.5 kb in size and contains an open reading frame of 3015 nucleotides. The conceptual translate consists of 1005 amino acids (Fig. 1B). Examination of the translate demonstrates that the N- and C-terminal regions contain stretches of hydrophobic amino acid sequences (amino acid residues 1-17 and 972-1005) which are a typical signal peptide and a GPI-anchorage motif, respectively. The mature peptide therefore consists of 988 amino acid residues, having a molecular weight of 109517 D.

Alignment of the translate with the chicken and mammalian proteins reveals great similarities between them, with identical amino acids at about 70% of the residues (Fig. 1B). However, amino acid residues in the N-terminal signal sequence and the C-terminal GPI anchor motif are not conserved well among the species. The signal sequence in the *Xenopus* peptides is 17 amino acid residues in contrast to




Fig. 1. (A) Restriction map of the cDNA encoding *Xenopus contactin*. The coding and non-coding sequences are represented by shaded and open boxes, respectively. The sites are: A, *Apa* I; B, *Bam* HI; E, *Eco* RI; K, *Kpn* I; N, *Nde* I; X, *Xho* I. Lines a and b represent fragments used as probes for Northern and Southern hybridization, and lines c and d for construction of the expression vectors (see text). (B) Amino acid sequence comparison of *Xenopus contactin* (X) with the chicken homologue (C). Dots and dashed lines in the chicken sequence indicate identical amino acid residues and gaps, respectively. Cysteine residues marked with closed circles are conserved to form immunoglobulin-like intrapeptide disulfide bonds, and asparagine residues marked with open squares potential N-linked glycosylation sites. The asparagine residue 616 (closed triangle) in the *Xenopus* sequence is not a glycosylation site because of the substitution of S to Q at the residue 618 (asterisk). Boxed sequences at N- and C-terminal regions indicate putative signals for secretion and GPI anchorage, respectively. The cDNA sequence data reported here will appear in the DDBJ, EMBL and GenBank Nucleotide Sequence Databases with the accession No. D86505.

24-26 residues in chicken and mammals. The 12 cysteine residues, which can form intrapeptide disulfide bonds of 6 immunoglobulin type C2-like domains, are conserved in all the species. In addition, 4 fibronectin type III-like repeats contain conserved characteristic tryptophan residues (residues 600, 713, 815, and 911) and tyrosine/phenylalanine residues (residues 613, 663, 728, 766, 826, 863, 926 and 958). Chicken and mammalian contactin peptides have 9 possible N-glycosylation sites (NXT/S). The *Xenopus* peptide conserves 8 of them. However, one of the sites (N at amino acid residue 616) does not have the consensus sequence by substitution of S to Q (618). There is an insertion of 3 amino acid residues in the second immunoglobulin C2 domain as compared with the chicken translate (or 2 residue insertion over the mammalian translates).

Contactin sequences in the *Xenopus* genome

Xenopus laevis is known to be pseudotetraploid (Kobel and Du Pasquier, 1986) and so has two functional non-allelic loci for many genes. To estimate the copy number of the contactin genes in the *Xenopus* genome, Southern blot analyses were performed using two separate cDNA fragments as probes. Both of the probe hybridized to two bands or a single broad band in each digest of 5 restriction enzymes (Fig. 2), suggesting that *Xenopus* has two non-allelic loci for the contactin gene as for the NCAM genes (Tonissen and Kreig, 1993).

Expression of the contactin mRNA and proteins

Northern blot hybridization using the radio-labeled cDNA fragment as a probe detected in the brain a single 6.5 kb band, which is approximately the same in size to the isolated cDNA (Fig. 3). No band was found in the other tissues examined even after an extended exposure period. The previous papers

describe expression of a major 6.5 kb and minor 3.2-4.4 kb contactin mRNAs in chicken and mammalian brains (Ranscht, 1988; Gennarini *et al.*, 1989; Reid *et al.*, 1994), suggesting alternative polyadenylation of the transcript. Our effort was unsuccessful to detect such a shorter mRNA in *Xenopus* brains at various developmental stages (data not shown).

All the monoclonal and polyclonal antibodies detected a single 135 kD band on the Western blot of brain extracts (Fig. 4A-C). A similar 135 kD band was also detected in the spinal chord extract. However, as in above-described Northern blot analysis, no band was found on the blots of other tissues examined (data not shown). The rabbit antiserum prepared against rat contactin also detected a similar 135 kD band in the brain (Fig. 4D). From the difference between the molecular weight estimated by Western blotting and that deduced from the cDNA sequence, *Xenopus* contactin is, similar to mammalian and chicken contactin proteins, probably highly glycosylated.

The differential extraction and PI-PLC digestion experiments were performed to assess subcellular localization of contactin proteins in *Xenopus* brains (Fig. 5). A majority of immunoreactive contactin proteins was recovered in the membrane fraction and a part of them was solubilized from the membrane in the buffer containing Nonidet P-40. A smaller amount of contactin proteins was also detectable in Nonidet P-40-insoluble fraction, which may represent a "membrane skeleton"-associated form (Moss and White, 1992). The PI-PLC treatment clearly increased the release of contactin proteins from the membrane fraction as compared with the control incubated without PI-PLC. This implies that in *Xenopus* brains contactin molecules bind, at least in part, to the membrane by GPI anchor.




Fig. 2. Southern blot analysis of contactin sequences in *Xenopus* genomic DNA. Genomic DNA isolated from blood cells were digested with *Bam* HI (Bm), *Bgl* II (Bg), *Eco* RI (E), *Hind* III (H) and *Pst* I (P), separated by agarose gel electrophoresis and transferred to the nitrocellulose membrane. (A) and (B) were probed with labeled cDNA fragment a and b in Fig. 1A, respectively.




Fig. 3. Northern blot analysis of contactin expression in various *Xenopus* tissues. Each lane was loaded with 5 μ g of total RNA and the blot was hybridized with the radiolabeled cDNA probe (upper panel). Expression of the 6.5 kb contactin transcript was detected in the brain (lane 1), but not in the other tissues: lanes 2-9, heart; kidney; liver; lung; skeletal muscle; spleen; testis; thymus. The same membrane was stained with methylene blue as a control for RNA loading (lower panel).




Fig. 4. (A-C) Western blot analysis of contactin proteins in *Xenopus* brains. The *Xenopus* brain extracts were fractionated by SDS-PAGE, transferred to nitrocellulose membranes, and probed with an affinity-purified rabbit antiserum (A), monoclonal antibodies 5C9 (B) and 1H1 (C) prepared against recombinant *Xenopus* contactin peptides. (D) Western blot analysis of contactin proteins in rat (lane 1) and *Xenopus* (lane 2) brain extracts using the rabbit anti-rat contactin antiserum.




Fig. 5. PI-PLC digestion of the membrane fraction of *Xenopus* brains. The membrane fraction was incubated with NP-40 (NP), PI-PLC (PLC) or the buffer alone (C), and the supernatants (S) and the precipitates (P) were separated by centrifugation. Contactin in each fraction was detected by Western blotting using monoclonal antibody 5C9.

Tissue distribution of contactin in the brain

To examine tissue distribution of contactin proteins in the brain, frozen sections were immunocytochemically stained using the affinity-purified rabbit antiserum to recombinant GST-contactin peptides. Figure 6 shows a part of typical transverse sections through the cerebellum and the brain stem which are not clearly demarcated. Similar to other frog species (Kemali and Braitenberg, 1969; Sotelo, 1976), the layered arrangement of *Xenopus* cerebellum differs from that of the mammalian cerebellum, consisting of the dorsal molecular layer, rather diffuse Purkinje cell layer and granular layer intermingled with white matter. No prominent contactin immunoreactivity was detected in the Purkinje cell layer and the most part of the molecular layer (Fig. 6A, C). Sparsely-distributed immunoreactive fibers were found in the dorsal midline area of the molecular layer. Accumulation of the immunoreactive fibers was found in the granular layer and brain stem. In the brain stem, the strongly immunoreactive fibers are accumulated in the medulla.

Weak staining was occasionally found on the neuronal cell bodies. The immunoreactivities on the nerve fibers were eliminated by preabsorption of the antiserum with GST-contactin peptides (Fig. 6B), while the weak staining on neuronal cell bodies remained after the preabsorption. Similar staining of the neuronal cell bodies was obtained using a normal rabbit serum (data not shown), indicating the staining to be nonspecific. The preabsorption of the antiserum with GST did not affect the staining.

The contactin immunoreactivity was also found on the nerve fibers in the optic tectum (Fig. 6D), white matter of the medulla oblongata and cerebral cortex and medulla (data not shown).

DISCUSSION

The results described above show that *Xenopus* contactin is a highly glycosylated 135 kD protein that has 6 immunoglobulin C2-like domains and 4 fibronectin type III-like repeats. It is also shown from the cDNA sequence and the PI-PLC digestion experiment that the proteins are, at least in part, associated with the membrane by GPI anchor. The




Fig. 6. Distribution of contactin immunoreactivities in *Xenopus* brain sections. (A) The frontal section through the cerebellum which is stained with an affinity-purified rabbit antiserum to *Xenopus* contactin. (B) The control section at the similar portion to A incubated with the antiserum preabsorbed with the GST-contactin peptide. (C) An enlargement of a parts of the figure A. (D) Distribution of immunoreactive fibers in the dorso-lateral portion of the optic tectum. bs, brain stem; gl, granular layer; ml, molecular layer; pl, Purkinje cell layer. Bars represent 100 μ m.

mRNA and proteins are preferentially expressed in the brain. The immunocytochemical analysis using an affinity purified antiserum localizes contactin proteins to the fibers of a subset of cerebral and cerebellar neurons. In cerebellum, the contactin-positive fibers are densely accumulated in the

granular layer, whereas they are sparsely distributed in the molecular layer and virtually absent in the Purkinje cell layer. Although the present study does not define the types of neurons that extend contactin-positive fibers, these observations are consistent with the report that contactin expression

in the mouse cerebellum is restricted to mossy fibers, granule cells and Golgi neurons, while Purkinje, stellate and basket cells and non-neuronal cells are contactin negative (Faivre-Sarraiilh *et al.*, 1992). Thus, the structural characteristics and tissue distribution of *Xenopus* contactin are basically similar to those of previously reported chicken and mammalian contactin.

Contactin is thought to be involved in cell-to-cell and cell-to-extracellular matrix interactions that regulate neurite outgrowth, axonal path finding, fasciculation and synaptogenesis. In chicken and mammalian contactins, these interactions are shown to be based on homophilic and heterophilic recognition of several ligands by specific domains of the contactin molecule (Gennarini *et al.*, 1991; Brümmendorf *et al.*, 1993; Morales *et al.*, 1993; Peles *et al.*, 1995). From the identical domain structures and the highly conserved amino acid sequences in each domain, similar molecules may serve as ligands for *Xenopus* contactin.

Contactin binds to the membrane phospholipid by GPI anchor and has neither membrane spanning nor cytoplasmic domains. It is unknown how the ligand-binding signal can be transduced to the interior of the cell. The recent papers suggest direct or indirect non-covalent association of contactin proteins with the intracellular Fyn tyrosine kinase (Olive *et al.*, 1995; Cervoello *et al.*, 1996). This may pose an important possibility that this tyrosine kinase is an intracellular component of contactin signaling. More detailed studies are obviously required to characterize molecular components and their interaction which transduce contactin signaling. A soluble form of contactin is reported to occur in the brain and cerebro-spinal fluid (Gennarini *et al.*, 1991; Durbec *et al.*, 1992). These proteins have been shown to induce neurite outgrowth in cultured neurons, suggesting their roles as growth or trophic factors in the brain. However, it is unknown how the GPI-anchored contactin proteins are released from the membrane (Moss and White, 1992). An alternative possibility that the secreted form of contactin is produced from the alternatively-spliced transcript has not been substantiated by cDNA cloning studies including the present one.

As assumed from the functional activities, contactin is thought to play a major role in postmitotic neurons after their migration and settlement in the specific sites of the nervous system. In fact, in mouse and rat brains contactin expression markedly increases after birth (Gennarini *et al.*, 1990; Faivre-Sarraiilh *et al.*, 1992; Watanabe *et al.*, 1995). This makes sharp contrast with other immunoglobulin superfamily cell surface molecules such as NCAM. In *Xenopus* and chicken embryos, NCAM proteins are expressed in the cells of the neural plate and play an important role in early neural morphogenesis (Kintner, 1988). However, little is known about mechanisms regulating cell type- and developmental stage-specific expression of the contactin gene. The structure of the contactin gene in the mouse genome was reported recently (Buttiglione *et al.*, 1995). This study pointed out possible binding sites for homeodomain- and bHLH-containing transcription factors in the 5' regulatory region, postulating that contactin gene

expression is regulated under a global developmental program.

Our preliminary study shows that contactin gene transcripts become detectable in *Xenopus* embryos before the closure of the neural tube (Nagata *et al.*, unpublished data), which is much earlier than the stages previously reported in other systems (Faivre-Sarraiilh *et al.*, 1992; Vaughan *et al.*, 1994; Hosoya *et al.*, 1995; Watanabe *et al.*, 1995). The *in situ* hybridization experiment to locate the transcripts in embryos is under way. Furthermore, the contactin protein is immunocytochemically detectable in the restricted areas of embryonic brain and spinal cord (Nagata *et al.*, unpublished data). Thus, *Xenopus* may provide experimental approaches to examine the mechanisms of developmentally regulated contactin gene expression and its roles in the development of the vertebrate nervous system.

ACKNOWLEDGMENT

This study was supported by a Grant-in-Aid for Scientific Research (No. 08680801) from the Ministry of Education, Science, Sports and Culture of Japan.

REFERENCES

- Brümmendorf T, Hubert M, Treubert U, Leuschner R, Tárnok A, Rathjen FG (1993) The axonal recognition molecule F11 is a multifunctional protein: Specific domains mediate interactions with Ng-CAM and restrictin. *Neuron* 10: 711–727
- Brümmendorf T, Rathjen FG (1993) Axonal glycoproteins with immunoglobulin- and fibronectin type III-related domains in vertebrates: structural features, binding activities, and signal transduction. *J Neurochem* 61: 1207–1219
- Brümmendorf T, Wolff JM, Frank R, Rathjen FG (1989) Neural cell adhesion molecule F11, homology with fibronectin type III and immunoglobulin type C domains. *Neuron* 2: 1351–1361
- Buttiglione M, Cangiano G, Goridis C, Gennarini G (1995) Characterization of the 5' and promoter regions of the gene encoding the mouse neuronal cell adhesion molecule F3. *Mol Brain Res* 29: 297–309
- Cervoello M, Matrangola V, Durbec P, Rougon G, Gomez S (1996) The GPI-anchored adhesion molecule F3 induces tyrosine phosphorylation: involvement of the FNIII repeats. *J Cell Sci* 109: 699–704
- Chomczynski P, Sacchi N (1987) Single step method of RNA isolation by acid guanidinium thiocyanate-phenol chloroform extraction. *Anal Biochem* 162: 156–159
- Durbec P, Gennarini G, Goridis C, Rougon G (1992) A soluble form of the F3 neuronal cell adhesion molecule promotes neurite outgrowth. *J Cell Biol* 117: 877–887
- Faivre-Sarraiilh C, Gennarini G, Goridis C, Rougon G (1992) F3/F11 cell surface molecule expression in the developing mouse cerebellum is polarized at synaptic sites and within granule cells. *J Neurosci* 12: 257–267
- Gennarini G, Cibelli G, Rougon G, Mattei MG, Goridis C (1989) The mouse neuronal cell surface protein F3: A phosphatidylinositol-anchored member of the immunoglobulin superfamily related to chicken contactin. *J Cell Biol* 109: 775–788
- Gennarini G, Durbec P, Boned A, Rougon G, Goridis C (1991) Transfected F3/F11 neural cell surface protein mediates intercellular adhesion and promotes neurite outgrowth. *Neuron* 6: 595–606
- Gennarini G, Rougon G, Goridis C (1990) F3: a new developmentally regulated member of the HNK-1 family. *Acta Histochem* 38

- (Suppl): 65–69
- Harlow E, Lane D (1988) *Antibodies: A Laboratory Manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY
- Hosoya H, Shimada K, Kobayashi S, Takahashi H, Shirasawa T, Takenawa T, Watanabe K (1995) Developmental expression of the neural adhesion molecule F3 in the rat brain. *Neurosci Lett* 186: 83–86
- Hynes RO, Lander AD (1992) Contact and adhesive specificities in the associations, migrations, and targeting of cells and axons. *Cell* 68: 303–322
- Kemali M, Braitenberg V (1969) *Atlas of the Frog's Brain*. Springer-Verlag, Berlin-Heidelberg-New York
- Kobel HR, Du Pasquier L (1986) Genetics of polyploid *Xenopus*. *Trends Genet* 2: 310–315
- Kintner C (1988) Effects of altered expression of the neural cell adhesion molecule, NCAM, on early neural development of *Xenopus* embryos. *Neuron* 1: 545–555
- Kreig PA, Sakaguchi DS, Kintner CR (1989) Primary structure and developmental expression of a large cytoplasmic domain form of *Xenopus laevis* neural cell adhesion molecule (NCAM). *Nucleic Acids Res* 17: 10321–10335
- Lee JE, Hollenberg SM, Snider L, Turner DL, Lipnick N, Weintraub H (1995) Conversion of *Xenopus* ectoderm into neurons by NeuroD, a basic helix-loop-helix protein. *Science* 268: 836–844
- Morales G, Hubert M, Brümmendorf T, Treubertm U, Tárnok A, Schwarz U, Rathjen F G (1993) Induction of axonal growth by heterophilic interactions between the cell surface recognition protein F11 and Nr-CAM/Brabo. *Neuron* 11: 1113–1122
- Moss DJ, White CA (1992) Solubility and posttranslational regulation of GP130/F11-a neuronal GPI-linked cell adhesion molecule enriched in the neuronal membrane skeleton. *Eur J Cell Biol* 57: 59–65
- Norenberg U, Wille H, Wolffe JM, Frank R, Rathjen FG (1992) The chicken neural extracellular matrix molecule restrictin: similarity with EGF-, fibronectin type III-, and fibrinogen-like motifs. *Neuron* 8: 849–863
- Olive S, Dubois C, Schachner M, Rougon G (1995) The F3 neuronal glycosylphosphatidylinositol-linked molecule is localized to glycolipid-enriched membrane subdomains and interacts with L1 and Fyn kinase in cerebellum. *J Neurochem* 65: 2307–2317
- Peles E, Nativ M, Campbell P, Sakurai T, Martinez R, Lev S, Plowman GD, Grumet M, Schlessinger J (1995) The carbonic anhydrase domain of receptor tyrosine phosphatase β is a functional ligand for the axonal cell recognition molecule contactin. *Cell* 82: 251–260
- Pesheva P, Gennarini G, Goridis C, Schachner M (1993) The F3/F11 cell adhesion molecules mediates the repulsion of neurons by the extracellular matrix glycoprotein J1-160/180. *Neuron* 10: 69–82
- Ranscht B (1988) Sequence of contactin, a 130-kD glycoprotein concentrated in areas of interneuronal contact, defines a new member of the immunoglobulin supergene family in the nervous system. *J Cell Biol* 107: 1561–1573
- Rathjen FG, Jessel TM (1991) Glycoproteins that regulate the growth and guidance of vertebrate axons: domains and dynamics of the immunoglobulin/fibronectin type III subfamily. *Semin Neurosci* 3: 297–307
- Rathjen FG, Wolf JM, Chiquet-Ehrismann R (1991) Restrictin: a chick neural extracellular matrix protein involved in cell attachment copurified with the cell recognition molecule F11. *Development* 113: 151–164
- Reid RA, Bronson DD, Young KM, Hemperly JJ (1994) Identification and characterization of the human cell adhesion molecule contactin. *Brain Res Mol Brain Res* 21: 1–8
- Sambrook J, Fritsch EF, Maniatis, T (1989) *Molecular Cloning: A Laboratory Manual*. 2nd Ed, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY
- Sonderegger P, Rathjen FG (1992) Regulation of axonal growth in the vertebrate nervous system by interactions between glycoproteins belonging to two subgroups of the immunoglobulin superfamily. *J Cell Biol* 119: 1387–1394
- Sotelo C (1976) Morphology of the cerebellar cortex. In "Frog Neurobiology, A Handbook" Ed by Linás R, Precht W, Springer-Verlag, Berlin-Heidelberg-New York, pp 864–891
- Tochinai S and Katagiri Ch (1975) Complete abrogation of immune responses to skin allografts and rabbit erythrocytes in the early-thymectomized *Xenopus*. *Dev Growth Differ* 17: 383–394
- Tonissen KF, Kreig PA (1993) Two neural-cell adhesion molecule (NCAM)-encoding genes in *Xenopus laevis* are expressed during development and in adult tissues. *Gene* 127: 243–247
- Vaughan L, Weber P, D'Alessandri L, Winterhalter AH (1994) Tenascin-contactin/F11 interactions: a clue for a developmental role? *Perspect Dev Neurobiol* 2: 43–52
- Vaughn DE, Bjorkman PJ (1996) The (Greek) key to structures of neural adhesion molecules. *Neuron* 16: 261–273
- Watanabe K, Shimada K, Hosoya H, Fukamauchi F, Takenawa T (1995) Cloning of the cDNA encoding neural adhesion molecule F3 from bovine brain. *Gene* 160: 245–248
- Wolff JM, Brümmendorf T, Rathjen FG (1989) Neural cell recognition molecule F11: membrane interaction by covalently attached phosphatidylinositol. *Biochem Biophys Res Comm* 161: 931–938
- Yoshihara Y, Kawasaki M, Tamada A, Nagata S, Kagamiyama H, Mori K (1995) Overlapping and differential expression of BIG-1, BIG-2, TAG-1, and F3: four members of an axon-associated cell adhesion molecule subgroup of the immunoglobulin superfamily. *J Neurobiol* 28: 51–69
- Zisch AH, D'Alessandri L, Ranscht B, Falchetto R, Winterhalter KH, Vaughan L (1992) Neuronal cell adhesion molecule contactin/F11 binds to tenascin via its immunoglobulin-like domains. *J Cell Biol* 119: 203–213

(Received July 15, 1996 / Accepted September 9, 1996)