

# GnRH Analog Stimulates Gonadotropin II Gene Expression in Maturing Sockeye Salmon

Authors: Kitahashi, Takashi, Alok, Deoraj, Ando, Hironori, Kaeriyama, Masahide, Zohar, Yonathan, et al.

Source: Zoological Science, 15(5): 761-765

Published By: Zoological Society of Japan

URL: https://doi.org/10.2108/zsj.15.761

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <a href="http://www.bioone.org/terms-of-use">www.bioone.org/terms-of-use</a>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# GnRH Analog Stimulates Gonadotropin II Gene Expression in Maturing Sockeye Salmon

Takashi Kitahashi<sup>1</sup>\*, Deoraj Alok<sup>2</sup>, Hironori Ando<sup>1</sup>, Masahide Kaeriyama<sup>3</sup>, Yonathan Zohar<sup>2</sup>, Hiroshi Ueda<sup>4</sup> and Akihisa Urano<sup>1</sup>

 <sup>1</sup>Division of Biological Sciences, Graduate School of Science, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan
 <sup>2</sup>Center of Marine Biotechnology, University of Maryland, Biotechnology Institute, Baltimore, Maryland 21202, USA
 <sup>3</sup>National Salmon Resources Center, Sapporo, Hokkaido 062-0922, Japan
 <sup>4</sup>Toya Lake Station for Environmental Biology, Faculty of Fisheries, Hokkaido University, Abuta, Hokkaido 049-5723, Japan

**ABSTRACT**—Our previous study suggested that, in the pituitaries of pre-spawning chum salmon, salmon gonadotropin-releasing hormone (sGnRH) stimulates expression of genes for gonadotropin (GTH) II $\beta$  but not for I $\beta$ , since the levels of mRNAs encoding sGnRH and GTH II but not I were increased during the final stages of spawning migration. In the present study, a capsule of GnRH analog (GnRHa) was implanted into the dorsal muscle of maturing sockeye salmon to clarify function of GnRH on expression of GTH subunit genes in pre-spawning homing salmonids. The amounts of GTH subunit mRNAs in the individual pituitaries were analyzed by a quantitative dot blot analysis using single-stranded sense DNA as the standard. The levels of GTH  $\alpha$  and II $\beta$  mRNAs in the GnRHa-implanted fish were significantly higher than those in the control fish in both the males and females, whereas the levels of GTH I $\beta$  mRNA did not show any significant differences in both sexes. These results indicate that GnRH elevates expression of GTH subunit genes which encode the components of GTH II,  $\alpha$  and II $\beta$  chains, in the pituitary of maturing sockeye salmon, and then accelerates final maturation.

# INTRODUCTION

The presence of two gonadotropins (GTH) in teleosts, namely GTH I and II, was well established by Kawauchi et al. (1989). GTH I is considered as vitellogenic, and GTH II as maturational. They are composed of common  $\alpha$  and unique  $\beta$ subunits. Complete amino acid sequences of GTH  $\alpha$  (Itoh et al., 1990) and  $\beta$  subunits (Itoh et al., 1988) were first determined in chum salmon. Further, complete nucleotide sequences of cDNAs encoding GTH subunits were determined in several salmonids (Trinh et al., 1986; Sekine et al., 1989; Gen et al., 1993; Kato et al., 1993; Suzuki et al., 1995). Because a certain sequence of the GTH subunit mRNA conserves high similarity (about 95%) with their homologues in salmonids, a cDNA probe designed for particular subunit mRNA in one species can be used as a hybridization probe in the pituitaries of various salmonid species (see Hiraoka et al., 1993).

Secretion of GTH is mainly controlled by gonadotropinreleasing hormone (GnRH), which is involved in the control of

\* Corresponding author: Tel. +81-11-706-2995; FAX. +81-11-706-4923.

reproduction in a wide range of vertebrate species including teleosts (see Sherwood et al., 1993). In salmonids, salmon GnRH (sGnRH) functions as reproductive GnRH despite the presence of two GnRH variants, sGnRH and chicken GnRH-II (Amano et al., 1997; Kobayashi et al., 1997). sGnRH neurons in the ventral telencephalon (VT) and the preoptic area (POA) which send their axons to the pars distalis may control pituitary GTH cells to release GTH into circulation. Recent studies in masu salmon indicated that expression of sGnRH gene in these neurons seemed to correlate well with sexual maturation (Amano et al., 1995a, b). Further, these VT and POA sGnRH neurons in chum salmon showed increase in the levels of sGnRH mRNA and also the number of hybridization positive cells during the final stages of spawning migration (Kudo et al., 1996). Arising question here was whether GTH gene expression is controlled by sGnRH in pre-spawning salmon.

In our previous study, we first developed a quantitative dot blot analysis by which the levels of GTH subunit mRNAs in the salmonid pituitary were determined by use of single-stranded sense DNA (ssDNA) encoding the same mRNA sequence. Utilizing this method, we found that the levels of GTH  $\alpha$  and II $\beta$  mRNAs in freshwater chum salmon ready to spawn

were higher than those in coastal seawater animals, however such difference was not seen in the level of GTH I $\beta$  mRNA (Kitahashi *et al.*, 1997). Since, as mentioned above, the number of hybridization positive sGnRH neurons in the VT and POA increased in pre-spawning chum salmon, it is highly probable that sGnRH stimulates expression of genes encoding GTH  $\alpha$  and II $\beta$  but not I $\beta$ , and promotes final sexual maturation in homing salmonids. In the present study, we examined whether this hypothesis is true by implantation of GnRH analog (GnRHa) into the dorsal muscle of maturing sockeye salmon, *Oncorhynchus nerka*. Because of difficulty in obtaining chum salmon at appropriate stages of sexual maturation, we used freshwater sockeye salmon which were maintained for artificial fertilization.

# MATERIALS AND METHODS

#### Fish

Maturing sockeye salmon were caught in the Bibi river upstream to Lake Utonai, Hokkaido, Japan, in the end of June 1996, and were kept in a holding pond in National Salmon Resources Center (Chitose, Hokkaido) under natural photoperiod condition. Fish were tagged by an intraperitoneal pit tag to be discriminated individually. On September 6, experimental animals were randomly selected, anesthetized with 0.05% ethyl m-aminobenzoate methanesulfonate (MS222, Nakalai tesque, Kyoto, Japan) buffered with sodium bicarbonate, and were measured of body weight and fork length. Body weights were about 1.88 kg in the males and 1.18 kg in the females, while fork length were about 55.0 cm in the males and 49.5 cm in the females. Blood samples (1 ml) were collected from the caudal vein for radioimmunoassay of plasma steroid hormones the result of which will be published elsewhere with the observation on improvement of gonadal fecundity (Fukaya et al., in preparation). Afterward, fish received in the dorsal muscle a 2 mm implant capsule containing 75 or 150  $\mu$ g of GnRHa, [Des-Gly<sup>10</sup>, D-Ala<sup>6</sup>, Pro<sup>9</sup>]-GnRH ethyl amide in an ethylene vinyl acetate copolymer matrix (Zohar et al., 1990; Zohar, 1996). The dose of implanted GnRHa was determined by consulting the previous papers (Zohar et al., 1990). Sham operated animals served as controls. Control and GnRHa-implanted fish were sampled three weeks after the implantation. Those fish sampled on the day of implantation were assigned as initial controls. When fish were sampled, gonads were taken out and measured of their weights to calculate gonadosomatic index (GSI). The GSI and the number of animals in each group are shown in Table 1.

 
 Table 1.
 Changes in gonadosomatic index in the initial, control and GnRHa-implanted sockeye salmon

	Male	n	Female	n
Initial	$1.94 \pm 0.11$	3	$7.81 \pm 0.55$	6
Control	$1.22 \pm 0.24^{*}$	5	$13.39 \pm 0.28^{**}$	4
75 μg GnRH	not tested		$12.40 \pm 0.46^{**}$	5
150 μg GnRH	$0.85 \pm 0.13^{**}$	5	$11.94 \pm 0.83^{**}$	3

#### Mean $\pm$ SEM.

Differences to initial: \*, P < 0.05; \*\*, P < 0.01; by the *t*-test.

#### **Tissue preparation**

The pituitaries were taken out upon decapitation, frozen in liquid nitrogen and stored at -80°C. Total RNA was extracted from single pituitaries by the acid guanidium thiocyanate-phenol-chloroform method (Chomczynski and Sacchi, 1987). The pituitary weight was about 20 mg and the amount of total RNA extracted from a single

pituitary was 50-100 µg.

#### Assay method

ssDNAs which have the same sequences of mRNAs encoding GTH  $\alpha$ , I $\beta$  and II $\beta$  were synthesized by use of a polymerase chain reaction (PCR) method with a single common sense primer. Template DNA for PCR was prepared by the restriction enzyme digestion at one site of plasmid DNA containing cDNA insert for the salmon GTH subunit (Sekine *et al.*, 1989). The GTH  $\alpha$  cDNA used in the present study is a cDNA encoding a 2 subunit, which is commonly obtained from GTH I and II according to Itoh's designation (Itoh et al., 1990). Synthetic 20mer oligonucleotide, 5'-GTTACTTCTGCTCTA-AAAGC-3', was used as the PCR primer. The PCR mixture contained 1.6 pmol template DNA described above, PCR buffer (50 mM KCl, 1.5 mM MgCl<sub>2</sub> and 10 mM Tris-HCl, pH 8.3), 200 µM dNTPs, 1.0 µM primer, 1 unit Taq DNA polymerase (TaKaRa) and sterile distilled water to 50 µl, overlaid with 30 µl light mineral oil. The profile of amplification was: 30 cycles of incubation at 93°C for 1.5 min, at 53°C for 1.5 min and 72°C for 2 min after denaturation at 95°C for 4 min. Amplified DNA was subjected to low melting temperature agarose gel electrophoresis to separate the band that contained ssDNA. Afterward, ssDNA was purified by phenol extraction. We obtained ssDNAs for GTH  $\alpha$ , I $\beta$  and II $\beta$  whose lengths were about 780, 790 and 840 bases, respectively.

Labeled cDNA probes which were specific to particular GTH subunit mRNAs were prepared by a primer extension method using Megaprime DNA labeling system (Amersham) and [ $\alpha$ -<sup>32</sup>P]dCTP (Amersham) with synthetic oligonucleotide primers. They are about 200 bases long and specific to corresponding standard DNAs. The sequences used as cDNA probes were carefully selected to avoid cross-hybridization.

Total RNA extracted from the pituitaries was blotted to Hybond<sup>TM</sup>-N<sup>+</sup> membranes (Amersham) using a MilliBlot<sup>TM</sup>-D (Millipore) in duplicate. As the standard to determine the amount of mRNA, serially diluted ssDNA (1-333 amol) were blotted to each membrane in triplicate.

Hybridization with the labeled probe was performed in a solution containing  $5 \times SSPE$  (0.05 M phosphate buffer, pH approx. 7.4, containing 0.76 M NaCl and 5 mM EDTA),  $5 \times Denhardt's$  solution (0.1% solution of bovine serum albumin, Ficoll and polyvinylpyrrolidone), 100 µg/ml denatured calf thymus DNA (Sigma) and  $1.5-2 \times 10^6$  cpm of labeled cDNA probe/ml hybridization buffer at  $65^{\circ}C$  for 20 hr. The membranes were then washed with  $2 \times SSPE/0.1\%$  SDS at room temperature (RT) for 15 min, twice with  $2 \times SSPE/0.1\%$  SDS at  $65^{\circ}C$  for 30 min, and twice with  $0.1 \times SSPE/0.1\%$  SDS at  $65^{\circ}C$  for 30 min. The membranes were then exposed to a Fuji imaging plate (Fuji Photo Film Co., Ltd.) for 24-48 hr. Radioactivity expressed as the intensity of photostimulated luminescence (PSL) was analyzed by a Bioimaging analyzer (Fuji Photo Film Co., Ltd.), and the intensity of signals were estimated by subtraction of background.

#### Statistical analysis

For statistical analysis, Student's or Welch's *t*-test was applied after Scheffé's F test for variance. Correlations between the plasma steroid levels and GTH subunit mRNA levels were analyzed using Pearson's correlation coefficient method.

#### RESULTS

# Changes in gonadosomatic index (GSI)

The stock of sockeye salmon used in the present study usually spawn in the middle to end of October. Hence, the initial controls sampled in the beginning of September were still in immature state. The GSI values in Table 1 indicate that, in both the male and female control groups, gonadal matura-



**Fig. 1.** Changes in the levels of GTH subunit mRNAs in terms of whole pituitary in GnRHa-implanted maturing sockeye salmon. GTH  $\alpha$ , I $\beta$  and II $\beta$  subunit mRNA levels in each group are shown. I, initial control; C, control; 75, 75  $\mu$ g of GnRHa-implanted; 150, 150  $\mu$ g of GnRHa-implanted. Mean  $\pm$  SEM (n = 3-6). \*, *P* < 0.05; \*\*, *P* < 0.01; by the *t*-test.

tion proceeded during the three-week experimental period. However, most of the control females were still preovulatory state, although all the control males excrete a little milt by pressing the abdomen. Implantation of GnRHa accelerated this naturally occurring gonadal maturation. All females which received GnRHa actually ovulated by three week GnRHa treatment (data not shown). The amount of excreted milt was increased in GnRHa treaded males.

# The levels of GTH subunit mRNAs

**Males:** Probably due to progress of sexual maturation, the pituitary levels of GTH  $\alpha$  and II $\beta$  mRNAs but not I $\beta$  mRNA were elevated 2-3 fold in the control males compared to those in the initial control (Fig. 1). The treatment with GnRHa significantly increased these  $\alpha$  and II $\beta$  mRNA levels 2-fold or more, whereas it had little effect on the level of GTH I $\beta$  mRNA (Fig. 1). The pattern of changes in the levels of GTH subunit mRNAs which are expressed in terms of the unit weight of pituitary (mg pituitary), (data not shown) is nearly the same with that in Fig. 1 expressed in terms of whole pituitary.

**Females:** The changes in the levels of GTH subunit mRNAs in the female pituitaries showed similar pattern to those in the male pituitaries (Fig. 1). The levels of GTH  $\alpha$  and II $\beta$  mRNAs were drastically elevated in the GnRHa-implanted females compared to those in the controls. The more than 3-fold increases in the levels of  $\alpha$  and II $\beta$  mRNAs in the GnRHa-implanted females were more prominent than the 2-fold increases in the males. There was little difference in the levels of GTH

subunit mRNAs between animals received 75  $\mu g$  of GnRHa and those received 150  $\mu g$  of GnRHa.

#### DISCUSSION

In the present study, we examined effects of GnRHa on expression of GTH subunit genes in the pituitary gland of maturing sockeye salmon. The effects of GnRHa on expression of GTH  $\alpha$  and II $\beta$  genes were markedly different from that of GTH I $\beta$  gene. The levels of GTH  $\alpha$  and II $\beta$  mRNAs in the GnRHa-implanted fish were significantly higher than those in the control fish, whereas no significant difference was seen in the level of GTH I $\beta$  mRNA between the controls and GnRHa-implanted fish. These findings indicate that, in maturing salmonids, GnRH stimulates expression of GTH  $\alpha$  and II $\beta$  subunit genes in pituitary GTH II-cells.

The previous immunohistochemical and *in situ* hybridization study clearly demonstrated that GTH I and II are synthesized in separate pituitary GTH cells in salmonids (Naito *et al.*, 1991). GTH I gene expression was elevated in association with initiation of vitellogenesis, whereas GTH II gene expression was concomitantly elevated with final gonadal maturation in the pituitary of rainbow trout. Taking the previous studies (Kudo *et al.*, 1996; Kitahashi *et al.*, 1997) mentioned in Introduction into consideration, we consider that sGnRH selectively stimulates GTH II cells, but not GTH I cells, to elevate expression of GTH  $\alpha$  and II $\beta$  genes in pre-spawning salmonids. The levels of GTH  $\alpha$  and II $\beta$  mRNAs in the control fish were slightly higher than those in the initial controls. The magnitudes of increases and the amounts of mRNAs were comparable to our previous results in pre-spawning chum salmon in which the levels of GTH  $\alpha$  and II $\beta$  mRNAs were also elevated. It is therefore possible that endogenous sGnRH synthesis and release is stimulated during this period to accelerate final maturation through GTH II secretion.

Differences of GTH  $\alpha$  and II $\beta$  mRNA levels between the control and sGnRHa-implanted fish were prominent in the females when compared to those in the males. Difference in maturity between both sexes may account for this phenomenon, since the previous paper from our research group reported sexual differences in homing profiles and shortening of homing duration by GnRHa implantation in lacustrine sockeye salmon (Sato *et al.*, 1997). GTH II synthesis in the control males in the present study may be already somewhat stimulated.

It is well known in salmonids that plasma levels of steroid hormones, in particular  $17\alpha$ ,  $20\beta$ -dihydroxy-4-pregnen-3-one (DHP), are elevated in association with the final gamete maturation (Nagahama and Adachi, 1985). Hence we compared the present results of GTH subunit mRNA levels with plasma levels of steroid hormones, such as DHP, testosterone (T), and 11-ketotestosterone (11KT) in the males and DHP, T and estradiol- $17\beta$  (E<sub>2</sub>) in the females in the same animals (Fukaya *et al.*, in preparation). The results show that GnRHa induced a significant elevation of plasma DHP levels in both sexes, however, did not induce any noticeable changes in plasma T and 11KT levels in the males, and T and E<sub>2</sub> levels in the females. The increase in the levels of GTH  $\alpha$  and II $\beta$  mRNAs in the GnRHa-implanted fish correlated well with the increase in the plasma DHP level, particularly in the males.

In conclusion, implantation of GnRHa stimulates GTH II synthesis at the transcriptional level, and accelerates final maturation in maturing sockeye salmon. The acceleration of final maturation may be induced through elevation of plasma DHP level, which is stimulated by increase in plasma GTH II level.

# ACKNOWLEDGMENTS

We thank Dr. S. Itoh (Kyowa Hakko Kogyo Co., Ltd.), and Dr. D. S. Doering (AquaPharm Technologies Corp.) for the gift of cDNAs for chum salmon GTH subunits and GnRHa implants, respectively. We expressed our gratitude to colleagues in our laboratory for their help in sampling efforts. This study was supported in part by Grants-in-Aid from the Fisheries Agency of Japan, the Ministry of Education, Science, Sports and Culture, Japan and the Takeda Science Foundation.

# REFERENCES

- Amano M, Hyodo S, Kitamura S, Ikuta K, Suzuki Y, Urano A, Aida K (1995a) Salmon GnRH synthesis in the preoptic area and the ventral telencephalon is activated during gonadal maturation in female masu salmon. Gen Comp Endocrinol 99: 13–21
- Amano M, Hyodo S, Kitamura S, Ikuta K, Suzuki Y, Urano A, Aida K

(1995b) Short photoperiod accelerates preoptic and ventral telencephalic salmon GnRH synthesis and precocious maturation in under yearling male masu salmon. Gen Comp Endocrinol 99: 22–27

- Amano M, Urano A, Aida K (1997) Distribution and function of gonadotropin-releasing hormone (GnRH) in the teleost brain. Zool Sci 14: 1–11
- Chomczynski P, Sacchi N (1987) Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. Anal Biochem 162: 156–159
- Gen K, Maruyama O, Kato T, Tomizawa K, Wakabayashi K, Kato Y (1993) Molecular cloning of cDNAs encoding two types of gonadotropin  $\alpha$  subunit from the masu salmon, *Oncorhynchus masou*: construction of specific oligonucleotides for the  $\alpha$ 1 and  $\alpha$ 2 subunits. J Mol Endocrinol 11: 265–273
- Hiraoka S, Suzuki M, Yanagisawa T, Iwata M, Urano A (1993) Divergence of gene expression in neurohypophysial hormone precursors among salmonids. Gen Comp Endocrinol 92: 292–301
- Itoh H, Suzuki K, Kawauchi H (1988) The complete amino acid sequences of  $\beta$ -subunits of two distinct chum salmon GTHs. Gen Comp Endocrinol 71: 438–451
- Itoh H, Suzuki K, Kawauchi H (1990) The complete amino acid sequences of alpha subunits of chum salmon gonadotropins. Gen Comp Endocrinol 78: 56–65
- Kato Y, Gen K, Maruyama O, Tomizawa K, Kato T (1993) Molecular cloning of cDNAs encoding two gonadotrophin  $\beta$  subunits (GTH-I $\beta$  and -II $\beta$ ) from the masu salmon, *Oncorhynchus masou*. J Mol Endocrinol 11: 275–282
- Kawauchi H, Suzuki K, Itoh H, Swanson P, Naito N, Nagahama Y, Nozaki M, Nakai Y, Itoh S (1989) The duality of teleost gonadotropins. Fish Physiol Biochem 7: 29–38
- Kitahashi T, Ando H, Ban M, Urano A (1997) Changes in expression of gonadotropin gene in chum salmon during spawning migration. Zool Sci 14 (Suppl): 13
- Kobayashi M, Amano M, Kim MH, Yoshiura Y, Sohn YC, Suetake H, Aida K (1997) Gonadotropin-releasing hormone and gonadotropin in goldfish and masu salmon. Fish Physiol Biochem 17: 1–8
- Kudo H, Hyodo S, Ueda H, Hiroi O, Aida K, Urano A, Yamauchi K (1996) Cytophysiology of gonadotropin-releasing hormone neurons in chum salmon (*Oncorhynchus keta*) forebrain before and after upstream migration. Cell Tissue Res 284: 261–267
- Nagahama Y, Adachi S (1985) Identification of maturation-inducing steroid in a teleost, the amago salmon (*Oncorhynchus rhodurus*). Dev Biol 109: 428–435
- Naito N, Hyodo S, Okumoto N, Urano A, Nakai Y (1991) Differential production and regulation of gonadotropins (GTH I and GTH II) in the pituitary gland of rainbow trout, *Oncorhynchus mykiss*, during ovarian development. Cell Tissue Res 266: 457–467
- Sato A, Ueda H, Fukaya M, Kaeriyama M, Zohar Y, Urano A, Yamauchi K (1997) Sexual differences in homing profiles and shortening of homing duration by gonadotropin-releasing hormone analog implantation in lacustrine sockeye salmon (*Oncorhynchus nerka*) in Lake Shikotsu. Zool Sci 14: 1009–1014
- Sekine S, Saito A, Itoh H, Kawauchi H, Itoh S (1989) Molecular cloning and sequence analysis of chum salmon gonadotropin cDNAs. Proc Natl Acad Sci USA 86: 8645–8649
- Sherwood NM, Lovejoy DA, Coe IR (1993) Origin of mammalian gonadotropin-releasing hormones. Endocrinol Rev 14: 241–254
- Suzuki K, Liu D, Hew CL (1995) A gene encoding chinook salmon (*Oncorhynchus tschawytscha*) gonadotropin alpha subunit: gene structure and promoter analysis in primary pituitary cells. Mol Marine Biol Biotechnol 4: 10–19
- Trinh KY, Wang NC, Hew CL, Crim LW (1986) Molecular cloning and sequencing of salmon gonadotropin beta subunit. Eur J Biochem 159: 619–624
- Zohar Y (1996) New approaches for the manipulation of ovulation and spawning in farmed fish. Bull Natl Res Inst Aquacult Suppl

Downloaded From: https://complete.bioone.org/journals/Zoological-Science on 06 May 2025 Terms of Use: https://complete.bioone.org/terms-of-use

# 2: 43–48

Zohar Y, Pagelson G, Gothilf Y, Dickhoff WW, Swanson P, Duguay S, Gombotz W, Kost J, Langer R (1990) Controlled release of

gonadotropin releasing hormones for the manipulation of spawning in farmed fish. Control Rel Bioact Mater 17: 51-52

(Received March 20, 1998 / Accepted May 7, 1998)