

Localization of mRNA Encoding Thyrotropin-Releasing Hormone Precursor in the Brain of Sockeye Salmon

Authors: Ando, Hironori, Ando, Junko, and Urano, Akihisa

Source: Zoological Science, 15(6) : 945-953

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.15.945>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Localization of mRNA Encoding Thyrotropin-Releasing Hormone Precursor in the Brain of Sockeye Salmon

Hironori Ando*, Junko Ando and Akihisa Urano

*Division of Biological Sciences, Graduate School of Science, Hokkaido University,
Sapporo, Hokkaido 060-0810, Japan*

ABSTRACT—The distribution of mRNA encoding thyrotropin-releasing hormone (TRH) precursor in the brain of sockeye salmon was studied by *in situ* hybridization histochemistry using digoxigenin-labeled riboprobes as a basis to investigate its physiological functions in the salmon brain. Since seasonal variation in TRH gene expression was expected in relation to smolting or maturation, fish were sampled in February and October. In both groups, TRH mRNA was widely distributed in discrete brain regions including the internal cellular layer (ICL) of the olfactory bulb, postcommissural nucleus of area ventralis telencephali (Vp), nucleus preopticus parvocellularis anterioris (PPa), nucleus preopticus magnocellularis, dorsal zone of periventricular hypothalamus (Hd), torus semicircularis, and also the motor nucleus of vagus nerve in the medulla oblongata. TRH neurons in ICL and Hd are round and small with diameters of 5–10 μm . In contrast, TRH neurons in the ventral telencephalon and the preoptic area are medium-sized (10–20 μm), and appear to have multiple processes. Most of these cells are restrictively localized along the lateral margin of the preoptic nuclei. The number of TRH neurons in Vp and PPa was smaller in February than in October, suggesting a seasonal change of TRH neurons in the preoptic area. In the medulla oblongata, a cluster of large oval-shaped cells (20–30 μm) showed signals for TRH mRNA. The present results suggest that TRH may function as a neurotransmitter or neuromodulator involved in olfactory activity and also autonomic motor integration, in addition to neurohormonal control of secretion of pituitary hormones.

INTRODUCTION

Thyrotropin-releasing hormone (TRH) is a tripeptide neurohormone (pGlu-His-Pro-NH₂), which was first isolated from the mammalian hypothalamus as a factor that stimulates secretion of thyroid-stimulating hormone (TSH) (Schally *et al.*, 1969; Burgus *et al.*, 1970). Stimulatory roles of TRH on the pituitary-thyroid axis have been well studied in mammals (Jackson 1982). TRH immunoreactivity has been shown in the hypothalamic tuberoinfundibular system and also in the extrahypothalamic brain regions by radioimmunoassay (RIA) (Jackson and Reichlin, 1974) and immunohistochemistry (Lechan and Jackson, 1982; Merchenthaler *et al.*, 1988; Fliers *et al.*, 1994). Extrahypothalamic TRH may function as a neurotransmitter or neuromodulator in the central nervous system as well as in the gastrointestinal tract (Jackson 1982). In nonmammalian vertebrates, however, the physiological functions and distribution of TRH have been controversial (Jackson and Reichlin, 1974; Ball, 1981).

TRH stimulated TSH release in amphibians (Denver, 1988) and reptiles (Preece and Licht, 1987), whereas earlier studies had shown no effects (see Ball, 1981). TRH also stimulated release of prolactin (PRL) and α -melanocyte-stimulat-

ing hormone (α -MSH) in amphibians (Toron *et al.*, 1983; Seki and Kikuyama, 1986). In birds, TRH elevated plasma levels of thyroid hormones and growth hormone (GH) (Leung *et al.*, 1984). Actions of TRH in fishes are more confusing (see Crim *et al.*, 1978). Some studies showed that TRH stimulates pituitary-thyroid axis in various teleosts (Tsuneki and Fernholm, 1975; Eales and Himick, 1988), whereas other studies could not find any effects (Bromage *et al.*, 1976). Furthermore, Bromage (1975) suggested an inhibitory role of TRH on TSH release in guppy. As in other vertebrate classes, TRH stimulated release of pituitary hormones other than TSH in teleosts, such as PRL (Barry and Grau, 1986), GH (Trudeau *et al.*, 1992) and α -MSH (Lamers *et al.*, 1994).

Localization of TRH immunoreactive (TRH-ir) cells in the fish brains, particularly in the preoptic area, was not consistent among several studies. Hamano *et al.* (1990) showed TRH-ir cell bodies only in the nucleus recessus lateralis (NRL) of carp. Batten *et al.* (1990) demonstrated a number of TRH-ir cells in the NRL, but only a few TRH-ir cells in the nucleus preopticus (PON) of sea bass. On the other hand, Matz and Takahashi (1994) showed the presence of TRH-ir cells in the olfactory bulb (OB), ventral telencephalon and the preoptic area in the brain of chinook salmon. The distribution of TRH-ir cells in the PON along with TRH-ir fibers in the hypothalamus and pituitary is consistent with the hypophysiotropic role of TRH. Recently, Ohno *et al.* (1995) demonstrated TRH-ir

* Corresponding author: Tel. +81-11-706-2995;
FAX. +81-11-706-4923.

cells in the preoptic area of sockeye salmon. TRH-ir cells in the OB of chum salmon were reported by Hamano *et al.* (1996).

Immunohistochemical localization of TRH in the brain has two major problems: (1) difficulty of raising potent and specific antibodies to TRH, and (2) loss of the peptide or alterations of its antigenic determinants during tissue fixation (Merchenthaler *et al.*, 1988). To overcome these difficulties, and more importantly to elucidate the mechanism of TRH biosynthesis, some investigators used an antiserum that recognize TRH prohormone (pro-TRH) (Lechan *et al.*, 1986; Merchenthaler *et al.*, 1989; Hamano *et al.*, 1996). In the rat brain, pro-TRH-ir cells were more widely distributed than TRH-ir cells, suggesting that TRH prohormone may be differentially processed in certain regions of the brain. *In situ* hybridization (ISH) of mRNAs for TRH precursor and also prohormone convertase (PC) demonstrated that PC genes were coexpressed in a subset of pro-TRH-expressing neurons in the rat (Segerson *et al.*, 1987; Pu *et al.*, 1996) and frog (Pu *et al.*, 1995).

It is, therefore, important to determine in the brain the distribution of cells containing TRH mRNA to address physiological roles of TRH and also of non-TRH peptides or extended forms of TRH peptides differentially processed from pro-TRH. Recently, we have isolated cDNAs encoding TRH precursors of sockeye salmon (Ohide *et al.*, 1996). In the present study, the distribution of TRH mRNA in the sockeye salmon brain was studied by ISH. In addition, we found a seasonal variation of TRH gene expression in the ventral telencephalon and anterior part of the preoptic area.

MATERIALS AND METHODS

Animals

Three-year-old sockeye salmon, *Oncorhynchus nerka*, of both sexes were obtained from the Toya Lake Station for Environmental Biology, Japan in October 1995 and February 1996. The fork length of the fish ranged from 20.6 to 23.5 cm, weighing from 82 to 139 g in the former fish, and from 21.2 to 24.4 cm, weighing from 107 to 164 g in the latter fish. Three males and three females of each month were used for ISH. They were not reproductively matured yet (GSI 0.03–0.11% for males, 0.50–0.89% for females) except for one male (GSI 3.5%) and two females (GSI 15 and 17%) in October.

Tissue preparation

Fish were lightly anaesthetized in 0.05% phenoxy ethanol. Immediately after decapitation, brains were dissected out and were fixed in 4% paraformaldehyde in 0.05 M phosphate buffer (pH 7.3) at 4°C for 20 hr. After fixation, the tissues were washed in 70% ethanol at 4°C for 3 days, dehydrated through a series of ethanol solutions, and were embedded in paraplast. When embedded, the brains were placed ventral side up on the bottom of a paraffin cup, and were sectioned at 6 µm in either the sagittal or transverse plane at right angles to the plane parallel to the cranial bone (Fig. 1). They were separated into five groups, and were mounted on 3-amino-propyltriethoxysilane coated slides.

In situ hybridization (ISH)

ISH was performed as previously described (Zwingman *et al.*, 1994) with some modifications. Briefly, after rehydration, sections were treated with 0.3% Triton X-100 and 0.2 M HCl, immersed in acetic anhydride (0.25% in triethanolamine 0.1 M, pH 8.0), and treated with

0.2% glycine. Digoxigenin-labelled riboprobes were synthesized with the sockeye salmon pro-TRH-A cDNA (Ohide *et al.*, 1996) cloned in pGEM-T (Promega, WI, USA). After linearization with *Spe* I, sense and antisense probes were transcribed *in vitro* with digoxigenin-UTP using a RNA labelling kit (Boehringer Mannheim) according to the manufacturer's instruction.

After prehybridization in 5 × SSC (1 × SSC = 150 mM NaCl, 15 mM sodium citrate, pH 4.5) containing 50% formamide at room temperature for 30 min, hybridization was carried out at 42°C overnight in a hybridization buffer (50% formamide, 10% dextran sulfate, 5 × SSC, pH 4.5, 1 × Denhardt's solution, 0.1 mg/ml heparin, 0.1 mg/ml calf thymus DNA, 0.2 mg/ml yeast tRNA), containing digoxigenin-labelled riboprobes. Then, the sections were washed in 4 × SSC at 42°C for 20 min, treated with RNase A (0.02 mg/ml) in buffer (0.5 M NaCl, 10 mM Tris-HCl, pH 7.5) at 37°C for 1 hr, and washed sequentially with 2 × SSC and 0.1 × SSC at 50°C for 30 min twice. Hybridization signals were visualized using a nucleic acid detection kit (Boehringer Mannheim) with alkaline-phosphatase conjugated antidigoxigenin antibody and nitroblue tetrazolium (NBT), according to the manufacturer's instruction.

Data analysis

TRH mRNA-containing neurons in parasagittal and frontal sections, taken from different levels of the brain, were drawn under a microscope with a camera lucida. For histological identification of the localization of these neurons, one group of sections were stained with cresyl violet. The nomenclature for brain loci was taken from Northcutt and Davis (1983) for the telencephalon, Braford and Northcutt (1983) for the diencephalon and pretectum, and Nieuwenhuys and Pouwels (1983) for the brain stem. The abbreviations of brain loci are listed in Table 1.

Cell measurement and statistics

The number of TRH mRNA-containing cells in the preoptic area was counted by examining every section at 30 µm intervals to estimate the total number of TRH neurons contained in each brain. Differences in the number of TRH neurons in the Vp and PPa regions between the fish in February and October were tested by Mann-Whitney U-test.

RESULTS

Localization and morphology of TRH mRNA-containing cells

ISH for TRH mRNA demonstrated the presence of hybridization positive cells in several regions of the forebrain and the brain stem. An overview of the localization of these cells is shown in Fig. 1.

Olfactory bulb: TRH mRNA-containing cells were scattered in the internal cellular layer (ICL), whereas no hybridization signals were observed in the external cellular layer (ECL) (Fig. 2a–c). TRH cells in the ICL were dispersed among non-TRH neurons. They were round and small with diameters of 5–10 µm (Fig. 3a). Some TRH cells were located in the neighboring region of the ventral nucleus of area ventralis telencephali (Vv) (Fig. 2c).

Telencephalon: TRH cells were located in the postcommissural nucleus of area ventralis telencephali (Vp) (Fig. 2d–f). Most of TRH cells were scattered in the lateral region of Vp. The diameter of round somata ranged 5–10 µm. No hybridization positive cells were observed in the anterior region to the anterior commissure (AC) in the telencephalon.

Table 1. Abbreviations

AC	anterior commissure	PMg	nucleus preopticus magnocellularis, pars gigantocellularis
C	cerebellum	PMm	nucleus preopticus magnocellularis, pars magnocellularis
D	area dorsalis telencephali	PMp	nucleus preopticus magnocellularis, pars parvocellularis
DF	nucleus diffusus of the inferior lobe	PON	nucleus preopticus
Dc	central zone of D	PPa	nucleus preopticus parvocellularis anterioris
Dd	dorsal zone of D	PPp	nucleus preopticus parvocellularis posterioris
Dld	dorsal part of lateral zone of D	Psm	nucleus preopticus superficialis, pars magnocellularis
Dlv	ventral part of lateral zone of D	SC	nucleus suprachiasmaticus
Dm	medial zone of D	SV	sacculus vasculosus
Dp	posterior zone of D	T	telencephalon
E	entopeduncular nucleus	TL	nucleus tuberis lateralis
ECL	external cellular layer of olfactory bulb	TL0	torus longitudinalis
G	nucleus glomerulosus	TP	nucleus tuberis posterior
HA	habenula	TS	torus semicircularis
HT	hypothalamus	V	area ventralis telencephali
Hd	dorsal zone of periventricular hypothalamus	VCg	granular layer of the valvula cerebelli
Hv	ventral zone of periventricular hypothalamus	Vcm	molecular layer of the valvula cerebelli
ICL	internal cellular layer of olfactory bulb	VL	nucleus ventrolateralis
LH	nucleus lateralis hypothalami	VM	nucleus ventromedialis
LV	nucleus lateralis valvulae	Vd	dorsal nucleus of V
M	medulla oblongata	Vp	postcommissural nucleus of V
nMLF	nucleus of the medial longitudinal fasciculus	Vv	ventral nucleus of V
OB	olfactory bulb		
OC	optic chiasm		
ON	optic nerve		
OT	optic tectum		
PIT	pituitary		




Fig. 1. Schematic illustration of the distribution of TRH mRNA-containing cells in the brain of sockeye salmon. Large dots represent TRH-expressing perikarya, and small dots indicate the Nissl-stained neurons. Planes of section for Fig. 2a-o are indicated. These planes are vertical to the plane parallel to the cranial bone.

Diencephalon: The largest number of cells containing TRH mRNA was observed in the hypothalamus. In the preoptic area, they were located in the nucleus preopticus parvocellularis anterioris (PPa) and nucleus preopticus magnocellularis (PM) (Fig. 2e-k). TRH cells in the PPa were mostly dispersed in the lateral margin of the PPa region (Figs. 2e-g, 3b), and their diameter ranged 10-15 μ m. On the other hand, slightly larger cells (10-20 μ m in diameter) showing intense hybridization signals were observed in the nucleus

preopticus magnocellularis pars magnocellularis (PMm) and the nucleus preopticus magnocellularis pars gigantocellularis (PMg) (Fig. 2h-k). They were confined to the lateral border of the PM region, and appeared to have multiple processes (Fig. 3c and d). Some cells formed clusters, whereas the others were scattered among other neurons (Fig. 3d). In the ventral hypothalamus, weakly hybridization positive cells were located in the dorsal zone of periventricular hypothalamus (Hd) (Fig. 2l and m). They were round and small with diameters of





Fig. 2. (a-o) Schematic representation of TRH mRNA-containing perikarya (dots) in the salmon brain in frontal sections. Abbreviations are shown in Table 1.



5-10 μ m.

Mesencephalon: Weakly hybridizing cells were distributed in the region lateral to the torus semicircularis (TS) in the midbrain tegmentum (Figs. 2n and o, 3e).

Myelencephalon: A cluster of large oval-shaped cells (20-30 μ m in diameter) showed TRH mRNA signals (Fig. 3f). They were located in the motor nucleus of vagus nerve in the medulla oblongata (Fig. 1).

Controls: In contrast to the specific hybridization with the antisense probe, parallel *in situ* hybridization studies with

the sense riboprobe did not show any hybridization signals in all regions of the brain studied.

Seasonal variation in the number of TRH cells in the PPa and Vp regions

The number of TRH mRNA-containing cells in the preoptic area was examined in February (prior to smolting season) and October (spawning season). The number in the PPa and Vp regions was significantly smaller in the fish in February than in October, whereas no difference was observed in the




Fig. 3. TRH mRNA-containing perikarya in the salmon brain. (a) Coronal section through the internal cellular layer (ICL) of the olfactory bulb. (b) Coronal section through the nucleus preopticus parvocellularis anterioris (PPa). (c) Coronal section through the nucleus preopticus magnocellularis (PM). Note that TRH cells are confined to the lateral margin of the PM region. (d) Sagittal section through the PM. A cluster of TRH cells was observed. (e) Coronal section through the torus semicircularis (TS) in the mesencephalon. TRH cells were dispersed laterally. (f) Sagittal section through the motor nucleus of vagus nerve in the medulla oblongata. A cluster of large oval-shaped cells (20–30 μm in diameter) showed TRH mRNA signals. Scale bar, 50 μm .




Fig. 4. The numbers of TRH mRNA-containing cells in the postcommissural nucleus of area ventralis telencephali (Vp) and nucleus preopticus parvocellularis anterioris (PPa) regions, and the nucleus preopticus magnocellularis pars magnocellularis (PMm) and nucleus preopticus magnocellularis pars gigantocellularis (PMg) regions of the salmon brain in October (open column) and February (solid column). * $P < 0.05$.

PMm and PMg regions (Fig. 4). Other regions containing TRH cells also did not show any apparent difference in the number of hybridization positive cells between February and October.

DISCUSSION

The present study demonstrated the distribution of TRH mRNA-containing cells for the first time in fish brain. ISH clearly showed TRH mRNA distribution in various regions of the salmon brain. We used riboprobes which were synthesized with the sockeye salmon TRH-A cDNA (Ohide *et al.*, 1996). Ohide *et al.* (1996) showed that in sockeye salmon there are at least two types of TRH encoding genes (TRH-A and -B), which are highly homologous with each other (90% in a 242 bp region). Therefore, the ISH in this study may reveal distribution of both types of TRH mRNAs in the sockeye salmon brain. Differences in the distribution and amount of these mRNAs are unknown at present and should be determined in future using specific probes.

TRH mRNA-containing cells were distributed in several regions of the forebrain and the brain stem. Most prominent hybridization signals were observed in cells of the preoptic area. TRH cells were located in the nucleus preopticus parvocellularis anterioris (PPa) and nucleus preopticus magnocellularis (PM). Matz and Takahashi (1994) have demonstrated the presence of TRH-ir cell bodies in the PM but not in the PPa in the chinook salmon. Lack of TRH-ir cell bodies in the PPa is possibly because they used the fish in March and April when the level of expression of TRH encoding gene may be low in the PPa region as shown in the sockeye salmon in February (Fig. 4). It is also possible that in the PPa region the pro-TRH might be differentially processed to generate non-TRH peptides or extended forms of TRH peptides as proposed

in the rat brain (Pu *et al.*, 1996). Otherwise, it may be because of the differences in species or stage. Ohno *et al.* (1995) have shown TRH-ir cells in the preoptic area of the sockeye salmon, but the exact location of the TRH-ir cell bodies was not specified. The results of the immunohistochemical and present ISH studies clearly demonstrate the presence of TRH cells in the preoptic area in salmon as reported in other vertebrates (rat, Lechan and Jackson, 1982; mallard, Jozsa *et al.*, 1988; bullfrog, Seki *et al.*, 1983), in concordance with the hypophysiotropic role of TRH. In the earlier studies, however, TRH-ir cell bodies could not be localized in the preoptic area in the carp (Hamano *et al.*, 1990) and sea bass (Batten *et al.*, 1990). This discrepancy may be because of the difficulty of immunohistochemistry of TRH peptide (Merchenthaler *et al.*, 1988).

TRH cells in the preoptic area were frequently seen in the lateral marginal part of the nuclei (Fig. 3b and c). This feature is most apparent in the PMm. Because TRH-ir cells were shown to be dispersed within the preoptic area in the other classes of vertebrates (Lechan and Jackson, 1982; Seki *et al.*, 1983; Jozsa *et al.*, 1988), the distribution of TRH cells in the lateral border of the preoptic area may be unique to the salmon brain. In chinook salmon such tendency in the localization of TRH-ir cells in the PM was not mentioned (Matz and Takahashi, 1994).

In the preoptic area, vasotocin (VT) and isotocin (IT) neurons are densely localized in the PM. IT neurons are further distributed in the PPa (Ota *et al.*, 1996). Moreover, in the preoptic area, distributed are neurons containing hypothalamic releasing hormones, such as gonadotropin-releasing hormone (GnRH) (Amano *et al.*, 1991) and corticotropin-releasing hormone (CRH) (Ando *et al.*, 1999). Meister *et al.* (1990) have demonstrated a small number of both vasopressin (VP) and oxytocin (OT) neurons colocalized with TRH in the rat paraventricular nucleus. VP has been shown to elicit release of TSH (Lumpin *et al.*, 1987), indicating that this colocalization may be functional. In salmon brain, although the colocalizations of TRH and VT, IT and/or CRH remain to be determined, TRH cells are likely to interact with these neurosecretory neurons and control pituitary functions.

It is of interest to note that the number of TRH cells in the preoptic area showed seasonal variation (Fig. 4). Seasonal variations in neurosecretory neurons are deeply involved in control of seasonal changes in physiological activities (Jokura and Urano, 1985; Hofman and Swaab, 1992). In the present study, the number of TRH cells in the PPa and Vp region was significantly different between February and October, whereas that in the PM region was not. This suggests that physiological functions of the TRH cells in the preoptic area are different between the PPa and PM nuclei. In mammals, the magnocellular neurons contain VP and OT, while the parvocellular neurons contain the hypothalamic releasing hormones. However, such anatomical distinctions have not been observed in salmon brain; IT (Ota *et al.*, 1996) and GnRH (Amano *et al.*, 1991) neurons are distributed in both PPa and PM nuclei. Therefore, it is difficult to correlate the location of

TRH cells with their function at present. Precise studies on colocalization of TRH and other neurohormones and also studies to determine to which types of pituitary cells the preoptic TRH neurons project their axons are necessary to clarify the physiological significance of the seasonal variation of TRH cells in the preoptic area.

Localization of TRH mRNA-containing cells in the ICL of the olfactory bulb (Fig. 3a) corresponds to the distribution of TRH-ir cell bodies (Matz and Takahashi, 1994). TRH-ir cell bodies in the ICL have been suggested to contribute TRH fibers in the olfactory tracts, and to project to olfactory targets in the ventral telencephalon, the preoptic area and the hypothalamic regions, all of which are known to receive olfactory projections in teleosts (Matz and Takahashi, 1994). Therefore, TRH is considered to play a role in the olfactory system, may act as a transmitter or neuromodulator in the salmon brain. Hamano *et al.* (1996) showed proTRH-ir neurons in the ECL, but not in the ICL in the chum salmon. This discrepancy might be due to the difference of animal stock. Hamano *et al.* (1996) used a wild stock of the chum salmon during the spawning migration, while we and Matz and Takahashi (1994) used salmon fed in the hatchery. In homing salmon, activities and functions of TRH neurons in the olfactory bulb might be different from those in the salmon fed in hatchery.

The present study showed the presence of TRH mRNA in the motor nucleus of vagus nerve in the medulla oblongata. Here the TRH cells are larger (20–30 μm) than those in other brain regions. The presence of TRH mRNA and TRH-ir in the dorsal motor nucleus of the vagus were reported in the rat (Lechan *et al.*, 1986; Segerson *et al.*, 1987). It was suggested that TRH in the dorsal vagal complex may function in the control of gastric secretion and motility in the rat (Rinaman *et al.*, 1989). Furthermore, TRH in the rat medulla was involved in modulation of respiratory function (McCown *et al.*, 1986). In teleost medulla, the motor nucleus of vagus nerve was suggested to be involved in the control of rhythmic breathing movements (Shelton, 1970). Therefore, TRH in the motor nucleus of vagus nerve may function as a neurotransmitter or neuromodulator involved in the autonomic control of breathing also in salmon.

The present study demonstrates a precise distribution of the TRH mRNA-containing neurons in the salmon brain. Based upon the comparison with the distribution of TRH-ir cells, the TRH mRNA-positive cells are more widely distributed than TRH-ir cells. Particularly, the possibility of differential splicing of pro-TRH was suggested in the PPa region. Interestingly, in this region the number of TRH cells was seasonally variable. It should be of considerable interest and importance to determine the physiological roles of TRH cells in the preoptic area in salmon brain. The present study also suggests the multiple TRH neuronal systems; TRH may function in the extrahypothalamic neuromodulatory systems involved in olfactory and respiratory functions as well as the conventional hypophysiotropic system.

ACKNOWLEDGMENTS

We thank Dr. H. Ueda (the Toya Lake Station for Environmental Biology) for providing sockeye salmon, and Mr. T. Ohya for technical assistance. This study was supported in part by the Grants-in-Aid from the Fisheries Agency and the Ministry of Education, Science, Sports and Culture, Japan.

REFERENCES

- Amano M, Oka Y, Aida K, Okumoto N, Kawashima S, Hasegawa Y (1991) Immunocytochemical demonstration of salmon GnRH and chicken GnRH-II in the brain of masu salmon, *Oncorhynchus masou*. *J Comp Neurol* 314: 587–597
- Ando H, Hasegawa M, Ando J, Urano A (1999) Expression of salmon corticotropin-releasing hormone precursor gene in the preoptic nucleus in stressed rainbow trout. *Gen Comp Endocrinol*, in press
- Ball JN (1981) Hypothalamic control of the pars distalis in fishes, amphibians, and reptiles. *Gen Comp Endocrinol* 44: 135–170
- Barry TP, Grau EG (1986) Estradiol-17 β and thyrotropin-releasing hormone stimulate prolactin release from the pituitary gland of a teleost fish *in vitro*. *Gen Comp Endocrinol* 62: 306–314
- Batten TFC, Moons L, Cambre ML, Vandesande F, Seki T, Suzuki M (1990) Thyrotropin-releasing hormone-immunoreactive system in the brain and pituitary gland of the sea bass (*Dicentrarchus labrax*, Teleostei). *Gen Comp Endocrinol* 79: 385–392
- Braford MR, Northcutt RG (1983) Organization of the diencephalon and pretectum of the ray-finned fishes. In "Fish Neurobiology Vol 2" Ed by RE Davis, RG Northcutt, University of Michigan Press, Ann Arbor, pp 117–164
- Bromage NR (1975) The effects of mammalian thyrotropin-releasing hormone on the pituitary-thyroid axis of teleost fish. *Gen Comp Endocrinol* 25: 292–297
- Bromage NR, Whitehead C, Brown TJ (1976) Thyroxine secretion in teleosts and the effects of TSH, TRH, and other peptides. *Gen Comp Endocrinol* 29: 246 (abstract)
- Burgus R, Dunn TF, Desiderio D, Ward DN, Vale W, Guillemin R (1970) Characterization of ovine hypothalamic hypophysiotropic TSH-releasing factor. *Nature* 226: 321–325
- Crim JW, Dickhoff WW, Gorbman A (1978) Comparative endocrinology of piscine hypothalamic hypophysiotropic peptides: Distribution and activity. *Am Zool* 18: 411–424
- Denver RJ (1988) Several hypothalamic peptides stimulate *in vitro* thyrotropin secretion by pituitaries of anuran amphibians. *Gen Comp Endocrinol* 72: 383–393
- Eales JG, Himick BA (1988) The effects of TRH on plasma thyroid hormone levels of rainbow trout (*Salmo gairdneri*) and arctic charr (*Salvelinus alpinus*). *Gen Comp Endocrinol* 72: 333–339
- Fliers E, Noppen NWAM, Wiersinga WM, Visser TJ, Swaab DF (1994) Distribution of thyrotropin-releasing hormone (TRH)-containing cells and fibers in the human hypothalamus. *J Comp Neurol* 350: 311–323
- Hamano K, Inoue K, Yanagisawa T (1990) Immunohistochemical localization of thyrotropin-releasing hormone in the brain of carp, *Cyprinus carpio*. *Gen Comp Endocrinol* 80: 85–92
- Hamano K, Yoshida K, Suzuki M, Ashida K (1996) Changes of thyrotropin-releasing hormone concentration in the brain and levels of prolactin and thyroxine in the serum during spawning migration of the chum salmon *Oncorhynchus keta*. *Gen Comp Endocrinol* 101: 275–281
- Hofman MA, Swaab DF (1992) Seasonal changes in the suprachiasmatic nucleus of man. *Neurosci Lett* 139: 257–260
- Jackson IMD (1982) Thyrotropin-releasing hormone. *New Engl J Med* 306: 145–155
- Jackson IMD, Reichlin S (1974) Thyrotropin-releasing hormone (TRH): Distribution in hypothalamic and extra hypothalamic brain tissues

- of mammalian and submammalian chordates. *Endocrinology* 95: 854–862
- Jokura Y, Urano A (1985) An immunohistochemical study of seasonal changes in luteinizing hormone-releasing hormone and vasotocin in the forebrain and the neurohypophysis of the toad, *Bufo japonicus*. *Gen Comp Endocrinol* 59: 238–245
- Jozsa R, Korf HW, Csernus V, Mess B (1988) Thyrotropin-releasing hormone (TRH)-immunoreactive structures in the brain of the domestic mallard. *Cell Tissue Res* 251: 441–449
- Lamers AE, Flik G, Wendelaar Bonga SE (1994) A specific role for TRH in release of diacetyl α -MSH in tilapia stressed by acid water. *Am J Physiol* 267: R1302–R1308
- Lechan RM, Jackson IMD (1982) Immunohistochemical localization of thyrotropin-releasing hormone in the rat hypothalamus and pituitary. *Endocrinology* 111: 55–65
- Lechan RM, Wu P, Jackson IMD (1986) Immunolocalization of the thyrotropin-releasing hormone prohormone in the rat central nervous system. *Endocrinology* 119: 1210–1216
- Leung FC, Taylor JE, van-Iderstine A (1984) Thyrotropin-releasing hormone stimulates body weight gain and increases thyroid hormones and growth hormone in plasma of cockerels. *Endocrinology* 115: 736–740
- Lumpin MD, Samson WK, McCann SM (1987) Arginine vasopressin as a thyrotropin-releasing hormone. *Science* 235: 1070–1073
- Matz SP, Takahashi TT (1994) Immunohistochemical localization of thyrotropin-releasing hormone in the brain of chinook salmon (*Oncorhynchus tshawytscha*). *J Comp Neurol* 345: 214–223
- McCown TJ, Hedner JA, Towle AC, Breese GR, Mueller RA (1986) Brainstem localization of a thyrotropin-releasing hormone-induced change in respiratory function. *Brain Res* 373: 189–196
- Meister B, Villar MJ, Ceccatelli S, Hokfelt T (1990) Localization of chemical messengers in magnocellular neurons of the hypothalamic supraoptic and paraventricular nuclei: An immunohistochemical study using experimental manipulations. *Neuroscience* 37: 603–633
- Mercenthaler I, Csernus V, Csontos C, Petrusz P, Mess B (1988) New data on the immunocytochemical localization of thyrotropin-releasing hormone in the rat central nervous system. *Am J Anat* 181: 359–376
- Mercenthaler I, Meeker M, Petrusz P, Kizer JS (1989) Identification and immunocytochemical localization of a new thyrotropin-releasing hormone precursor in rat brain. *Endocrinology* 124: 1888–1897
- Nieuwenhuys R, Pouwels E (1983) The brain stem of actinopterygian fishes. In "Fish Neurobiology Vol 1" Ed by RE Davis, RG Northcutt, University of Michigan Press, Ann Arbor, pp 25–88
- Northcutt RG, Davis RE (1983) Telencephalic organization in ray-finned fishes. In "Fish Neurobiology Vol 2" Ed by RE Davis, RG Northcutt, University of Michigan Press, Ann Arbor, pp 203–236
- Ohide A, Ando H, Yanagisawa T, Urano A (1996) Hydropathy profiles of predicted thyrotropin-releasing hormone precursors are highly conserved despite low similarity of primary structures. *J Neuroendocrinol* 8: 695–701
- Ohno N, Ohide A, Sugimura N, Sato T, Yangisawa T, Parhar I, Iwata M (1995) Distribution of thyrotropin-releasing hormone in the sockeye salmon brain and its putative physiological role. *Proc Japan Soc Comp Endocrinol* 10: 78
- Ota Y, Ando H, Ban M, Ueda H, Urano A (1996) Sexually different expression of neurohypophysial hormone genes in the preoptic nucleus of pre-spawning chum salmon. *Zool Sci* 13: 593–601
- Preece H, Licht P (1987) Effects of thyrotropin-releasing hormone *in vitro* in thyrotropin and prolactin release from the turtle pituitary. *Gen Comp Endocrinol* 67: 247–255
- Pu LP, Hayes WP, Mill JF, Ghose S, Freidman TC, Loh YP (1995) Frog prohormone convertase PC2 mRNA has a mammalian-like expression pattern in the central nervous system and is colocalized with a subset of thyrotropin-releasing hormone-expressing neurons. *J Comp Neurol* 354: 71–86
- Pu LP, Ma W, Barker JL, Loh YP (1996) Differential coexpression of genes encoding prothyrotropin-releasing hormone (Pro-TRH) and prohormone convertases (PC1 and PC2) in rat brain neurons: Implications for differential processing of Pro-TRH. *Endocrinology* 137: 1233–1241
- Rinaman L, Miselis RR, Kreider MS (1989) Ultrastructural localization of thyrotropin-releasing hormone immunoreactivity in the dorsal vagal complex in rat. *Neurosci Lett* 104: 7–12
- Schally A, Redding T, Bowers C, Barret J (1969) Isolation and properties of porcine thyrotropin-releasing hormone. *J Biol Chem* 244: 4077–4088
- Segerson TP, Hoefler H, Childers H, Wolfe HJ, Wu P, Jackson IMD, Lechan RM (1987) Localization of thyrotropin-releasing hormone prohormone messenger ribonucleic acid in rat brain by *in situ* hybridization. *Endocrinology* 121: 98–107
- Seki T, Nakai Y, Shioda S, Mitsuma T, Kikuyama S (1983) Distribution of immunoreactive thyrotropin-releasing hormone in the forebrain and hypophysis of the bullfrog, *Rana catesbeiana*. *Cell Tissue Res* 233: 507–516
- Seki T, Kikuyama S (1986) Effect of thyrotropin-releasing hormone and dopamine on the *in vitro* secretion of prolactin by the bullfrog pituitary gland. *Gen Comp Endocrinol* 61: 197–202
- Shelton G (1970) The regulation of breathing. In "Fish Physiology Vol 4" Ed by WS Hoar, DJ Randall, Academic Press, New York and London, pp 293–359
- Toron MC, Leroux P, Stockel ME, Jegou S, Pelletier G, Vaudry H (1983) Catecholaminergic control of α -melanocyte stimulating hormone (α -MSH) release by frog neurointermediate lobe *in vitro*: Evidence for direct stimulation of α -MSH release by thyrotropin-releasing hormone. *Endocrinology* 112: 133–141
- Trudeau VL, Somoza GM, Nahorniak CS, Peter RE (1992) Interactions of estradiol with gonadotropin-releasing hormone and thyrotropin-releasing hormone in the control of growth hormone secretion in the goldfish. *Neuroendocrinology* 56: 483–490
- Tsuneki K, Fernholm B (1975) Effect of thyrotropin-releasing hormone on the thyroid of a teleost, *Chasmichthys dolichognathus*, and a hagfish, *Eptatretus burgeri*. *Acta Zool* 56: 61–65
- Zwingman T, Fujimoto H, Lai LW, Boyer T, Ao A, Stalvey JRD, Blecher SR, Erickson RP (1994) Transcription of circular and noncircular forms of *Sry* in mouse testes. *Mol Reprod Develop* 37: 370–381

(Received July 10, 1998 / Accepted September 16, 1998)