

A Comparative Study on Histochemical Distribution of Some Enzymes Related to Steroid and Glucuronide Synthesis in Seminal Vesicle and Testis of the Catfish, Clarias batrachus

Authors: Singh, Madhu S., and Joy, Keerikkattil P.

Source: Zoological Science, 15(6): 955-961

Published By: Zoological Society of Japan

URL: https://doi.org/10.2108/zsj.15.955

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A Comparative Study on Histochemical Distribution of Some Enzymes Related to Steroid and Glucuronide Synthesis in Seminal Vesicle and Testis of the Catfish, *Clarias batrachus*

Madhu S. Singh and Keerikkattil P. Joy*

Department of Zoology, Banaras Hindu University, Varanasi-221 005, India

ABSTRACT—In both seminal vesicle (SV) and testis of the catfish *Clarias batrachus*, a large number of interstitial cells are distributed, singly or in groups close to blood capillaries in the interstices of the secretory lobules and seminiferous tubules, respectively. They showed strong reactions for 3 β -HSD, G-6-PD, NADH diaphorase and UDPGD activities. Both the germinal epithelium and SV epithelium showed moderate to strong reaction for UDPGD activity. These results strongly indicate that the SV is steroidogenic and capable of steroid glucuronidation, like the testis. It is suggested that the interstitial cells of the testis and SV are homologous having a common embryonic origin from the genital ridge.

INTRODUCTION

In teleosts, as in other vertebrates, testicular interstitial (Leydig) cells, which are distributed usually in the interstices of seminiferous tubules singly or in clusters are the main site of production of androgens (Lofts and Bern, 1972; Guraya, 1976; Fostier et al., 1983; Nagahama, 1983; Lofts, 1987). In addition, lobule boundary cells or Sertoli cells have also been implicated in steroidogenesis in some teleosts (references as above). The steroidogenic cells are characterized by the presence of lipids which serve as the precursor of steroid hormones, and enzymes of steroid hydroxylations, such as Δ^{5} -3β-hydroxysteroid dehydrogenase (3β-HSD) which catalyzes the conversion of Δ^5 -hydroxysteroids, such as pregnenolone and dehydroepiandrosterone, to Δ^4 -ketosteroids, such as progesterone and androstenedione, glucose-6-phosphate dehydrogenase (G-6-PD), a key enzyme of the pentose-phosphate pathway which supplies NADPH (TPNH) for steroid hydroxylations, diaphorases (both NAD and NADP), flavoproteins which link the NAD or NADP-linked dehydrogenases to the electron transfer system and oxidize the reduced NAD or NADP for their reuse in steroid hydroxylations (Baillie et al., 1966; Kime, 1987). The localization of these enzymes provides a highly reliable procedure to identify the steroidogenic potential of a given tissue (Lofts and Bern, 1972). The teleost testes are unique in that they can convert androgens into watersoluble glucuronides (Ozon, 1972; Kime, 1980, 1987; Fostier et al., 1983). During glucuronidation, uridine diphosphoglucose dehydrogenase (UDPGD) catalyzes the conversion of UDP- glucose into UDP-glucuronic acid which supplies the glucuronyl group to different hydroxylated compounds, such as steroids (Dutton, 1980). The presence of the enzyme is taken as a histochemical index of glucuronidation (Resink *et al.*, 1987; Van den Hurk *et al.*, 1987a, b).

Accessory sex organs in the form of seminal vesicles (SV), testicular glands and testicular blind pouches are present in catfishes (Siluriformes), and gobies and blennies (Perciformes). Anatomical studies have shown that the SV lobules are embedded in a connective tissue stroma which contains interstitial cells, fibroblasts, blood capillaries, and nerve elements (Van den Hurk et al., 1987a; Fishelson et al., 1994) as in the testicular glands of blennies (Seiwald and Patzner, 1989; Lahnsteiner et al., 1990). In previous studies, 3B-HSD activity was reported in SV interstitial cells of Heteropneustes fossilis (Nayyar and Sundararaj, 1969) and Clarias gariepinus (Van den Hurk et al., 1987a, b) and testicular glands of blennies (Lahnsteiner et al., 1990). In contrast, in urohaze-goby, the enzyme activity was reported in the SV epithelial cells (Asahina et al., 1989). In C. gariepinus, the SV epithelial cells showed UDPGD and G-6-PD activities (Van den Hurk et al., 1987a, b). Biochemical studies in this species further showed that the SV is capable of both steroid biosynthesis and glucuronide formation (Schoonen and Lambert, 1986,1987). In view of these information, a histochemical study was conducted to demonstrate the steroidogenic and glucuronidation sites in the SV of the catfish C. batrachus and the results were compared with that of the testis.

^{*} Corresponding author: Tel. +91-542-317325;

FAX. +91-542-317074.

MATERIALS AND METHODS

Collection and maintenance of fish

Male adult *Clarias batrachus* weighing 60 ± 5 g were collected from local fish markets in and around Varanasi in resting (January), late preparatory (April), early prespawning (July) and postspawning (October) phases. They were maintained in the laboratory under natural conditions and fed goat liver daily.

Tissue processing

For demonstrating the spermatogenetic stages, testes of resting (January), late preparatory (April) and late prespawning (June) and postspawning (October) fish were fixed in Bouin's fluid. In the late preparatory phase, pieces of the SV were also fixed in Bouin's fluid. Paraffin sections (7.5 μ m) were stained with Ehrlich's hematoxylineosin. For histoenzymology, SVs and testes of late preparatory fish were dissected out and quickly frozen. Fresh cryostat sections (10-15 μ m) were taken in the transverse plane and mounted on clean slides and air-dried. The sections were utilized for localization of various enzymes.

a. Δ^5 , 3β-hydroxysteroid dehydrogenase (3β-HSD) activity was localized according to the method of Baillie *et al.* (1966) using dehydroepiandrosterone (DHA; Sigma) as the substrate. The medium was prepared fresh at the time of incubation as follows: 10 mg nicotinamide adenine dinucleotide (NAD), 10 mg ethylene diamine tetraacetic acid disodium salt (EDTA), and 10 mg nitroblue tetrazolium (NBT) were dissolved in 8.0 ml 0.2 M phosphate buffer (pH 7.4). DHA was dissolved separately in 2.0 ml dimethyl formamide (DMF). Both the solutions were then mixed and used for the incubation. Sections were incubated at 37°C for 20 min. As a control, some sections were incubated in a substrate-free medium for checking the specificity of the reaction.

b. Reduced nicotinamide adenine dinucleotide (NADH) diaphorase: The diaphorase activity was localized according to the method of Baillie *et al.* (1966) using NADH (Sigma) as the substrate. The incubation medium was prepared fresh at the time of incubation as follows: 5 mg NBT and 5 mg NADH were dissolved in 10 ml 0.2 M phosphate buffer (pH 7.4). Cryostat sections were incubated at 37°C for 5 min. As a control, a few sections were incubated in a substrate-free medium.

c. Glucose-6-phosphate dehydrogenase (G-6-PD): G-6-PD activity was localized by the method of Cohen (1959), using G-6-P (Sigma) as the substrate. The medium was prepared at the time of incubation as follows: 5 mg NADP (NAD phosphate), 5 mg NBT, 5 mg EDTA, 10 mg G-6-P were dissolved in 10 ml 0.2 M phosphate buffer (pH 7.4). Frozen sections were incubated at 37°C for 20 min. As a control, a few sections were incubated in a substrate-free medium. **d. Uridine diphosphoglucose dehydrogenase (UDPGD):**

a. Uridine diprosphoglucose denyarogenase (UDPGD): UDPGD activity was localized by the method of Jacobsen and Jørgensen (1973). The incubation medium was prepared by dissolving 0.82 mM uridine-5'-diphosphoglucose sodium salt (Sigma), 2.15 mM EDTA, 0.24 mM NBT, 0.75 mM NAD and 5% polyvinyl alcohol in 0.02 M Tris- HCl buffer, pH 8.3. The incubation was carried out for 1 hr at 30°C. As a control, sections were also incubated in a substratefree medium.

After the incubations, sections were fixed in 15% formol-saline for 15 min. They were washed in tap water, rinsed in distilled water and mounted in glycerine-jelly.

RESULTS

Spermatogenetic cycle

In *C. batrachus*, the testis and SV show correlated seasonal activity. The testicular cycle can be divided into 5 phases. **Resting or quiescent phase:** During this phase (November- January), the testis is thin and filament-like (Fig. 1). Histologically, it consists of several small undifferentiated seminiferous cords containing spermatogonia and spermatogonial mother cells embedded within thick interstitium. Interstitial cells are embedded in the connective tissue.

Preparatory phase: In this phase (February-April), the size of the testis increases progressively (Figs. 2 and 4). The seminiferous tubules increase in size with a reduction in the size of the interstitium. Spermatogenic cysts were seen in different stages of meiosis with numerous division figures. The lumen is filled with spermatids and spermatozoa. Sertoli cells are clearly visible (Fig. 4).

Prespawning phase: In this phase (May-June), the wave of spermatogenesis continues and reaches the peak. The tubules are greatly distended and the testis is enlarged. This stage is characterized by the abundance of spermatozoa in the lumen (Fig. 3). The germinal epithelium contains several cysts of actively dividing spermatocytes. The interstitial cells are numerous and active.

Spawning phase: In this phase (July-August), the testis becomes totally opaque, fragile and attains maximum size. The interstitial cells are active and numerous. The lumen is filled with spermatozoa or empty due to release of the sperm. **Postspawning phase:** In this phase (September-October), the testis is almost empty and spent. The germinal epithelium shows residual spermatogenic activity, giving rise to a second crop of spermatozoa.

In the SV, interstitial cells are present in groups or singly between the lobules and in the periphery close to blood capillaries (Fig. 5). They are round or oval in shape and are comparable in distribution and morphology to the Leydig cells of the testis.

Localization of 3 β -HSD, G-6-PD, NADH diaphorase, and UDPGD activities: In both testis and SV, the interstitial cells showed strong reaction for 3 β -HSD (Figs. 6 and 7), G-6-PD (Figs. 8 and 9), and NADH diaphorase (Figs. 10 and 11) activities, as indicated by purple-blue diformazan deposits. The interstitial cells of both testis and SV (Figs. 12-14) showed strong UDPGD activity. The germinal epithelium of the testis and the secretory epithelium of the SV also showed moderate to strong enzyme activity. In the testis, the Sertoli cells seemed to be strongly reactive than the germ cells (Fig. 13). In control sections, which were incubated without the respective substrates, diformazan deposits were not formed indicating the specific nature of the reactions.

DISCUSSION

In the present study, the interstitial (Leydig) cells of the testis showed strong 3β-HSD activity, as has been reported by Kirubagaran and Joy (1988). In several teleosts, the Leydig cells are the main sites of steroidogenesis: *Tilapia mossambica* (Yaron, 1966); *Cymatogaster aggregata* (Wiebe, 1969); *Carassius auratus* (Yamazaki and Donaldson, 1969); *Poecilia reticulata* and *Oryzias latipes* (Takahashi and Iwasaki, 1973a, b); *Salmo gairdneri* (Van den Hurk *et al.*, 1978; reviews:



Figs. 1-5. Histology of the testis of *Clarias batrachus*. Spermatocytes (SC), Spermatids (ST). Leydig cells (L). Ehrlich's hematoxylin - eosin. **Fig. 1.** A transverse section (TS) of the testis in the resting phase (January). Note a large number of spermatogonia (thick arrows) and differentiating seminiferous cords. Interstitium is thick. × 240.

Fig. 2. A TS of the testis in early preparatory phase (March) showing well developed and distended seminiferous tubules with spermatocytes and numerous spermatids and a few spermatozoa. × 240.

Fig. 3. A TS of the testis in late prespawning phase (June). Note the tubules filled with spermatozoa. Some tubules still contain spermatids. \times 240.

Fig. 4. A TS of the testis of *C. batrachus* in late preparatory phase (April) showing interstitial (Leydig) cells (arrows), Sertoli cells (arrow-heads) and blood capillaries (C). \times 384.

Fig. 5. A TS of the seminal vesicle of *C. batrachus* in late preparatory phase (April) showing interstitial cells (thick arrows) and blood capillaries (C). × 925.



Figs. 8 and 9. Frozen sections of the testis and SV of *C. batrachus* in late preparatory phase (April) showing intense G-6-PD activity in the interstitial cells of the testis (Fig. 8) and SV (Fig. 9). × 240. Figs. 10 and 11. Frozen sections of the testis and SV of *C. batrachus* in late preparatory phase (April) showing NADH diaphorase activity in the interstitial cells of the testis (Fig. 10) and SV (Fig. 11). × 240. indicating strong 3 β -HSD activity. \times 240.



Figs. 12 and 13. Frozen sections of the testis of *C. batrachus* in late preparatory phase (April) showing strong UDPGD activity in the interstitial cells (arrows) of the testis. Note strong reaction in the germinal epithelium, most likely in Sertoli cells (arrow-heads). \times 80; \times 384. **Fig. 14.** Frozen section of the SV of *C. batrachus* in late preparatory phase (April) showing UDPGD activity in the SV epithelium and interstitial cells (arrows). \times 240.

Fostier *et al.*, 1983; Nagahama, 1983). The interstitial cells showed strong reaction for G-6-PD and NADH diaphorase activities, providing additional histochemical evidence for steroidogenic activity. The teleost testis is an important site of steroid glucuronidation (Colombo *et al.*, 1977; Kime, 1980; Resink *et al.*, 1987; Van den Hurk *et al.*, 1987c). It has been reported that the presence of UDPGD activity indicates sites of steroid glucuronide synthesis (Van den Hurk *et al.*, 1987a,

b). These authors showed that the interstitial cells showed weak enzyme activity in the testis of the African catfish. In contrast, a strong enzyme activity was reported in the interstitial cells of the testis in *C. batrachus*. Furthermore, the germinal epithelium showed moderate enzyme activity with Sertoli cells perhaps showing intense activity. Thus, the combined presence of UDPGD and 3β -HSD activities in the interstitial and Sertoli cells suggests glucuronidation of steroids in these

sites. The strong reaction of UDPGD activity in *C. batrachus* may be due to the fact that they were collected from the wild. Van den Hurk *et al.* (1987b) observed that the enzyme activity was weak in captive fish which was also supported by subsequent biochemical studies (Schoonen and Lambert, 1987; Schoonen *et al.*, 1987a). The moderate to strong UDPGD activity noticed in the germinal epithelium suggests perhaps incorporation of UDP-glucuronic acid into polysaccharides or glucuronidation of nonsteroidal compounds, as has been suggested in vitellogenic follicles (Van den Hurk *et al.*, 1987b).

The interstitial cells of the SV of C. gariepinus and testicular glands of blennies show ultrastructural and histoenzymological characteristics of steroid-producing cells and are homologised to the Leydig cells (Van den Hurk et al., 1987a; Lahnsteiner et al., 1990). In C. batrachus they showed intense reaction for 3β-HSD, G-6-PD, NADH diaphorase and UDPGD activities. The presence of 3β-HSD activity was reported in the SV of *H. fossilis* (Nayyar and Sundararaj, 1969), C. gariepinus (Van den Hurk et al., 1987a, b), and blennies (Lahnsteiner et al., 1990). In contrast, Asahina et al. (1989) reported 3β-HSD activity only in the epithelial cells of the SV of urohaze-goby. Van den Hurk et al. (1987a) reported G-6-PD activity in the epithelial cells of the SV along with other enzymes, the activity being proportionate to the height of the epithelium. It was weak in squamous cells indicating decreased cellular activity. In C. batrachus, UDPGD activity was noticed both in the interstitial and epithelial cells. In C. gariepinus, the epithelial cells exhibited strong enzyme activity (Van den Hurk et al., 1987a, b). According to these workers, the interstitial cells of the SV of specimens obtained from wild only showed the enzyme activity. The occurrence of UDPGD along with 3β-HSD activity in the interstitial cells of the SV suggests that these cells synthesize steroid glucuronides. In testicular glands of Gobius niger, both 3β-HSD and UDPGD activities were localized in the same cells (Seiwald and Patzner, 1989). Apparently, steroid glucuronides are not produced in the testicular glands of blennies (Lahnstenier et al., 1990). The presence of UDPGD activity in the epithelial cells suggests two possibilities: (1) the steroids are taken up from the interstitial cells, converted into steroid glucuronides, and then released into the lumen, (2) incorporation of UDP-glucuronic acid into polysaccharides or nonsteroidal compounds secreted by the epithelium (Van den Hurk et al., 1987a, b). Biochemical studies have shown that the SV of C. gariepinus produces a large number of steroids and steroid glucuronides (Schoonen and Lambert, 1987). Studies of Schoonen and Lambert (1987) and Schoonen et al. (1987a, b) have shown that the SV seems to be more important in steroid glucuronide production, judging from its greater capacity (than testis) to synthesize them and from the greater diversity of glucuronides formed in it. They opined that, in those species in which the SV is present, the functions normally performed by the testis are divided between them; the steroid glucuronides being synthesized mostly in the SV. The present histochemical localization of UDPGD activity both in the testis and SV suggests that steroid glucuronide formation may occur in both the structures but quantitative differences, if any, in glucuronide formation can be known only by biochemical analysis. The steroid glucuronides of the SV origin are implicated in pheromonal roles (Stacey *et al.*, 1987; Van Weerd, 1990; Stacey and Sorensen, 1991; Van den Hurk and Resink, 1992). In the African catfish, steroid glucuronides of the SV serve as olfactory stimuli to stimulate ovulation (Resink *et al.*, 1989).

A comparison of the distribution of the different enzymes in the interstitial cells of the SV and testis shows that both structures are steroidogenic and are capable of steroid glucuronidation, indicating a common origin from the gential ridge. The homology between the two structures is further supported by other evidences. They are: (i) correlative changes in histology and some biochemical constituents (proteins, fructose, hexosamines, sialic acid), (ii) presence of scattered spermatogenic cysts in different stages of development in the SV and SV-testis interphase, (iii) extirpation of testicular component (hemicastration) leads to hyperactivity of the SV in terms of increase in size, testosterone production and secretory activity of the epithelium, and (iv) similarities in androgen and cyproterone acetate responses (Rai, 1996). The anterior part of the gonad is differentiated for a genital function (testis) and the posterior part for a secretory function with some residual spermatogenic activity (SV). Both discharge their contents through the efferent duct systems - the testis into the lateral sperm ducts and SV into the common sperm duct-to the exterior. Apart from the secretory function (both exocrine and endocrine), the SV is used for temporary sperm storage which has adaptive value under extreme drought conditions. The fish survives in moist mud burrows and sperm is protected in the SV for ensuing spawning.

REFERENCES

- Asahina K, Suzuki K, Hibiya T, Tamaoki B (1989) Structure and steroidogenic enzymes of the seminal vesicle of the urohaze-goby (*Glossogobius olivaceus*). Gen Comp Endocrinol 74: 385–391
- Baillie AH, Ferguson MM, Hart DMCK (1966) Developments in Steroid Histochemistry. Academic Press, New York
- Cohen RB (1959) Histochemical localization and metabolic significance of glucose-6-phosphate dehydrogenase system in adrenal cortex. Proc Soc Exp Biol Med 101: 405–407
- Colombo L, Colombo-Belvédère P, Pilati A (1977) Biosynthesis of free and conjugated 5β-reduced androgens by testes of the black goby, *Gobius jozo* (L). Boll Zool 44: 131–134
- Dutton GJ (1980) Glucuronidation of Drugs and Other Compounds. CRC Press, Boca Raton, Florida
- Fishelson L, Van Vuren JHJ, Tyran A (1994) Ontogenesis and ultrastructure of seminal vesicles of the catfish, *Clarias gariepinus*. J Morph 219: 59–71
- Fostier A, Jalabert B, Billard R, Breton B, Zohar Y (1983) The gonadal steroids. In "Fish Physiology Vol IX A" Ed by Hoar WS, Randall DJ and Donaldson EM, Academic Press, New York, pp 277–372
- Guraya SS (1976) Recent advances in the morphology, histochemistry, and biochemistry of steroid-synthesizing cellular sites in the testes of nonmammalian vertebrates. Int Rev Cytol 47: 99–136
- Jacobsen WO, Jørgensen F (1973) Further enzyme histochemical observations on the segmentation of the proximal tubules in the kidney of the male rat. Histochemistry 34: 11–32

- Kime DE (1980) Androgen biosynthesis by testis of the goldfish, *Carassius auratus, in vitro*: the effect of temperature on the formation of steroid glucuronides. Gen Comp Endocrinol 41: 164– 172
- Kime DE (1987) The steroids. In "Fundamentals of Comparative Vertebrate Endocrinology" Ed by Chester-Jones I, Ingleton PM and Phillips JG, Plenum Press, New York, pp 3–56
- Kirubagaran R, Joy KP (1988) Inhibition of testicular 3β-hydroxy-Δ⁵steroid dehydrogenase (3β-HSD) activity in the catfish *Clarias batrachus* (L.) by mercurials. Indian J Expt Biol 26: 907–908
- Lahnsteiner F, Richtarski U, Patzner RA (1990) Functions of the testicular gland in two blenniid fishes, *Salaria* (= *Blennius*) *pavo* and *Lipophrys* (= *Blennius*) *dalmatinus* (Blenniidae, Teleostei) as revealed by electron microscopy and enzyme histochemistry. J Fish Biol 37: 85–97
- Lofts B (1987) Testicular function. In "Hormones and Reproduction in Fishes, Amphibians, and Reptiles" Ed by Norris DO and Jones RE, Plenum Press, New York, pp 283–325
- Lofts B, Bern HA (1972) The functional morphology of steroidogenic tissues. In "Steroids in Nonmammalian Vertebrates" Ed by Idler DR, Academic Press, New York, pp 37–125
- Nagahama Y (1983) The functional morphology of teleost gonads. In "Fish Physiology Vol IX A" Ed by Hoar WS, Randall DJ and Donaldson EM, Academic Press, New York, pp 223–275
- Nayyar SK, Sundararaj BI (1969) Effects of pituitary hormones, and drogens, and corticosteroids on the seminal vesicles of the castrate catfish, *Heteropneustes fossilis* (Bloch). J Expt Zool 172: 385–398
- Ozon R (1972) Androgens in fishes, amphibians, reptiles, and birds. In "Steroids in Nonmammalian Vertebrates" Ed by Idler DR, Academic Press, New York, pp 328–389
- Rai MS (1996) Reproductive Endocrinology of Catfish: Aspects of Structure and Physiology of Seminal Vesicle in *Clarias batrachus* (L.). PhD Thesis, Banaras Hindu University, Varanasi, India
- Resink JW, Van den Hurk R, Voorthuis PK, Terlou M, De Leeuw R, Viveen WJAR (1987) Quantitative enzyme histochemistry of steroid and glucuronide synthesis in testes and seminal vesicle, and its correlation to plasma gonadotropin level in *Clarias gariepinus*. Aquaculture 63: 97–114
- Resink JW, Voorthuis PK, Van den Hurk R, Peters RC, Van Oordt PGWJ (1989) Steroid glucuronides of the seminal vesicle as olfactory stimuli in African catfish, *Clarias gariepinus*. Aquaculture 83: 153–166
- Schoonen WGEJ, Lambert JGD (1986) Steroid metabolism in the seminal vesicles of the African catfish, *Clarias gariepinus* (Burchell), during the spawning season, under natural conditions, and kept in ponds. Gen Comp Endocrinol, 61: 355–367
- Schoonen WGEJ, Lambert JGD (1987) Gas chromatographic-mass spectrometric analysis of steroids and steroid glucuronides in the seminal vesicle fluid of the African catfish, *Clarias gariepinus*. Gen Comp Endocrinol, 68: 375–386
- Schoonen WGEJ, Granneman JCM, Lambert JGD, Van Oordt PGWJ (1987a) Steroidogenesis in the testes and seminal vesicles of spawning and non-spawning African catfish, *Clarias gariepinus*. Aquaculture 63: 77–78

- Schoonen WGEJ, Lambert JGD, Resink JW, Viveen WJAR, Van Oordt PGWJ (1987b) A quantitative study of steroid bioconversions in the testes of the African catfish, *Clarias gariepinus* (Burchell), under natural spawning and natural and cultivated non-spawning conditions. J Endocrinol 112: 323–332
- Seiwald M, Patzner RA (1989) Histological, fine-structural and histochemical differences in the testicular glands of gobiid and blenniid fishes. J Fish Biol 35: 631–640
- Stacey N, Sorensen P (1991) Function and evolution of fish hormonal pheromones. In "Biochemistry and Molecular Biology of Fishes Vol I" Hochachka PL and Mommsen TP, Elsevier, Toronto, pp 109–135
- Stacey NE, Sorensen PW, Dulka JG, Van der Kraak GJ, Hara TJ (1987) Teleost sex pheromones: recent studies on identity and function. In "Proceedings of the Third International Symposium on Reproductive Physiology of Fish" Ed by Idler DR, Crim LW and Walsh JM, Memorial University Press, St John's, Canada, pp 150–153
- Takahashi H, Iwasaki Y (1973a) The occurrence of histochemical activity of 3β -hydroxysteroid dehydrogenase in the developing testes of *Poecilia reticulata*. Dev Growth Different 15: 241–253
- Takahashi H, Iwasaki Y (1973b) Histochemical demonstration of Δ⁵-3β-hydroxysteroid dehydrogenase activity in the testis of the medaka, *Oryzias latipes*. Endocrinol Japon 20: 529–534
- Van den Hurk R, Vermeij JAJ, Stegenga J, Peute J, Van Oordt PGWJ (1978) Cyclic changes in the testis and vas deferens of the rainbow trout (*Salmo gairdneri*) with special reference to sites of steroidogenesis. Ann Biol Anim Biochim Biophys 18: 899–904
- Van den Hurk R, Resink JW, Peute J (1987a) The seminal vesicle of the African catfish, *Clarias gariepinus*. A histological, histochemical, enzyme-histochemical, ultrastructural and physiological study. Cell Tissue Res 247: 573–582
- Van den Hurk R, Resink JW, Voorthuis PK (1987b) An enzyme histochemical study concerning the localization of steroid glucuronide production in the reproductive organs of African catfish, *Clarias gariepinus*. Aquaculture 63: 89–96
- Van den Hurk R, Schoonen WGEJ, Van Zoelen GA, Lambert JGD (1987c) The biosynthesis of steroid glucuronides in the testis of the zebrafish, *Brachydanio rerio*, and their pheromonal function as ovulation inducers. Gen Comp Endocrinol 68: 179–188
- Van den Hurk R, Resink JW (1992) Male reproductive system as sex pheromone producer in teleost fish. J Exp Zool 261: 204–213
- Van Weerd JH (1990) Pheromones and Ovarian Growth in the African Catfish *Clarias gariepinus*. Ph D Thesis, Agricultural University, Wageningen, The Netherlands
- Wiebe JP (1969) Steroid dehydrogenases and steroids in gonads of the seaperch *Cymatogaster aggregata* Gibbons. Gen Comp Endocrinol 12: 256–266
- Yamazaki F, Donaldson EM (1969) Involvement of gonadotropin and steroid hormones in the spermiation of the goldfish (*Carassius auratus*). Gen Comp Endocrinol 12: 491–497
- Yaron Z (1966) Demonstration of 3β-hydroxysteroid dehydrogenase in the testis of *Tilapia mossambica*. J Endocrinol 34: 127–128

(Received July 16, 1997 / Accepted July 10, 1998)