

Evolution of Hypervariable Regions, V4 and V7, of Insect 18S rRNA and Their Phylogenetic Implications

Authors: Hwang, Ui Wook, Ree, Han Il, and Kim, Won

Source: Zoological Science, 17(1) : 111-121

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.17.111>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Evolution of Hypervariable Regions, V4 and V7, of Insect 18S rRNA and Their Phylogenetic Implications

Ui Wook Hwang^{1,2}, Han Il Ree² and Won Kim^{1*}

¹*Department of Molecular Biology, Seoul National University, Seoul 151-742, Korea and*

²*Department of Parasitology, Yonsei University College of Medicine,
Seoul 120-752, Korea*

ABSTRACT—We compared primary and secondary structures of V4 (helices E23-2 to E23-5) and V7 (helix 43) regions of 18S rRNAs in insects and the other three major arthropod groups (crustaceans, myriapods, and chelicerates) known so far. We found that the lengths of primary sequences and the shapes of secondary structures of these two hypervariable regions of insect 18S rRNA even at infraclass levels are phylogenetically informative and reflect major steps in insect evolution. The long sequence insertion and bifurcated shape of helices E23-2 to E23-5 in the V4 region are unique synapomorphic characters for winged insects (Pterygota). The long sequence insertion and expanded stem length of helix 43 in the V7 region are synapomorphic characters for holometabolous insects which conduct complete metamorphosis. The strongly conserved secondary structures suggest the possibility that these hypervariable regions may be related with certain important cellular functions unknown thus far. The comparison with insect fossil records revealed that the pterygote synapomorphy (V4) and the holometabolous synapomorphy (V7) were established prior to the acquisition of insect wings (flight system) and prior to the development of complete metamorphosis, respectively. These synapomorphies have been also relatively stable over at least 300 Myr and 280 Myr, respectively as well. It implies that the expansion events of the V4 and V7 regions have not occurred simultaneously but independently at different periods during the insect evolution. Then this suggests that V4 and V7 regions are not functionally correlated as recently suggested by Crease and Coulbourn.

INTRODUCTION

The slowly evolving property of 18S rRNA sequences has been widely used for phylogenetic studies among remotely related animal groups such as among phyla, classes, and orders. In the phylogenetic studies of major arthropods and related groups, a number of authors have also used primary sequence information from slowly evolving parts of the 18S rDNA (Carmean *et al.*, 1992; Pashley *et al.*, 1993; Campbell *et al.*, 1994; Friedrich and Tautz 1995; Kim *et al.*, 1996; Chalwatzis *et al.*, 1996; Giribet *et al.*, 1996; Friedrich and Tautz, 1997). The fast evolving regions (especially V4 and V7) have been excluded in those analyses because of difficulties in obtaining a reliable alignment, of difficulties in constructing an unambiguous secondary structure and of saturation of phylogenetic informations on nucleotide sequences due to multiple hit. Only since recently the fast evolving parts of 18S rDNA have been used for the phylogenetic studies among close relatives at lower catego-

ries such as the family level in tiger beetles (Volger and Pearson, 1996; Volger *et al.*, 1997; Hancock and Volger, 1998).

We had previously determined the 18S rDNA sequences from a number of collembolan species (Lee *et al.*, 1995a, b; Hwang *et al.*, 1995). We found that the sequence length of the collembolan 18S rDNA is far shorter than those of dipteran insects and the differences of the sequence lengths are mainly caused by expansions of V4 and V7 regions in dipteran insects (Hwang *et al.*, 1995). This finding made it possible to deduce that primary and secondary structures of these two variable regions may provide us with some critical information related to insect phylogeny and 18S rRNA evolution. Recently, Crease and Coulbourn (1998) have reported that the coordinated and perhaps functionally correlated increases occur between V4 and V7 regions of many arthropod 18S rRNAs.

The phylogenetic relationships among major subgroups of insects were relatively well documented on the basis of morphological and paleontological characters (Kristensen 1991, Kukalová-Peck, 1991). In addition, insect phylogeny was also examined and discussed on the basis of molecular data such as the alignable sequences of 18S and 28S

* Corresponding author: Tel. +82-2-880-6695;
FAX. +82-2-872-1993.
E-mail: wonkim@plaza.snu.ac.kr

rRNAs (Chalwatzis *et al.*, 1996, Whiting *et al.*, 1997). Yet, in the molecular studies, variable regions of 18S rRNA have never been employed for phylogenetic studies on higher categorical levels (above Order) of insects so far. In this paper, we conduct comparative analyses of primary and secondary structures of two hypervariable regions, V4 and V7, of 18S rRNA. We suggest that these regions have independently evolved during the insect evolution and could provide phylogenetic informations in higher categorical levels (above Order) of insects.

MATERIALS AND METHODS

All arthropod 18S rRNA sequences accessible from EMBL data bank were retrieved and examined. At least one or more 18S rDNA sequences from most major insect orders have been published. Because primary and secondary structures of 18S rRNAs are similar within each insect order, one representative species in each insect order is selected (except for Collembola and Diptera; 3 and 2 species, respectively) and their sequence alignments and the secondary structures are presented in this paper. Two representative 18S rRNA sequences from the other major arthropod groups (crustaceans, chelicerates, and myriapods) are also shown as reference groups. However, the extreme cases of sequence expansion obtained from tiger beetles (Volger *et al.*, 1997), strepsipteran species such as *Xenos vesparum*, *Mengenilla chobauti*, and *Stylops melittae* (Chalwatzis *et al.*, 1995), pea aphid, *Acyrtosiphon*

Table 1. List of representative arthropod species employed in this analysis and the abbreviations

Classification	Species	Source	Abbreviation
Subphylum Hexapoda			
Class Ellipura ^{a)d)}			
Order Collembola	<i>Podura aquatica</i>	X89485 (V4 & V7)	Col1
	<i>Hypogastrura dolsana</i>	Z26765 (V4 & V7)	Col2
	<i>Crossodonthina koreana</i>	Z36893 (V4 & V7)	Col3
Class Insecta ^{e)}			
Subclass Archaeognatha ^{a)}			
Order Archaeognatha	<i>Petrobius brevistylis</i>	X89808 (V4)	Arc
Subclass Dicondylia			
Infraclass Thysanura ^{a)}			
Order Thysanura	<i>Lepisma saccharina</i>	X89484 (V4 & V7)	Thy
Infraclass Pterygota			
Division Ephemeroptera ^{b)c)}			
Order Ephemeroptera	<i>Ephemera</i> sp.	X89489 (V4 & V7)	Eph
Division Odonata ^{b)c)}			
	<i>Aeschna cyanea</i>	X89482 (V4 & V7)	Odo
Division Neoptera			
Order Dermaptera ^{c)}	<i>Labidura riparia</i>	U65114 (V4)	Der
Order Orthoptera ^{c)}	<i>Melanoplus</i> sp.	U65115 (V4)	Ort
Order Hemiptera ^{c)}	<i>Philaenus spumarius</i>	U06480 (V4 & V7)	Hem
Subdivision Holometabola			
Order Hymenoptera	<i>Polistes dominulus</i>	X74762 (V4 & V7)	Hym
Order Coleoptera	<i>Tenebrio molitor</i>	X07801 (V4 & V7)	Ten
Order Lepidoptera	<i>Galleria mellonella</i>	X89491 (V4 & V7)	Lep
Order Siphonaptera	<i>Archaeopsylla erinacea</i>	X89486 (V4 & V7)	Sip
Order Diptera	<i>Drosophila melanogaster</i>	M21017 (V4 & V7)	Dip1
	<i>Aedes albopictus</i>	X57172 (V4 & V7)	Dip2
Subphylum Myriapoda			
Class Chilopoda			
Order Lithobiomorpha	<i>Lithobius forficatus</i>	X90653 (V4)	Myr1
	<i>Bothropylus asperatus</i>	Ref* (V7)	Myr3
Class Diplopoda			
Order Chilognatha	<i>Megaphyllum</i> sp.	X90658 (V4)	Myr2
Subphylum Crustacea			
Class Branchiopoda			
Order Anostraca	<i>Artemia salina</i>	X10723 (V4 & V7)	Cru1
Class Maxillopoda			
Subclass Branchiura			
Order Arguloida	<i>Argulus nobilis</i>	M27187 (V4 & V7)	Cru2
Subphylum Chelicerata			
Class Arachnida			
Order Araneae	<i>Eurypelma californica</i>	X13457 (V4 & V7)	Che1
Class Merostomata			
Order Xiphosura	<i>Limulus polyphemus</i>	X90467 (V4)	Che2

a) Apterygota b) Paleoptera c) Hemimetabolous insect d) Entognathous insect e) Ectognathous insect

*Ref: The sequence was obtained from the paper of Min *et al.* (1998).

pisum (Kwon *et al.*, 1991), the branchiopod crustacean *Daphnia pulex* (Crease and Colbourne, 1998), and the isopod crustacean *Armadillidium vulgare* (Choe *et al.*, 1999a) are not included in the alignment set due to the difficulties for constructing their stable secondary structures. We will discuss these exceptional cases in detail in another subsequent paper. The classification, the representative species names, EMBL accession numbers, and abbreviations of taxon names are listed in Table 1. The classification scheme is followed by Kristensen (1991) for the Hexapoda, by Brusca and Brusca (1990) for the Crustacea, and by Hickman *et al.* (1984) for the Chelicerata and the Myriapoda.

The sequences of V4 and V7 regions of the 18S rDNAs from the 22 and 17 species (Table 1) respectively were aligned by Clustal X (Thompson *et al.*, 1997). Then the alignments of the primary sequences of helices E23-2 to E23-5 of the V4 region and helix 43 of the V7 region were adjusted by the observation of compensatory substitutions in our predicted secondary structure model. The nomenclature of these helices is after Neefs *et al.* (1993) and the positions are indicated in the putative secondary structure of 18S rRNA of *Hypogastrura dolsana* (order Collembola) for convenience (Fig. 1A).

In the present study, our putative secondary structure of helices




Fig. 1. (A) Putative 18S rRNA secondary structure of *Hypogastrura dolsana*, order Collembola. It was constructed by hand after Rijk's *D. melanogaster* model (Rijk *et al.*, 1992). The two arrows indicate the stem-loop structures used for comparative analyses of primary and secondary structures; the helices E23-2 to E23-5 in the V4 region and the helix 43 in the V7 region, respectively. The helices are numbered according to the method of Neefs *et al.* (1993). These two regions have the highest sequence variability in insect 18S rDNA. Based on this complete secondary structure, the helix E23-2 and the basal part of the helix 43 were determined as anchored pairings for predicting putative secondary structures in the present comparative analyses. The bold lines mark the two regions of anchored pairings. The entire regions of the V4 and the V7 are shown in boxes. (B) Secondary structures of the helices E23-2 to E23-5 in *D. melanogaster* 18S rRNA derived from three different secondary structure models; Rijk *et al.*'s model (Rijk *et al.*, 1992), Hancock *et al.*'s model (Hancock *et al.*, 1988), and our present model. In the Rijk *et al.*'s model, they were originally designated the helices E21-2 to E21-5. However, the helices were renamed helices E23-2 to E23-5 by Neefs *et al.* (1993). All three models have the helix E23-5 in common, which are marked by thin lines. The anchored pairings (helix E23-2) determined from (A) are shown in bold lines in the Rijk *et al.*'s model and our present model. These secondary structures were folded without any constraints after excluding flanking sequences.


E23-2 to E23-5 in *D. melanogaster* was predicted by using helix E23-2 as an anchored pairing and was compared with two previously published secondary structure models by Rijk *et al.* (1992) and Hancock *et al.* (1988) (Fig. 1B). Our present model is quite different from that of Rijk *et al.* (1992) which is folded into three helices (E23-3, E23-4, and E23-5) from helix E23-2 (anchored pairings). However our model is rather similar to that of Hancock *et al.* (1988). Both models are folded into two helices and the base pairings of stems in these helices are completely the same, though the anchored pairings are absent and their nucleotides are involved in flanking stems or loops. Of the two helices in our secondary structure model (Fig. 1B), the right helix includes helix E23-5 of Rijk *et al.* (1992) and the left one is the same as the hypervariable region discussed in 18S rDNA of tiger beetles by Volger *et al.* (1997). The

model recently revised by Peer *et al.* (1998) agrees well with our present model. The putative secondary structures of helices E23-2 to E23-5 and helix 43 were finally drawn by loopDloop secondary structure drawing software (Gilbert, 1992).

RESULTS

Primary structure analysis

The multiple sequence alignments of arthropod V4 and V7 regions are shown in Fig. 2. The stem and loop regions are indicated according to the Kjer method in the alignments (Kjer, 1995). In this multiple alignment, the



B)





Fig. 2. Multiple alignments of (A) the V4 and (B) the V7 regions of 18S DNA in insects and the other three major arthropod groups. Borders of hairpin structures are bracketed, nucleotides assumed to participate in stem building, where it could be established, are underlined. Species names are abbreviated to the first three letters of the respective order or class name except for the case of "Ten", order Coleoptera (refer to Table 1). Asterisk (*) above the alignments denotes inclusion in comparative analyses of primary and secondary structures. When we predict the putative stem-loop structures of the helices E23-2 to E23-5 and the helix 43, the helix E23-2 and the basal part of the helix 43 are used as anchored pairings, respectively. The approximate regions of anchored pairings are shown in bold lines. The nucleotides used for anchored pairings are written in bold. Dip1' has the same sequences with Dip1 but the depicted secondary structure bases on Rijk's model (Rijk *et al.*, 1992).

sequence positions of V4 and V7 regions correspond to positions 643 to 855 and 1421 to 1608 of 18S rDNA of *Drosophila melanogaster*, respectively (Tautz *et al.*, 1988).

The sequences of the helices E23-2 to E23-5 of the V4 region range from 49 bp to 51 bp (apterygote insects), from 72 bp to 120 bp (pterygote insects), from 46 bp to 48 bp (myriapods), from 47 bp to 55 bp (crustaceans), and from 46 bp to 49 bp (chelicerates). The longest and the shortest sequences are those of *Philaenus spumarius* (Order Hemi-


ptera, Class Insecta, Subphylum Hexapoda) and *Artemia salina* (Order Anostraca, Class Branchiopoda, Subphylum Crustacea), respectively. The sequences of helices E23-2 to E23-5 in pterygote insects are longer than those of apterygote insects and other three major arthropod groups (chelicerates, crustaceans, myriapods). In pterygote insects, Orders of Dermaptera, Orthoptera, and Hemiptera have relatively long sequences compared to the other pterygote insects. In holometabolans of pterygote insects, *D.*

A)

Collembola

Thysanura


Archaeognatha

B)

Ephemeroptera

Hemiptera

Holometabola 1

C)

Collembola

Thysanura

Ephemeroptera

Hemiptera




Fig. 3. Putative secondary structures of the helices E23-2 to E23-5, (A) and (B), and the helix 43, (C) and (D), of 18S rRNAs from representative insects and the other three major arthropods listed in Table 1, except for Col1 and Col3. All these present structures are predicted by the same method used in our present model of Fig. 1B. The sequences of anchored pairings are also shown in Fig. 2. Watson-Crick base pairing is represented by a solid line (e.g. A-U), noncanonical G-U base pairs by a period. The 5'-end and 3'-end are marked. (A) Collembola (*Hypogastrura dolsana*), Thysanura (*Lepisma saccharina*), and Archaeognatha (*Petrobius brevistylis*); (B) Ephemeroptera (*Ephemer* sp.), Hemiptera (*Philaenus spumarius*), and Holometabola (*Drosophila melanogaster*); (C) Collembola (*Hypogastrura dolsana*), Thysanura (*Lepisma saccharina*), Ephemeroptera (*Ephemer* sp.), and Hemiptera (*Philaenus spumarius*); (D) Holometabola 1 (*Drosophila melanogaster*) and Holometabola 2 (*Tenebrio molitor*).

melanogaster has 108 bp that is about 30 bp longer than those of the other holometabolous insects (Fig. 2A).

The sequences of helix 43 of the V7 region range from 52 bp to 59 bp long in apterygote insects, from 55 bp to 66 bp in hemimetabolous insects, from 86 bp to 155 bp in holometabolous insects, and from 49 bp to 53 bp in crustaceans. Those of *Eurypelma californica*, a chelicerate and *Bothropylis asperatus*, a myriapod are 50 bp and 54 bp long, respectively. The longest and the shortest are those of *D. melanogaster* (155 bp) and *A. salina* (49 bp), respectively. The helix 43 of *P. spumarius* (order Hemiptera) is 66 bp long and about 10 bp longer than those of the apterygotes and the paleopterans (Ephemeroptera and Odonata). The sequences of holometabolous insects are longer than those of apterygote insects, hemimetabolous insects (Paleoptera and Hemiptera), and the other three major arthropod groups (Fig. 2B).

Secondary structure analysis

The shapes of the secondary structures of the helices E23-2 to E23-5 are well conserved with taxon-specific patterns as shown in Fig. 3 (A and B). Most of the secondary structures of E23-2 to E23-5 helices in pterygote look like bifurcated forms connected to the anchored pairings (helix E23-2) (Fig. 3B). These secondary structures indicate that

apterygote insects have one more helix compared to apterygote insects (Fig. 3A) and the other three major arthropod groups (data not shown), both of which have one elongated helix connected to anchored pairings. Putative secondary structures of the helix 43 (V7 region) were predicted and compared as shown in Fig. 3 (C and D). The stem lengths of helix 43 of holometabolous insects (Fig. 3D) are much longer compared to those of hemimetabolous insects, apterygote insects (Fig. 3C), and the other major three arthropod groups (data not shown).

DISCUSSION

We had previously reported that two dipteran insects (*D. melanogaster* and *Aedes albopictus*) are clearly distinguished from collembolan insects by their longer sequences in the V4 region (61–76 bp) and in the V7 region (89–104 bp). We also suggested that the comparison of more insect 18S rDNA sequences would give a significance to this apparently taxon-specific pattern (Hwang *et al.*, 1995). In our present study, it revealed that these two expanded/deleted regions corresponded to the helices E23-2 to E23-5 in the V4 and helix 43 in the V7 regions, respectively. Our present multiple alignments derived from most of the major orders of insects reconfirm that in insect groups, expansions of 18S

rRNA appear to be taxon specific mainly in these two hypervariable regions, V4 and V7.

Wheeler (1989) suggested that the ectognathous insects (or perhaps pterygote insects) are defined by a medium-sized sequence insertion in 18S rDNA on the basis of their size variation patterns digested by restriction enzymes XbaI/EcoRI. However, because his analysis did not include Thysanura and Archaeognatha, it was not clear if the observed insertions would define ectognathous insects or pterygote insects. Moreover, due to the limitations of the experiment, it was impossible to determine in which region of the 18S rDNA the insertions appeared. Our present analysis reveals that the sequence length differences observed are caused by the expansions of the helices E23-2 to E23-5 in pterygote insects and of the helix 43 in holometabolous insects rather than in ectognathous insects as a whole.

When Rijk *et al.* (1992) and Neefs *et al.* (1993) listed and named helices specific to eukaryotes, they included helices E23-2, 3, 4, and 5 in the class Insecta (Fig. 1B). At that time, just a few 18S rRNA sequences were available in insect groups and they were limited only to holometabolous insects (e.g. *Apis mellifera*, *Tenebrio molitor*, and a couple of *Drosophila* species). Our present analysis of secondary structure conducted with relatively abundant sequence data shows that among the class Insecta, only the infraclass Pterygota has formed two helices with expanded sequences. Furthermore, the Apterygota which consists of Collembola, Thysanura, and Archaeognatha, have only one elongated helix.

The present analyses of primary and secondary structures of 18S rRNAs are in accordance with the monophyly of the Pterygota and of the Holometabola already established by morphological evidence (Kristensen, 1991). The monophyly of Archaeognatha, Thysanura, and Pterygota (ectognathous insects) has been suggested based on the morphological characters and has been widely accepted, as well (Brusca and Brusca, 1990). However, one intriguing fact from our present study is that Archaeognatha and Thysanura have neither V4 sequence expansion nor one more helix in the helices E23-2 to E23-5 of the V4 as it was for collembolans as well as the other three major arthropod groups (crustaceans, myriapods, and chelicerates). Hence, this is in contrast with the other ectognathous insects like Pterygota, which have two helices in the helices E23-2 to E23-5. Among the ectognathous insects, Archaeognatha and Thysanura have been considered as intermediate-form insects between the development of ectognathous mouthparts and the development of wings. Thus they have shared both primitive morphological characters found in collembolans and advanced morphological characters found in pterygotes. Their collembolan-like single helix possession in the helices E23-2 to E23-5 is one another plesiomorphic character shown in the molecular level.

Our results imply that the bifurcated shape of helices E23-2 to E23-5 is a synapomorphy for the pterygote insects.

The large sequence insertion and the bifurcated form of the secondary structure of this region are present in most of the pterygote insects examined thus far, though there exists one exception, *Hydropsyche* sp. (data not shown; Trichoptera, X89483) which has only a single helix in the helices E23-2 to E23-5 region. Thus it is most parsimonious to assume that the second helix of the *Hydropsyche* case was lost independently in the evolutionary lineage of Trichoptera. Likewise, for the holometabolous insects, the stem elongation of helix 43 is the shared synapomorphic character. The remarkably conserved secondary structures suggest the possibility that these hypervariable regions are related with certain important cellular functions unknown so far.

In the present analysis, we do not include 18S rDNAs of some insect species to evolve fast as well as to have extremely expanded lengths because it is difficult to predict their stable secondary structures. Recently, we reconstructed the secondary structures from strepsipteran insects (Choe *et al.*, 1999b). The result showed that they have unique secondary structures highly deviated from our present general features. It is likely that such deviations shown in the excluded taxa including strepsipteran insects are autapomorphic characters appearing only on each evolutionary lineage.

With regard to the insect phylogeny, the cladistic exploitation of the structural changes in rDNA was pioneered by Wheeler (1989). In our previous publication (Hwang *et al.*, 1998), we have also attempted to interpret the phylogenetic meanings of the structural changes of R1/R2 elements of 28S rDNA, D3 stem of 28S rRNA, and ITS2a within 5.8S rDNA in insect phylogeny. We had concluded then that the yield of structural changes which are informative for higher insect systematics is poor in the 28S and 5.8S rDNA regions. Through our present study, however, it is now revealed that the primary and the secondary structures in V4 and V7 regions of 18S rRNA are phylogenetically informative in higher categorical level of insects and are evidently indicative of major steps in the insect evolution.

On the basis of insect fossil records (Kukalová-Peck, 1991), it can be deduced that the bifurcated form of helices E23-2 to E23-5 was established prior to the acquisition of insect wings (flight system) and has been relatively stable for over at least 300 Myr. Thereafter, the elongation event of helix 43, which has been relatively stable for over at least 280 Myr, was established prior to complete metamorphosis (Fig. 4). It should be noted here that these two structural changes in insects are highly conserved for relatively long period of time. All this seems to indicate that the expansion events of V4 and V7 regions in insect 18S rRNA have not occurred simultaneously but independently at different periods during the insect evolution. Crease and Coulbourn (1998) in their recent publication, have proposed two possibilities in order to explain the coordinated increases of V4 and V7 regions in Arthropoda; one possibility is that the variable regions are functionally correlated, as was suggested for the 28S rDNA (Hancock and Dover, 1988, 1990) and the




Fig. 4. Secondary structure evolution of the helices E23-2 to E23-5 (the V4 region) and the helix 43 (the V7 region) of 18S rRNA. The paleontological record of major insect subclades according to Kukalová-Peck (1991) is superimposed onto the canonical view of insect phylogeny (Kristensen, 1991). The arrows indicate Pterygota and Holometabola clades, respectively. Numbers above branches refer to conspicuous derived character states: (1) Ectognathous mouth part; (2) double articulation of mandible; (3) possession of wings; (4) complete metamorphosis; (5) modification of hind wings to halteres, acquisition of a labellum.

other possibility is that expansion mechanisms selectively operated only within certain variable regions where such changes are able to be tolerated. Our comparative analyses based on the insect fossil records strongly support that the expansions of these two variable regions are not functionally correlated and evolved separately. Without considering the times when expansions of the V4 and the V7 regions happened, it is apt to misinterpret as if lengths of these two regions co-increased simultaneously. Our results show that there exists a time-gap, at least 20 Myr, between expansion events of these two variable regions. This is a relatively long gap of time considering that, since the first pterygote insect emerged, most of the extant insects have evolved only within ca. 50 Myr. If these variable regions were functionally correlated, the expansion events must have occurred nearly at the same time. Therefore, it is likely that the coordinated-

like pattern of V4 and V7 expansions is due to selective operation of expansion mechanism rather than functional correlation.

In conclusion, the primary and the secondary structures in V4 and V7 regions of 18S rDNA are found to be phylogenetically informative and reflect major steps in insect evolution. Highly conserved secondary structures of these two hypervariable regions show that these regions may be in charge of unknown important cellular functions. Their coordinated-like increasing pattern is not caused by functional relationship of these two regions. Considering that hypervariable regions of 18S rDNA have been generally employed for phylogenetic studies on lower hierarchical levels (below Family) or removed before conducting comparative analyses so far, these new view points and findings are quite intriguing. Our present approach shows that the

secondary structures of those fast evolving regions of 18S rRNA are remarkably conserved and can be used in phylogenetic studies on higher hierarchical levels (above Order). In other animals, we could not find similar structural changes that are phylogenetically informative so far. Nevertheless, if more 18S rDNA sequence information for a wider range of taxa is obtained and if the same analyses be conducted and applied to the higher taxonomic levels of other animal groups besides insects, the secondary structures of variable regions of 18S rRNA should also be able to show distinct patterns that are phylogenetically informative along the examined taxa. Furthermore, additional sequence data could be helpful to unravel unknown cellular functions and evolutionary mechanisms of V4 and V7 regions

ACKNOWLEDGEMENTS

We heartily thank Dr. Markus Friedrich, Department of Biological Science, Wayne State University, 5047 Gullen Mall, Detroit, MI48202, for his valuable and critical comment for our manuscript. This study was partially supported by grants from KOSEF in 1995–1998 (95-0401-04-01-3), SRC (97K3-0401-03-03-1) and the Ministry of Education of Korea (Institute for Molecular Biology and Genetics) in 1998. Dr. Ui Wook Hwang was supported by post-doctoral fellowship (Sep. 98–Aug. 99) of KOSEF.

REFERENCES

- Brusca RC, Brusca GJ (1990) Invertebrates. Sinauer, Sunderland, MA
- Campbell BC, Steffen-Campbell JD, Gill RJ (1994) Evolutionary origin of whiteflies (Hemiptera: Sternorrhyncha: Aleyrodidae) inferred from 18S rDNA sequences. *Insect Mol Biol* 3: 73–88
- Carmean D, Kimsey LS, Berbee ML (1992) 18S rDNA sequences and the holometabolous insects. *Mol Phyl Evol* 2: 270–278
- Chalwatzis N, Baur A, Stetzer E, Kinzelbach R, Zimmermann FK (1995) Strongly expanded 18S rRNA gene correlated with a peculiar morphology in the insect order of Strepsiptera. *Zoology* 98: 115–126
- Chalwatzis N, Hauf J, Van de Peer Y, Kinzelbach R, Zimmermann FK (1996) 18S ribosomal RNA genes of insects - primary structure of the genes and molecular phylogeny of the holometabola. *Ann Entomol Soc America* 89: 788–803
- Choe CP, Hancock JM, Hwang UW, Kim W (1999a) Analysis of the primary sequence and secondary structure of the unusually long SSU rRNA of the soil bug, *Armadillidium vulgare*. *J Mol Evol* 49: 798–805
- Choe CP, Hwang UW, Kim W (1999b) Putative secondary structures of unusually long strepsipteran SSU rRNAs and its phylogenetic implications. *Mol Cells* 9: 191–199
- Crease TJ, Colbourne JK (1998) The unusually long small-subunit ribosomal RNA of the crustacean, *Daphnia pulex*: sequence and predicted secondary structure. *J Mol Evol* 46: 307–313
- Friedrich M, Tautz D (1995) Ribosomal DNA phylogeny of the major extant arthropod classes and the evolution of myriapods. *Nature* 376: 165–167
- Friedrich M, Tautz D (1997) An episodic change of rDNA nucleotide substitution rate has occurred at the time of the emergence and radiation of the insect order Diptera. *Mol Biol Evol* 14: 644–653
- Gilbert DG (1992) loopDloop, a Macintosh program for visualizing RNA secondary structure. Published electronically on the internet, available via anonymous ftp to ftp.bio.indiana.edu
- Giribet G, Carranza S, Baguña J, Riutort M, Ribera C (1996) First molecular evidence for the existence of a Tadirgrada + Arthropoda clade. *Mol Biol Evol* 13: 76–84
- Hancock JM, Dover GA (1988) Molecular coevolution among cryptically simple expansion segments of eukaryotic 26S/28S rRNAs. *Mol Biol Evol* 5: 377–391
- Hancock JM, Dover GA (1990) Compensatory slippage in the evolution of ribosomal RNA genes. *Nuc Acid Res* 18: 5949–5954
- Hancock JM, Tautz D, Dover GA (1988) Evolution of secondary structures and compensatory mutations of the ribosomal RNAs of *Drosophila melanogaster*. *Mol Biol Evol* 5: 393–414
- Hancock JM, Volger AP (1998) Modelling the secondary structures of slippage-prone hypervariable RNA regions: the example of the tiger beetle 18S rRNA variable region V4. *Nuc Acid Res* 26: 1689–1699
- Hickman CP, Roberts LS, Hickman FM (1984) Integrated principles of zoology. Times Mirror/Mosby College Publishing, St. Louis, Toronto, Santa Clara
- Hwang UW, Kim W, Tautz D, Friedrich M (1998) Molecular phylogenetics at the Felsenstein zone: approaching the Strepsiptera problem using 5.8S and 28S rDNA sequences. *Mol Phyl Evol* 9: 470–480
- Hwang UW, Lee BH, Kim W (1995) Sequences of the 18S rDNAs from two collembolan insects: shorter sequences in the V4 and V7 regions. *Gene* 154: 293–294
- Kim CB, Moon SY, Gelder SR, Kim W (1996) Phylogenetic relationships of annelid, molluscs, and arthropods evidenced from molecules and morphology. *J Mol Evol* 43: 207–215
- Kjer KM (1995) Use of ribosomal RNA secondary structure in phylogenetic studies to identify homologous positions — an example of alignment and data presentation from the frogs. *Mol Phyl Evol* 4: 314–330
- Kristensen NP (1991) Phylogeny of extant hexapods. In “The insects of Australia Vol. 1” Ed by ID Naumann, Melbourne University Press, Melbourne, pp 125–140
- Kukalová-Peck J (1991) Fossil history and evolution of hexapod structures. In “The insects of Australia Vol. 1” Ed by ID Naumann, Melbourne University Press, Melbourne, pp 141–179
- Kwon OY, Ogino K, Ishikawa H (1991) The longest 18S ribosomal RNA ever known. Nucleotide sequence and presumed secondary structure of the 18S rRNA of the pea aphid, *Acyrtosiphon pisum*. *Eur J Biochem* 202: 827–833
- Lee BH, Hwang UW, Kim W, Park KH, Kim JT (1995a) Systematic position of cave Collembola *Gulgastrura reticulosa* (Insecta) based on morphological characters and 18S rDNA nucleotide sequence analysis. *Memories de Biospeologie* 22: 83–90
- Lee BH, Hwang UW, Kim W, Park KH, Kim JT (1995b) Phylogenetic study of the suborder Arthropleona (Insecta: Collembola) based on morphological characters and 18S rDNA sequence analysis. *Polskie Pismo Entomologiczne* 64: 2611–277
- Min GS, Kim SH, Kim W (1998) Molecular phylogeny of Arthropoda and their relatives: Polyphyletic origin of arthropodization. *Mol Cells* 8: 75–83
- Neefs JM, Peer YV, Rijk PD, Chapelle S, Wachter RD (1993) Compilation of small subunit ribosomal RNA sequences. *Nuc Acid Res* 21: 3025–2049
- Pashley DP, MacPherson BA, Zimmer EA (1993) Systematics of holometabolous insect orders based on 18S ribosomal RNA. *Mol Phyl Evol* 2: 132–142
- Peer YV, Caers A, Rijk PD, Wachter RD (1998) Database on the structure of small subunit ribosomal RNA. *Nuc Acid Res* 26: 179–182
- Rijk PD, Neefs JM, Peer YV, Wachter RD (1992) Compilation of small subunit ribosomal RNA sequences. *Nuc Acid Res* 20 supplement: 2075–2089
- Tautz D, Hancock JM, Webb DA, Tautz C, Dover GA (1988)

- Complete sequences of the rRNA genes of *Drosophila melanogaster*. Mol Biol Evol 5: 366–376
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL-X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. Nuc Acid Res 25: 4876–4882
- Volger AP, Pearson DL (1996) A molecular phylogeny of the tiger beetles (Cicindelidae): congruence of mitochondrial and nuclear rDNA data sets. Mol Phyl Evol 6: 321–338
- Volger AP, Welsh A, Hancock JM (1997) Phylogenetic analysis of slippage-like sequence variation in the V4 rRNA expansion segment in Tiger Beetles (Cicindelidae). Mol Biol Evol 14: 6–19
- Wheeler WC (1989) The systematics of insect ribosomal DNA. In “The hierarchy of life. Molecules and morphology in phylogenetic analysis” Ed by B. Fernholm, K. Bremer, H. Jornvall, Elsevier, Amsterdam, pp 307–321
- Whiting MF, Carpenter JC, Wheeler QD, Wheeler WC (1997) The Strepsiptera problem: phylogeny of the holometabolous insect orders inferred from 18S and 28S ribosomal DNA sequences and morphology. Syst Biol 46: 1–68

(Received March 11, 1999 / Accepted July 3, 1999)