

A New Cerithioidean Genus *Megistocerithium* (Gastropoda; Mollusca) from the Miocene of Southeast Asia: A Possible Relict of Mesozoic “Eustomatidae”

Authors: Kase, Tomoki, Kurihara, Yukito, Aguilar, Yolanda M., Pandita, Hita, Fernando, Allan Gil S., et al.

Source: Paleontological Research, 19(4) : 299-311

Published By: The Palaeontological Society of Japan

URL: <https://doi.org/10.2517/2015PR013>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A new cerithioidean genus *Megistocerithium* (Gastropoda; Mollusca) from the Miocene of Southeast Asia: a possible relict of Mesozoic “Eustomatidae”

TOMOKI KASE¹, YUKITO KURIHARA², YOLANDA M. AGUILAR³, HITA PANDITA⁴,
ALLAN GIL S. FERNANDO⁵ AND HIROKI HAYASHI⁶

¹Department of Geology and Paleontology, National Museum of Nature and Science, 4-1-1 Amakubo, Tsukuba, Ibaraki 305-0005, Japan
(e-mail: kase@kahaku.go.jp)

²Department of Geology, Faculty of Education, Mie University, 1557 Kurimamachiya-cho, Tsu, Mie 514-8507, Japan

³Marine Geological Survey, Mines and Geosciences Bureau, North Avenue, Diliman, Quezon City, 1101, Philippines

⁴Teknik Geologi, Sekolah Tinggi Teknologi Nasional Yogyakarta, Jl. Babarsari, Depok Sleman, Yogyakarta 55281, Indonesia

⁵National Institute of Geological Sciences, the University of Philippines, Diliman, Quezon City, 1101, Philippines

⁶Department of Geoscience, Interdisciplinary Graduate School of Science and Engineering, Shimane University, 1060 Nishikawatsu-cho, Matsue, Shimane 690-8504, Japan

Received October 25, 2014; Revised manuscript accepted March 17, 2015

Abstract. Based upon large cerithiform gastropod specimens from the middle Miocene beds of Java, Indonesia and the Philippines, a new genus and new species, *Megistocerithium magoi*, is described herein. Ten fragmentary specimens from the lower and upper Miocene beds of the Philippines are tentatively referred to this species. *M. magoi* Kase sp. nov. is amongst the largest fossil cerithioidean gastropods (except for Turritellidae). It was a dweller in intertidal sandy mudflats, probably in close proximity to mangrove forests. No species considered to be congeneric with this species have been found in the Cenozoic. On the other hand, *M. magoi* Kase gen. et sp. nov. shares apertural characters with the Middle Jurassic species *Eustoma tuberculosa*, the type species of a genus belonging to the yet to be well-recognized family Eustomatidae. *M. magoi* Kase gen. et sp. nov. is possibly a relict of the Mesozoic Eustomatidae. It is assumed to have been a grazer on mangrove litter like the potamidid genera *Terebralia* and *Telescopium* in modern mangrove swamps.

Key words: Cerithioidea, Gastropoda, Java, *Megistocerithium*, Miocene, Philippines

Introduction

Cerithioidean gastropods are a diverse group and a major component of younger Cenozoic shallow marine molluscan assemblages in the Indonesian and Philippine archipelagos. They are mostly represented by cerithiform (elongate, multiwhorled, and high-spired) shells with sculpture of spiral rows of granules and/or axial ribs. Nonetheless, unlike modern examples, supraspecific identification of fossil specimens from these areas is often difficult, because the complete outer lips and anterior siphonal canals of the last whorl, important shell characters for supraspecific systematics, are usually destroyed during fossilization or broken during excavation. The majority of fossil cerithioidean species descriptions from tropical Southeast Asia have been based on

shells without complete apertures (e.g. Martin, 1899; Shuto, 1978). Therefore, acquisition of specimens with a complete aperture is indispensable for correct identification as well as for understanding cerithioidean biodiversity in this bioprovince. Our collecting surveys on islands in the Philippines and in Java, Indonesia have resulted in discoveries of complete specimens for many cerithioid species with previously undetermined generic and/or familial positions.

This paper presents descriptions and a discussion of remarkably large cerithiform gastropod specimens obtained during our collecting surveys of the middle to upper Miocene formations in Java and the Philippines. The specimens are very infrequent in occurrence (represented by only 18 specimens) and represented mostly by pieces of spire whorls and/or incomplete last whorls.

However, one specimen from the upper Miocene Nyalindung Formation in West Java has the last whorl almost completely preserved. We are now able to realize that these fragmentary specimens belong to a new genus and species of superfamily Cerithioidea, among the largest Cerithioidea recorded. We suggest that this new genus is not congeneric to any known genera from Cenozoic formations or in modern seas, but can possibly be interpreted as a close relative of the Mesozoic genus *Eustoma* Piette, 1855.

Abbreviations used in this paper are as follows: EM = Geological Collections in the University of Lyon 1, France; NMNS = National Museum of Nature and Science, Tsukuba, Japan; SMNH = Department of Palaeo-zoology, Swedish Museum of Natural History, Stockholm, Sweden; UMUT = University Museum, the University of Tokyo, Tokyo, Japan.

Localities and stratigraphy

The specimens discussed in this study were obtained from two localities in West Java, Indonesia and four localities in the Philippines, ranging in age from middle to late Miocene (Figure 1). What follows are brief descriptions of the collecting sites, stratigraphy, lithology, sedimentary environments, and ages.

THL-1.—A creek bank in Tjitalahab valley in the Nyalindung area of Pelabuhanratu, West Java, Indonesia ($7^{\circ}01'16.3"S$, $106^{\circ}56'35.2"E$); Nyalindung Formation (Figure 1A). According to our survey, the Nyalindung Formation exposed in the Tjitalahab valley is about 220 m thick. The lower to middle part of the formation consists of dark sandy siltstone and calcareous sandstone, and contains diverse shallow marine mollusks such as conid, cerithiid and nassariid gastropods, arcid and tridaenid bivalves, and branching corals, suggesting a deposition in a tropical lower intertidal to upper subtidal environment. Oostingh (1938) designated the Nyalindung Formation as the stratotype of the Preangerian Stage, an Indonesian Neogene stage approximately equivalent to the middle Miocene (Bemmelen, 1972). Recently, Batenburg *et al.* (2011) suggested an age of ~10–13 Ma (late Serravallian to earliest Tortonian) for the Nyalindung Formation based on larger foraminifers from the nearby locality Tjiangsana. Planktic foraminifers are very few in this formation and all we found are not age-diagnostic. However, a nannofossil analysis by one of the authors has shown that the mollusk-bearing lower to middle part of the formation in the Tjitalahab valley contains the marker species *Discoaster hamatus*, indicating Martini's (1971) NN9 nannofossil zone (*ca.* 10 Ma; early late Miocene) (A. G. S. Fernando, unpublished data).

CAG-3.—A small exposure in a small creek, Tjikantjaora, in Tjiangsana village in the Nyalindung area ($7^{\circ}01'53.3"S$, $106^{\circ}55'47.4"E$), about 2 km southwest of locality THL-1; Nyalindung Formation (Figure 1A). Calcareous with larger foraminifers, interbedded with dark silty sandstone. The age is probably the same as THL-1.

C-DNO-3 (1).—An abandoned small-scale coal-mining operation, Sitio Capilyahan, Masaba, Danao City, Cebu, Philippines ($10^{\circ}33'18.3"N$, $123^{\circ}57'48.3"E$) (Figure 1B). This site appears to be the same as F272 of Smith (1913; Kanno *et al.*, 1980), where Smith (1913, pl. 6, fig. 5) illustrated a fragmentary specimen of *Megistocerithium magoi* Kase gen. et sp. nov. Gray, bituminous sandy siltstone spoil from the mine contains the potamidid *Vicarya verneuili* and small mollusks such as conids, cypraeids, and turrids. Kanno *et al.* (1980) mentioned the occurrence of *Telescopium* and fragments of brackish-water shells from this site, suggesting deposition in an intertidal flat beside a mangrove forest. The site was mapped as the lower Miocene Malubog Formation (Mines and Geosciences Bureau, 1983), and A. G. S. Fernando (unpublished) identified *Sphenolithus belemnos* among others, indicating that the sediments from this site are within the NN3 nannofossil zone, early Miocene in age.

CDN-1.—A creek bed in Palaka Creek, Sitio Palaka, Barangay Bactalon, Sipalay City, Negros Occidental, Philippines ($9^{\circ}48'43.4"N$, $122^{\circ}34'49.6"E$) (Figure 1B). Basal part of an unnamed formation consisting of dark-gray silty sandstone interbedded with a thick coal bed in the lower part. Mollusks are prolific, beautifully preserved, and appear to be autochthonous in origin without any indication of strong currents during the deposition. Dominant mollusks are three potamidid species (*Menkrawia callosalabiata*, *Tymanonotus beberkirianus*, and *Vicarya verneuili*), and small cerithiids, *Thais* sp., *Volema* spp., small conids, and oysters. Additionally, *Telescopium titan*, *Rhizophorimurex capucinus* and *Terebralia* sp., whose modern species or congeners are dwellers in mangrove swamps, were also found. The sedimentary facies and fossil mollusks suggest that the mollusk-bearing beds in this locality were deposited on an intertidal flat within or close to a mangrove forest.

H. Hayashi (unpublished) identified planktic foraminifers from the topmost bed of this unnamed formation including age-diagnostic species such as *Menardella praemenardii* and *Paragloborotalia siakensis*, indicating a middle Miocene age spanning 14.50 Ma to 11.50 Ma according to Tian *et al.* (2008). Additionally, A. Gil S. Fernando (unpublished) identified the nannofossil *Sphenolithus heteromorphus*, among others, from a sandy siltstone bed slightly below the topmost bed. This species ranges from Martini's (1971) NN4 to NN5, 17.90 Ma to 13.83 Ma according to Tian *et al.* (2008). In summary,




Figure 1. Collecting sites of *Megistocerithium magoi* Kase gen. et sp. nov. in Java, Indonesia (A) and Visayan islands of Philippines (B).

the upper part of the unnamed formation ranges in age from 14.38 Ma to 13.83 Ma, late Langhian.

DNO-1.—Bank of the Camansihan Creek, Barangay Poblacion, Danao municipality, Bohol, Philippines

(9°56'27.2"N, 124°13'03.0"E) (Figure 1B). An unnamed formation consisting of dark-gray silty sandstone with mollusks and stony corals (family Poritidae), suggesting an intertidal to upper subtidal environment. Fernando *et*

al. (2008) has shown that the formation can be assigned to Martini's (1971) NN11 zone (*ca.* 8 Ma, late Miocene) based on nannofossils.

PSS-1.—A small sand quarry in Barangay Bacuranan, Passi City, Panay, Philippines ($11^{\circ}08'35.8''N$, $122^{\circ}39'03.7''E$) (Figure 1B). Dingle Formation. Calcareous silty sandstone about 10 m thick, with a 40 cm-thick lignite bed at the base. The sandstone contains cerithiid, conid and large strombid gastropods and fragments of corals, all allochthonous in occurrence, suggesting deposition in a shallow marine environment. Mines and Geosciences Bureau (2010) assigned the Dingle Formation to the upper Miocene, and H. Hayashi (unpublished) identified *Globorotalia plesiotumida* and *Globoturborotalia nepenthes*, among others, from this locality, also suggesting a late Miocene age between 8.58 Ma and 4.43 Ma based on Tian *et al.* (2008) and Hilgen *et al.* (2012).

Systematic description

(Tomoki Kase)

?Family Eustomatidae Cossmann, 1906
Genus ***Megistocerithium*** Kase gen. nov.

Type species.—*Megistocerithium magoi* Kase sp. nov. from the Miocene of Java, Indonesia and the Philippines.

Diagnosis.—Shell very large, elongate cerithiform; later whorls seven-sided pyramidal in shape formed by juxtaposed, weakly opisthocline, prominent axial ribs. Spiral sculpture of sharp primary cords, their interspaces densely packed with one to several secondary cords, all crossed by axial ribs. Aperture elliptical, flared, plane of aperture almost orthocline to shell axis. Last whorl constricted anteriorly, with an obtuse ventrolateral varix. Outer lip thick, flaring, varix-like, bevelled, and smooth within, crenate outside in accordance with strong spiral cords on last whorl; adapical end of outer lip well extended onto last whorl, forming flat triangular area, with a shallow groove due to deposition of shell material defined by a parietal plait extending from posterior anal canal. Anterior siphonal canal long, tube-like, extending antero-dorsally. Parietal area smooth with thick callus. Columellar lip covered with thick, expanded, reflected inductura, being broadly concave dorsally.

Etymology.—The genus name comes from the combination of the Greek *megistos*, meaning greatest, and *cerithium*, in reference to it having the largest shell size among cerithioideans.

Remarks.—The genus is monotypic at present. For comparison and relationship with other cerithioidean genera, see “Discussion” below.

***Megistocerithium magoi* Kase sp. nov.**

Figures 2, 3

Cerithium (Campanile) sp. Smith, 1913, p. 269, pl. 4, fig. 5.
?Pyrazus sp. 3. Dharma, 2005, p. 308, pl. 119, fig. 3a, b.

Material.—Holotype, NMNS PM26959, *ca.* 190 mm long (reconstructed), loc. CAG-1, Nyalindung Formation, late Miocene, West Java; five paratypes, NMNS PM26960–26964, fragmentary specimens, loc. THL-1, Nyalindung Formation, late Miocene, West Java; two specimens, NMNS PM26965, 26966, loc. CDN-1, middle Miocene, Philippines. Ten specimens are tentatively referred to this species: one fragmentary specimen, loc. C-DNO-3(1), NMNS PM26967, Malubog Formation, early Miocene, Philippines; eight specimens, BMNS PM26968–26975, all fragmentary, loc. PSS-1, late Miocene, Philippines; one specimen, NMNS PM26976, incomplete specimen without most of the last whorl, loc. DNO-1, late Miocene, Philippines.

Etymology.—Named after Wency Mago of Mining and Geosciences Bureau, Quezon City, Philippines for his assistance during our more than seven years of field work.

Diagnosis.—Same as for the genus.

Description.—Holotype (Figure 2C–E). Shell of five whorls preserved, very large (probably reaching 190 mm in height if complete), high-spired, cerithiform, pleural angle about 25° . Protoconch and early teleoconch whorls missing. Suture shallow, somewhat undulated. Spire whorls weakly convex in profile. Last three spire whorls with seven prominent axial ribs and much wider interspaces; axial ribs strong, round-topped, weakly opisthocline (extent of inclination slightly variable), becoming spiny at mid-whorl toward last whorl. Spire whorls covered with dense, slightly wavy spiral sculpture, consisting of spiral cords or threads of variable prominence; those just adapical and abapical to suture and at mid-whorl more prominent than others. Last whorl with one abapically thickened ventrolateral varix located about 270° volution back from outer lip; axial ribs on last whorl spiny, becoming gradually shorter and less prominent toward outer lip; spiral cords becoming thicker, two on base and one on mid-whorl prominent. Aperture elliptical; anterior siphon long, tube-like, extending antero-dorsally, narrowly open. Parietal lip broadly concave, covered with smooth, thin callus, fading gradually into last whorl, with small parietal plate. Columella covered with thick, reflected, rimmed inductura, dorsally concave as a whole. Outer lip flared and varix-like, rugged externally, smooth and beveled within due to thick callus; adapical part of outer lip forming triangular wedge covered by thick callus, flattened dorsally, with shallow




Figure 2. *Megistocerithium magoi* Kase gen. et sp. nov. **A**, NMNS PM26965, shell height 78.4 mm, middle Miocene unnamed formation, Loc. CDN-1, Negros, Philippines; **B**, NMNS PM26960, paratype, shell height 46.5 mm, Nyalindung Formation, THL-1, West Java, Indonesia; **C-E**, NMNS PM26959, holotype, shell height 133.0 mm, upper Miocene Nyalindung Formation, CAG-3, West Java, Indonesia; **C**, lateral view; **D**, apertural view; **E**, adapertural view.




Figure 3. *Megistocerithium magoi* Kase gen. et sp. nov. **A**, NMNS PM26968, shell height 65.3 mm, upper Miocene Dingle Formation, Loc. PSS-1, Panay, Philippines; **B**, **C**, NMNS PM26969, shell height 58.5 mm, upper Miocene Dingle Formation, Loc. PSS-1, Panay, Philippines; **B**, adapertural view; **C**, apertural view; **D**, **E**, NMNS PM26967, shell height 67.6 mm, lower Miocene Malubog Formation, Loc. C-DNO-3(1), Cebu, Philippines; **D**, adapertural view; **E**, apertural view; **F**, **G**, NMNS PM26976, apertural and adapertural views, shell height 95.4 mm, unnamed upper Miocene formation, loc. DNO-1, Bohol, Philippines; **F**, apertural view; **G**, adapertural view.

groove extending from adapical notch of aperture.

Dimensions.—NMNS PM26959, holotype, height 133 mm, width 61.6 mm; NMNS PM26960, paratype,

height 46.5 mm, width 27.1 mm; NMNS PM26965, paratype, height 78.4 mm, width 36.3 mm; NMNS PM26969, height 58.5 mm, width 49.6 mm; NMNS PM26976,

height 95.4 mm, width 47.8 mm.

Remarks.—The occurrence of this species is very infrequent both in Java and in the Philippines. Other than the holotype and the five paratypes, Smith (1913, pl. 6, fig. 5) illustrated a fragmentary specimen identified as *Cerithium (Campanile)* sp. from Cebu, Philippines and Dharma (2005, pl. 119, fig. 3a, b) illustrated two specimens of this species identified as ?*Pyrazus* sp. from the Nyalindung Formation in Tjitalahab, West Java. We here refer eleven fragmentary specimens from three localities in the Philippines to this species, ranging in age from the early to late Miocene. Two specimens from the unnamed middle Miocene formation at locality CDN-1, Negros have six and four preserved whorls, respectively, and are sculptured with densely packed fine spiral cords, as on the holotype and paratypes from Java (Figure 2A). In these specimens, the axial ribs are more angulated at the mid-whorl compared to the early whorls on the holotype. However, this difference is too minor to separate the specimens from CDN-1 as a different species.

We tentatively refer the ten fragmentary specimens from the three localities in the Philippines to *Megistocerithium magoi* Kase gen. et sp. nov. (Figure 3A–G). NMNS PM30009 from the Malubog Formation at locality C-DNO-3 (1) is a silicified specimen with only three preserved whorls including the posteriorly expanded outer lip of the last whorl (Figure 3D, E). The spiral sculpture of this specimen is composed of thick cords, unlike the densely packed fine spirals in addition to some thicker cords in the holotype. However, in silicified specimens found in coal-mining sites in the Philippines, the surface shell layer of many gastropods is unsilicified and easily eroded away, particularly in float specimens. We suggest that NMNS PM26967 is a case of such selective silicification and subsequent erosion.

Seven specimens from the upper Miocene Dingle Formation at locality PSS-1 are all fragmentary and their shell surfaces are strongly eroded. Two of the specimens have a preserved last whorl and have a posteriorly expanded outer lip as in the holotype (Figure 3B, C). The densely packed fine spiral cords still remain but are obscure in three specimens, and the axial ribs are pyramidal in early spire whorls with seven ribs per whorl (Figure 3A). In these specimens the axial ribs are laterally flattened unlike those in the holotype.

One specimen, NMNS PM26976, from the unnamed upper Miocene formation at locality DNO-1 in Bohol, has seven preserved whorls including the last whorl and a posteriorly expanded outer lip (Figure 3F, G). The shell surface is sculptured with densely packed fine spiral cords and strong axial ribs. However, the shell is more slender, the crests of the axial ribs are more rounded and not angulated at the middle of the whorls, and the spiral

band below the suture is thicker than those of the holotype. Taking a wide variability of shell sculpture in cerithioidean gastropods into consideration (e.g. Houbrick, 1992), it seems likely that the differences are within the morphological variation of *Megistocerithium magoi* Kase gen. et sp. nov.

To the best of our knowledge, no species definitely congeneric to *Megistocerithium magoi* Kase gen. et sp. nov. is known from the Cenozoic and Mesozoic formations or modern seas. *Pyrazus protebeninus* Vredenburg, 1928 from the lower Miocene of Gaj in Sindh, Pakistan, resembles *M. magoi* Kase sp. nov. Harzhauser (2007) reassigned *P. protebeninus* to the genus *Pyrazisinus* Heilprin, 1887, and also suggested that it was closely related to *Pyrazisinus monstrosus* (Grateloup, 1847) (Lesport and Cahuzac, 2002). Two specimens illustrated by Vredenburg (1928, pl. 18, figs. 6, 7) may attain well over 100 mm in shell length when complete, and have eight prominent opisthocline axial ribs that are raised into spines where they are crossed by rather strong spiral cords.. However, assignment of this Pakistan species either to *Pyrazisinus* or to *Megistocerithium* Kase gen. nov. is difficult because the shells of the Pakistan species are too poorly preserved and their apertures are completely missing.

Discussion

Generic comparison to Cenozoic cerithiform gastropods.—*Megistocerithium magoi* Kase gen. et sp. nov. is remarkably large. Cerithiform gastropods (gastropods of widely separated taxonomic groups similar in shell form to the genus *Cerithium*; Sälgeback and Savazzi, 2006) that approximate or exceed the shell height of this genus are known sporadically from the Cenozoic. A well known example is *Campanile giganteum* (Lamarck, 1804) from the Eocene of France, which attains over a meter in length, while during the Neogene and Quaternary, the largest species of *Campanile*, *C. symbolicum* Iredale, 1917 from the Pleistocene to Recent of southwestern Australia, is 244 mm long (Houbrick, 1981, 1984). *Campanile* is currently classified as Campanilidae of the superfamily Campaniloidea based on its anatomical characters (e.g. Haszprunar, 1988; Healy and Wells, 1998). *Campanile* is similar in shell characters to species of the superfamily Cerithioidea but differs in having an unflared opisthocline outer lip that sweeps backwards in a broad sinus abapical to the suture, and a calcified periostracum (usually not preserved in fossil species). *Megistocerithium* Kase gen. nov. is not related to *Campanile* because the outer lip is expanded and almost orthocline, without the backward-directed sinus abapical to the suture.

The younger Cenozoic shallow-marine carbonate sedimentary basins of southern Australia are known for the occurrence of large cerithiform gastropods (Ludbrook, 1971). Aside from two campanilid species, *Thericium (Chavanicerithium) torri* (Tate, 1899), *Thericium (Chavanicerithium) pritchardi* (Harris, 1897) and *Jetwoodsia apheles* (Tenison-Woods, 1879) reach over 100 mm in length. These species have a short, twisted anterior siphon and an unflared, simple outer lip, indicating that they are not related to *Megistocerithium* Kase gen. nov. In modern seas, the largest cerithiid species is *Pseudovergatus* (s. s.) *nobilis* (Reeve in Sowerby, 1855) in the tropical Indo-West Pacific, which reaches up to 167 mm in length (Houbrick, 1978). This species is not congeneric with *Megistocerithium* Kase gen. nov. as it has a smooth shell, a simple, unflared outer lip and a short twisted anterior canal. *Cerithium nodulosum* Bruguière, 1792 (type species of the genus) in the tropical Indo-West Pacific reaches 114 mm in length (Houbrick, 1992), has a slightly flared outer lip and thick axial ribs that are nodulous on later whorls as in *Megistocerithium magoi* Kase gen. et sp. nov., but this similarity is only superficial. The aperture has a distinct spout at the anal canal, the outer lip has deep spiral grooves interiorly and a claw-like anterior extension that crosses over the short anterior siphonal canal slightly reflexed to the left in *C. nodulosum*. In contrast, the anal canal is indistinct and the outer lip is smooth within and merges into the anteriorly extended siphonal canal in *M. magoi* Kase gen. et sp. nov. *Megistocerithium* Kase gen. nov. has a remarkable shell character unknown in Cerithiidae: a triangularly projected posterior margin of the outer lip lined internally with thick callus deposits and a shallow furrow within, indicating that this new genus may not belong to Cerithiidae.

Cerithiform gastropods with an expanded outer lip such as *Megistocerithium* Kase gen. nov. are also known in Potamididae and Batillariidae. The best example is the Miocene potamidid genus *Vicarya* d'Archiac and Haime, 1854, thought to have been an inhabitant of mangrove swamps in Southeast Asia and Japan (e.g. Oyama, 1950; Chinzei, 1978; Tojo and Masuda, 1999). Like *Megistocerithium* Kase gen. nov., the outer lip of *Vicarya verneuili* (d'Archiac, 1851) and *V. yokoyamai* Takeyama, 1933 is well expanded posteriorly to form a triangular projection with a thick callus lining internally and a shallow furrow within (Kanno, 1986; Sälgeback and Savazzi, 2006, fig. 13D). *Vicarya* clearly differs from *Megistocerithium* Kase gen. nov. in having a very short siphonal canal, a claw-like basal lip, a remarkable knob-like swelling of callus on the shell base and a different type of shell sculpture that consists of a subsutural spiral row of spiny tubercles and some beaded spiral cords. *Pyrazus*

ebeninus (Bruguière, 1792), the largest modern species of Batillariidae, living in the tidal flats and estuaries of Australian and Tasmanian waters, has strong axial and spiral sculpture similar to that of *M. magoi* Kase sp. nov. *P. ebeninus* also possesses a posteriorly projected outer lip but the outer lip is less extended than in *M. magoi* Kase gen. et sp. nov., and does not have a shallow groove within the interior surface. *P. ebeninus* clearly differs from *M. magoi* Kase sp. nov. in having a very short anterior siphon and a wide backward-directed sinus in the upper part of the outer lip.

Comparison to Mesozoic cerithiform gastropods.—As discussed above, no species considered to be congeneric with *Megistocerithium magoi* Kase gen. et sp. nov. have been found in the Cenozoic in spite of the rich fossil records of tropical faunas of this age. On the other hand, *Megistocerithium* Kase gen. nov. has apertural characters quite similar to *Eustoma* Piette, 1855, of which the type species is *E. tuberculosa* Piette, 1855 from the Bathonian (Jurassic) of France (Figure 4A). Cossmann (1906) proposed the new family Eustomatidae (as Eustomidae) with *Eustoma* as the type genus, and renamed it *Diatinostoma* because *Eustoma* and *Eustomum* Leidy, 1851 (a flatworm) appeared to be in homonymy. However, the latter name, according to Cossmann (1906), was a misspelling. Wenz (1940) regarded *Eustomum* as a senior homonym of *Eustoma*, and Williams and Richards (1968, p. 217) took the same view. However, *Eustoma* is not a homonym of *Eustomum* based on the current code (ICZN, 4th ed., 1999, Art. 32.2 and 32.5); *Diatinostoma* is, therefore, an unnecessary name.

Piette (1855) described *Eustoma tuberculosa*, accompanied with line drawings of two specimens (Figure 4B), and Piette (1856) described it again with more elaborate line drawings (Figure 4C, D). Six syntypes of *E. tuberculosa*, all incomplete specimens, are stored in the Geological Collections of the University of Lyon 1, France (Figure 4A). The shell may attain over 60 mm in height, is elongate cerithiform and has a long, anteriorly extended, tube-like siphonal canal. The outer lip is well expanded, its posterior tip reaches the penultimate whorl, and the posterior portion of the outer lip forms a triangular projection. This projection is lined internally with a thick callus deposit, with a shallow groove extending from the posterior corner of the aperture to the posterior tip of the outer lip. The parietal area is covered with a thick inductura that spreads over the basal area of the last whorl. The exterior sculpture is comprised of round-topped axial ribs on the adapical half of each whorl in addition to fine spiral cords. Although the shell size is smaller and the exterior sculpture is different, the other shell characteristics of *E. tuberculosa* fit well with those of *Megistocerithium magoi* Kase gen. et sp. nov., sug-




Figure 4. A, Six syntypes of *Eustoma tuberculosa* Piette, 1855, EM 31352, in the Geological Collections of the University of Lyon 1, France (courtesy of E. Robert); cardboard mount of syntypes 120 mm high and 70 mm wide, Bathonian of France; B, hand drawing of *E. tuberculosa* Piette, 1855, seemingly based on two of the syntypes (upper left and lower left in A) (after Piette, 1855, pl. 31, fig. 2); C, D, hand drawings of *E. tuberculosa* Piette (after Piette, 1856, pl. 2, fig. 10, 11); E–G, *Pyrazopsis* sp., previously identified by Sälgeback and Savazzi (2006) as ?*Diatinostoma* sp., SMNH Mo 113872, apertural, lateral and adapertural views, shell height 43.5 mm, Upper Cretaceous of Syria; H, I, *Pyrazus?* *scalariformis* Nagao, 1934, Aptian Miyako Group, Japan; H, UMUT MM15695, shell height 69.5 mm, apertural view; I, UMUT MM 15694 shell height 58.8 mm, apertural view.

gesting that the two species might be related to each other.

Cossmann (1906, p. 12–13) listed ten species referable to *Eustoma* (as *Diatinostoma*) ranging in age from Bathonian to Turonian (Cretaceous). However, complete apertures are unknown in all these species, so that the assignment of these species to *Eustoma* remains unconfirmed. Hikuroa and Kaim (2007) consider the genus *Silberlingiella* Frýda and Blodgett, 2003 (originally placed in the family Purpurinidae of Littorinoidea) to be a member of Eustomatidae. *Silberlingiella ornata* Frýda and Blodgett, 2003 (the type species of the genus) from the Norian (Triassic) of Nevada, USA and *Silberlingiella latadyensis* Hikuroa and Kaim, 2007 from the Middle/Upper Jurassic of Antarctica are both characterized by a small, elongate, turriculate shell with a row of nodes or spines at the narrow ramp angle slightly abapical to the

upper suture, a concave and vertical whorl surface and a keeled basal periphery. According to Hikuroa and Kaim (2007), the outer lip is expanded and sinuous, and the anterior margin is notched to form an incipient siphonal canal in *S. latadyensis*. These shell characteristics in *Silberlingiella* are quite different from *Megistocerithium magoi* Kase gen. et sp. nov.

Diatinostoma? eximium Sayn, 1932 from the Urgonian (Lower Cretaceous) of Spain is likely to be related to *Eustoma tuberculosa* and *Megistocerithium magoi* Kase gen. et sp. nov. In addition to the pyramidal spire whorls, the outer lip of this species is expanded onto the penultimate whorl, covered with a thick callus internally, and has a gutter-like groove extending from the posterior corner of the aperture to the posterior tip of the outer lip. Two more species worth mentioning are *Pyrazus?* *scalariformis* Nagao, 1934 (Kase, 1984) from the Aptian

(Cretaceous) of Japan (Figure 4H, I) and *Pyrazus stuieri* Cossmann (1906) from the Coniacian (Cretaceous) of France. These two species have a pyramidal upper spire and an internally callused posterior projection of the outer lip with an indistinct shallow groove within like that of *Megistocerithium* Kase gen. nov. However, the complete anterior siphonal canal is not known in these three species.

Sälgeback and Savazzi (2006, fig. 16A) illustrated an almost complete cerithiform shell from the Upper Cretaceous of Syria and identified it as *?Diatinostoma* sp. This specimen has five-sided pyramidal upper spire whorls, an obliquely elongate aperture rimmed with a callus deposit along the parietal and columellar area, an expanded outer lip, an obscurely defined notch at the posterior end of the aperture, and a moderately long, variously twisted, open siphonal canal (Figure 4E–G). The posterior part of the outer lip appears to form a small triangular projection that is not covered by a callus deposit. This Syrian specimen has many shell characters much in common with the Eocene species *Pyrazopsis pentagonatus* (Schlotheim, 1820) and *Pyrazopsis angulatus* (Solander in Brander, 1766) except that the anterior canal is longer, suggesting that it belongs to the Batillariidae. As already pointed out by Saul and Squires (2003), the shell apertures of *Pyrazopsis quinquecostatus* (Egojan, 1955) from the Upper Cretaceous of Armenia, the type species of *Pyrazopsis* Akopjan, 1972, and of seven other Cretaceous species Akopjan (1972, 1976) referred to *Pyrazopsis* are all unknown. Therefore, it is not known whether *P. pentagonatus* and *P. angulatus* belong to *Pyrazopsis*.

Saul and Squires (2003) described a remarkable new genus and species, *Alamirifica corona*, of unknown systematic position, based on a complete specimen from the Turonian of California. Interestingly, this specimen has an elongate cerithiform shell and pyramidal upper spire whorls like the above-mentioned Cretaceous species, which Akopjan (1972, 1976) referred to *Pyrazopsis*, but surprisingly its aperture is circular and rimmed without a posterior notch and has a moderately long, closed anterior siphonal canal. Their finding suggests that, without the complete aperture, the systematic position of cerithiform gastropods with pyramidal spire whorls from the Mesozoic cannot be determined definitely.

Cossmann (1906) pointed out that species of Eustomatidae have shell characteristics (expanded outer lip and straight, tube-like anterior canal) intermediate between those of Stromboidea (Alatacea of Cossmann) and Cerithioidea. However, recent anatomical and molecular phylogenetic analyses do not support such a close relationship between Cerithioidea and Stromboidea (e.g. Colgan *et al.*, 2007). Wenz (1940) placed Eustomatidae in the Cerithioidea with a query, and Sälgeback and

Savazzi (2006) placed *Eustoma* (as *Diatinostoma*) in the Procerithiidae. More recently, Hikuroa and Kaim (2007) suggested that Eustomatidae was an ancestor of Batillariidae or Potamididae. Eustomatidae is a poorly known family based only on a few morphological characters. Therefore, the status of Eustomatidae as a family-group taxon and the reciprocal relationships with other cerithioidean families require additional information on included species and genera and morphological characters such as larval shells. We agree with Hikuroa and Kaim (2007) that Eustomatidae is either a family close to Batillariidae or Potamididae, or assume that it is a clade within Potamididae because of the morphological similarity to some members of this family.

In conclusion, *Megistocerithium magoi* Kase gen. et sp. nov. is not related to any cerithioidean species from the Cenozoic, and is possibly related to or is a relict of the poorly known Mesozoic family Eustomatidae.

Paleoecology.—Most modern cerithioideans are herbivores and detritus feeders (Healy and Wells, 1998), and it may have been the same for *Megistocerithium magoi* Kase gen. et sp. nov. The possible affinity to Batillariidae or Potamididae as well as the sedimentary facies and associated autochthonous mollusks of the host sediments suggest that *M. magoi* Kase gen. et sp. nov. was an intertidal sandy mudflat dweller within or close to mangrove forests. The long, antero-dorsally extended siphonal canal further suggests a semi-infaunal mode of life, but this may have been facultative because the unflattened dorsal region of the last whorl and the prominent axial sculpture appear unsuitable for continuous burrowing in sediment (Sälgeback and Savazzi, 2006). The shallow groove in the triangular projection of the outer lip may have functioned as an exhalent sinus during an inactive semi-burrowing phase.

Megistocerithium magoi Kase gen. et sp. nov. is very large and is suggested to have been a dweller within or close to mangrove swamps. Today, the potamid species *Terebralia palustris* and *Telescopium telescopium* and the corbiculid bivalve *Geloina* are core species in the Indo-West Pacific mangrove molluscan communities (Plaziat, 1984, 1995; Plaziat *et al.*, 2001), and are extraordinarily large compared with other potamid species and bivalves in the community. Analogous assemblages with large potamids (e.g. *Telescopium titan* Martin, 1890a and *Terebralia wickhami* Martin, 1890b) and *Geloina* (e.g. *G. stachi* Oyama, 1950) are known from deposits of the latest Oligocene onwards in the Philippines (Iwasaki, 1970; Kanno, 1986; unpublished data) and from the middle Miocene of Japan (Oyama, 1950; Chinzei, 1978; Itoigawa, 1978; Ozawa *et al.*, 1986; Tsuda *et al.*, 1986; Inoue, 2007; etc.), whereas *Terebralia* and other potamid gastropods during the Paleogene are not as large

as those from the Miocene onwards (e.g., Kowalke, 2001, 2003; *Terebralia subcorvina* is an exception). Interestingly, ecological observations and isotope and fatty acid marker analyses have demonstrated that in modern mangrove swamps, these large mollusks exploit food resources mostly from mangrove litter such as leaves and fruits or detritus derived from mangrove trees (Alexander *et al.*, 1979; Nishihara, 1983; Rodelli *et al.*, 1984; Houbrick, 1991; Slim *et al.*, 1997; Bachok *et al.*, 2003; Bouillon *et al.*, 2004; Fratini *et al.*, 2004). Vermeij (2011, 2012) discussed how the body size of the largest species in shallow marine molluscan guilds (group of species with a similar habit and trophic role) reflects the relative amount of available food supply, and that a change in the maximum size within a guild over time chronicles a change in productivity. We here suggest that the appearance of large mangrove mollusks from the latest Oligocene onwards indicates an increase in mangrove litter in the tropical West Pacific. We assume that *M. magoi* Kase sp. nov. was another gastropod exploiting mangrove litter as a major food resource during the Miocene.

Acknowledgements

We would like to thank L. L. Jasareno (Director, Mines and Geosciences Bureau, Philippines) for a research permit, and W. Mago and E. Azurin (Mines and Geosciences Bureau, Philippines) for their assistance during field-work. We also thank C. Skovsted (Swedish Museum of Natural History), T. Sasaki (University Museum, the University of Tokyo), and Y. Kobayashi (Hokkaido University Museum, Hokkaido University) for the loan of the specimens under their care, E. Robert (University of Lyon 1, France), who provided high-resolution images of the syntypes of *Eustoma tuberculosus*, and R. Nakashima, who provided us with one of the references. Thanks are also due to A. Kaim (Polish Academy of Sciences, Poland), an anonymous reviewer for their helpful comments on the manuscript and A. G. Beu (Institute of Geological and Nuclear Sciences, New Zealand) for his improvement of the English text. This study was funded by the National Museum of Nature and Science, Tokyo and a Grant-In-Aid for Scientific Research from the Japan Society for the Promotion of Science (JSPS) (no. 18253007).

References

- Akopjan, V. T., 1972: On a new cerithioidean family. *Izvestiya Akademii Nauk Armyanskoy SSR. Nauki o Zemlye*, vol. 25, p. 3–14. (in Russian and Armenian; original title translated)
- Akopjan, V. T., 1976: *Late Cretaceous Gastropods from the Armenian SSR*, 443 p. Izdatel'stvo Akademii Nauk Armyanskoy SSR, Yerevan. (in Russian; original title translated)
- Alexander, C. G., Cutler, R. L. and Yellowless, D., 1979: Studies on the composition and enzyme content of the crystalline style of *Telescopium telescopium* (L.) (Gastropoda). *Comparative Biochemistry and Physiology*, vol. 64B, p. 83–89.
- Archiac, A. d', 1851: *Histoire des Progrès de la Géologie de 1834 à 1849*, Tome 3, 624 p. Au Lieu des Séances de la Société, Paris.
- Archiac, A. d' and Haime, J., 1853–1854: *Description des Animaux Fossiles du Groupe Nummulitique de l'Inde*, 373 p., 36 pl. Gide et J. Baudry, Paris.
- Bachok, Z., Mfilinge, P. L. and Tsuchiya, M., 2003: The diet of the mud clam *Geloina coaxans* (Mollusca, Bivalvia) as indicated by fatty acid markers in a subtropical mangrove forest of Okinawa, Japan. *Journal of Experimental Marine Biology and Ecology*, vol. 292, p. 187–197.
- Batenburg, S. J., Reichart, G.-J., Jilbert, T., Janse, M., Wesselingh, F. P. and Renema, W., 2011: Interannual climate variability in the Miocene: High resolution trace element and stable isotope ratios in giant clams. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 306, p. 75–81.
- Bemmelen, R. W. van, 1972: *The Geology of Indonesia. Volume 1A. General Geology of Indonesia and Adjacent Archipelagos, Second Edition*, 732 p. Martinus Nijhoff, The Hague.
- Bouillon, S., Moens, T., Overmeer, I., Koedam, N. and Dehairs, F., 2004: Resource utilization patterns of epifauna from mangrove forests with contrasting inputs of local versus imported organic matter. *Marine Ecology Progress Series*, vol. 278, p. 77–88.
- Brander, G., 1766: *Fossilia Hantoniensis Collecta, et in Musaeo Britannico Deposita, a Gustavo Brander*, 43 p, 9 pl. London. [Catalogue written by Daniel Solander.]
- Bruguière, J. G., 1789–1792: *Encyclopédie Méthodique, ou par Ordre de Matières, par une Société de Gens de Lettres, de Savans et d'Artistes: Précédée d'un Vocabulaire Universel, Servant de Table pour Tout l'Ouvrage, Ornée des Portraits de MM. Diderot & d'Alembert, Premiers Éditeurs de l'Encyclopédie. Histoire Naturelle des vers. Tome Premier*, [iii] + xviii + 758 p. Panckoucke, Paris.
- Chinzei, K., 1978: Neogene molluscan faunas in the Japanese islands: an ecologic and zoogeographic synthesis. *Veliger*, vol. 21, p. 155–170.
- Colgan, D. J., Ponder, W. F., Beacham, E. and Macaranas, J., 2007: Molecular phylogenetics of Caenogastropoda (Gastropoda: Mollusca). *Molecular Phylogenetics and Evolution*, vol. 42, p. 717–737.
- Cossmann, M., 1906: *Essais de Paléoconchologie Comparée. Tome 7*, 261 p., 14 pl. M. Cossmann and F. R. de Rudeval, Paris.
- Dharma, B., 2005: *Recent and Fossil Indonesian Shells*, 423 p. ConchBooks, Hackenheim.
- Egojan, V. L., 1955: *Upper Cretaceous Deposits of the Southwestern Part of the Armenian SSR*, 270 p. Izdatel'stvo Akademii Nauk Armyanskoy SSR, Yerevan. (in Russian; original title translated)
- Fernando, A. Gil S., Fernandez, A. R. C., Aguilar, Y. M., Kurihara, Y. and Kase, T., 2008: Late Miocene calcareous nannofossils from Danao basin, Bohol (Visayan Basin), Philippines. *Bulletin of the National Museum of Nature and Science, Series C*, vol. 34, p. 27–38.
- Fratini, S., Vigiani, V., Vannini, M. and Cannicci, S., 2004: *Terebralia palustris* (Gastropoda; Potamididae) in a Kenyan mangal: size structure, distribution and impact on the consumption of leaf litter. *Marine Biology*, vol. 144, p. 1173–1182.
- Frýda, J. and Blodgett, R. B., 2003: *Silberlingiella*, a new purpurinid genus (Littorinoidea, Gastropoda) from the Late Middle Norian

- (Late Triassic) of the Clan Alpine Range, western Canada. *Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg*, vol. 87, p. 47–54.
- Grateloup, J.-P. S. de, 1847: *Conchyliologie Fossile des Terrains Tertiaires du Bassin de l'Adour (Environs de Dax). Tome 1. Univalves*. Atlas 48 pl, index 12 p. T. Lafargue, Bordeaux.
- Harris, G. F., 1897: *Catalogue of the Tertiary Mollusca in the Department of Geology, British Museum (Natural History). Part 1. The Australasian Tertiary Mollusca*, 407 p. British Museum (Natural History), London.
- Harzhauser, M., 2007: Oligocene and Aquitanian gastropod fauna from the Sultanate of Oman and their biogeographic implications for the early western Indo-Pacific. *Palaeontographica, Abteilung A*, vol. 280, p. 75–121.
- Haszprunar, G., 1988: On the origin and evolution of major gastropod groups, with special reference to the Streptoneura (Mollusca). *Journal of Molluscan Studies*, vol. 54, p. 367–441.
- Healy, J. M. and Wells, F. E., 1998: Superfamily Cerithioidea. In, Beesley, P. L., Ross, G. J. B. and Wells, F. E. eds., *Mollusca: The Southern Synthesis. Part B. Fauna of Australia Volume 5*, p. 707–733. CISRO Publishing, Melbourne.
- Heilprin, A., 1887: Explorations on the west coast of Florida and in the Okeechobee wilderness. With special reference to the geology and zoology of the Floridian Peninsula. *Transactions of the Wagner Free Institute of Science of Philadelphia*, vol. 1, p. 1–134.
- Hikuroa, D. C. H. and Kaim, A., 2007: New gastropods from the Jurassic of Orville Coast, eastern Ellsworth Land, Antarctica. *Antarctic Science*, vol. 19, p. 115–124.
- Hilgen, F. J., Lourens, L. J. and Van Dam, J. A., 2012: The Neogene Period. In, Gradstein, F. M., Ogg, J. G., Schmitz, M. and Ogg, G., eds., *The Geologic Time Scale 2012*, p. 923–978. Elsevier, Amsterdam.
- Houbrick, R. S., 1978: The family Cerithiidae in the Indo-Pacific. Part 1: The genera *Rhinoclavis*, *Pseudovergatus* and *Clavocerithium*. *Monographs of Marine Mollusca*, no. 1, p. 1–130.
- Houbrick, R. S., 1981: Anatomy, biology and systematics of *Campanile symbolicum* with reference to adaptive radiation of the Cerithiace (Gastropoda: Prosobranchia). *Malacologia*, vol. 21, p. 263–289.
- Houbrick, R. S., 1984: The giant creeper, *Campanile symbolicum* Iredale, an Australian relict marine snail. In, Eldredge, N. and Stanley, S. M. eds., *Living Fossils*, p. 232–235. Springer-Verlag, New York.
- Houbrick, R. S., 1991: Systematic review and functional morphology of the mangrove snails *Terebralia* and *Telescopium* (Potamididae; Prosobranchia). *Malacologia*, vol. 33, p. 289–338.
- Houbrick, R. S., 1992: Monograph of the genus *Cerithium* Bruguière in the Indo-Pacific (Cerithiidae: Prosobranchia). *Smithsonian Contributions to Zoology*, no. 510, p. 1–211.
- Inoue, K., 2007: Batillariid and potamidid gastropods from the Middle Miocene Kukinaga Group, Tanegashima Island, southwest Japan, and their paleobiogeographic implications. *Paleontological Research*, vol. 11, p. 277–292.
- International Commission on Zoological Nomenclature, 1999: *International Code of Zoological Nomenclature, Fourth Edition*, 306 p. International Trust for Zoological Nomenclature, London.
- Iredale, T., 1917: More molluscan name-changes, generic and specific. *Proceedings of the Malacological Society of London*, vol. 12, p. 322–330.
- Itoigawa, J., 1978: Evidence of subtropical environment in the Miocene of Japan. *Bulletin of the Mizunami Fossil Museum*, no. 5, p. 7–22.
- Iwasaki, Y., 1970: A Miocene molluscan fauna in the Philippines. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 77, p. 205–228.
- Kanno, S., 1986: Revision of genus *Vicarya* (Gastropoda) from the Indo-Pacific region. *Bulletin of Joetsu University of Education*, vol. 5, p. 31–57.
- Kanno, S., Hashimoto, W., Aoki, N., Noda, H. and Alacantara, P. M., 1980: Stratigraphic horizons and geologic ages of the Philippine *Vicarya* (Gastropoda). *Geology and Palaeontology of Southeast Asia*, vol. 21, p. 135–161.
- Kase, T., 1984: *Early Cretaceous Marine and Brackish-water Gastropoda from Japan*, 263 p. National Science Museum, Tokyo.
- Kowalke, T., 2001: Cerithioidea (Caenogastropoda: Cerithiomorpha) of Tethyan coastal swamps and their relations to modern mangal communities. *Bulletin of the Czech Geological Survey*, vol. 76, p. 253–271.
- Kowalke, T., 2003: Verbreitung der Potamididae H. and A. Adams, 1854 (Caenogastropoda: Cerithiomorpha: Cerithioidea) im europäischen Känozoikum. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, vol. 227, p. 301–320.
- Lamarck, J. B. P. D. de M. de, 1804: Suite des mémoires sur les fossiles des environs de Paris. *Annales du Muséum National d'Histoire Naturelle*, vol. 3, p. 436–441.
- Leidy, J., 1851: Contributions to helminthology. *Proceedings of the Academy of Natural Sciences of Philadelphia*, vol. 5, p. 349–351.
- Lesport, J.-F. and Cahuzac, H., 2002: Sur un Potamididae méconnu du Miocène inférieur d'Aquitaine: *Pyrazisimus monstrosus* (Grateloup, 1847) [Mollusques Gastéropodes]; discussion générique. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, vol. 223, p. 1–52.
- Ludbrook, N. H., 1971: Large gastropods of the families Diastomatidae and Cerithiidae (Mollusca: Gastropoda) in southern Australia. *Transactions of the Royal Society of South Australia*, vol. 95, p. 29–42.
- Martin, K., 1890a: Ein neues *Telescopium* und die Beziehung dieser Gattung zu *Nerinea*. *Sammlungen des Geologischen Reichsmuseums in Leiden, Ite Serie*, vol. 4, p. 232–237.
- Martin, K., 1890b: Die Kei-Inseln und ihr Verhältniss zur Australisch-Asiatischen Grenzlinie, zugleich ein Beitrag zur Geologie von Timor und Celebes. *Tijdschrift Koninklijk Nederlandsch Aardrijkskundig Genootschap, Serie 2*, vol. 7, p. 241–280.
- Martin, K., 1899: Die Fossilien von Java, auf Grund einer Sammlung von Dr. R. D. M. Verbeek. Band 1, Gastropoda. Heft 6–8. *Sammlungen des Geologischen Reichs-Museum in Leiden, Neue Folge*, vol. 1, p. 133–220.
- Martini, E., 1971: Standard Tertiary and Quaternary calcareous nanoplankton zonation. In, Farinacci, A. ed., *Proceedings of the Second Planktonic Conference Roma 1970*, vol. 2, p. 739–785. Edizioni Tecnoscienza, Rome.
- Mines and Geosciences Bureau, 1983: *Geological Map of Danao Quadrangle, Sheet 3851 IV*. Ministry of Natural Resources, Philippines, Manila.
- Mines and Geosciences Bureau, 2010: *Geology of the Philippines, Second Edition*, 532 p. Department of Environment and Natural Resources, Quezon City.
- Nagao, T., 1934: Cretaceous Mollusca from the Miyako district, Honshu, Japan (Lamellibranchiata and Gastropoda). *Journal of the Faculty of Science, Hokkaido Imperial University, Series 4, Geology and Mineralogy*, vol. 2, p. 177–278.
- Nishihara, M., 1983: Grazing of the mangrove littoral by *Terebralia palustris* (Gastropoda: Potamididae) in the Okinawan mangal: Preliminary report. *Galaxea*, vol. 2, p. 45–58.
- Oostingh, C. H., 1938: Mollusken als gidsfossielen voor het Neogeen in Nederlandsch-Indië. *Handelingen van het Achste Nederland-*

- sch-Indisch Natuurwetenschappelijk Congres, Soerabaja, Juli 1938*, p. 508–516.
- Oyama, K., 1950: Studies of fossil molluscan biocoenosis. No. 1, bio-coenological studies on the mangrove swamps, with descriptions of new species from Yatsuo Group. *Report of Geological Survey of Japan*, no. 132, p. 1–16.
- Ozawa, T., Nakagawa, T. and Takeyama, K., 1986: Middle Miocene molluscan fauna of the Uchiura Group, Wakasa Province, Southwest Japan. *Palaeontological Society of Japan, Special Paper*, no. 29, p. 133–148.
- Piette, É., 1855: Observations sur les étages inférieurs du terrain jurassique dans les départements des Ardennes et de l'Aisne. *Bulletin de la Société Géologique de France, Série 2*, tome 12, p. 1083–1122.
- Piette, É., 1856: Notice sur les coquilles ailées trouvées dans la grande oolithe de l'Aisne, des Ardennes et de la Moselle. *Bulletin de la Société Géologique de France, Série 2*, tome 13, p. 85–100.
- Plaziat, J.-C., 1984: Mollusk distribution in the mangal. In, Por, F. D. and Dor, I. eds., *Hydrobiology of the Mangal*, p. 111–143. Dr. W. Junk, The Hague.
- Plaziat, J.-C., 1995: Modern and fossil mangroves and mangals: their climatic and biogeographic variability. In, Bosence, D. W. J. and Allison, P. A. eds., *Marine Palaeoenvironmental Analysis from Fossils*, p. 73–96. Geological Society Special Publication, no. 83. The Geological Society, London.
- Plaziat, J.-C., Cavagnetto, C., Koeniguer, J.-C. and Baltzer, F., 2001: History and biogeography of the mangrove ecosystem, based on a critical reassessment of the paleontological record. *Wetlands Ecology and Management*, vol. 9, p. 161–179.
- Rodelli, M. R., Gearing, J. N., Gearing, P. J., Marshall, N. and Sasekumar, A., 1984: Stable isotope ratio as a tracer of mangrove carbon in Malaysian ecosystems. *Oecologia*, vol. 61, p. 326–333.
- Sälgeback, J. and Savazzi, E., 2006: Constructional morphology of cerithiform gastropods. *Paleontological Research*, vol. 10, p. 233–259.
- Saul, L. E. and Squires, R. L., 2003: New Cretaceous cerithiform gastropods from the Pacific Slope of North America. *Journal of Paleontology*, vol. 77, p. 442–453.
- Sayn, G., 1932: Description de la faune de l'Urgonien de Barcelonne (Drôme). *Travaux du Laboratoire de Géologie de la Faculté des Sciences de Lyon*, fasc. 18, mémoire 15, p. 1–67.
- Schlotheim, E. F., 1820: *Die Petrefactenkunde auf ihrem jetzigen Standpunkte*, 437 p., 15 pl. Becker, Gotha.
- Shuto, T., 1978: Notes on Indonesian Tertiary and Quaternary gastropods mainly described by the late Professor K. Martin II. Potamididae and Cerithiidae. *Geology and Palaeontology of Southeast Asia*, vol. 19, p. 113–160, pls. 15–18.
- Slim, F. J., Hemminga, M. A., Ochieng, C., Jannink, N. T., Cocheret de la Morinière, E. and Velde, G. van der, 1997: Leaf litter removal by the snail *Terebralia palustris* (Linnaeus) and sesarmid crabs in an East African mangrove forest (Gazi Bay, Kenya). *Journal of Experimental Marine Biology and Ecology*, vol. 215, p. 35–48.
- Smith, W. D., 1913: Contributions to the stratigraphy and fossil invertebrate fauna of the Philippine Islands. *Philippine Journal of Science*, vol. 8, Section A, p. 235–300.
- Sowerby, G. B., Jun., 1855: Monograph of the genus *Cerithium*, Adanson. In, *Thesaurus Conchyliorum, or Monographs of Genera of Shells*, vol. 2, p. 847–899. Sowerby, London.
- Takeyama, T., 1933: Notes on the genus *Vicarya*, with description of two Japanese forms. *Japanese Journal of Geology and Geography*, vol. 10, p. 129–144.
- Tate, R., 1899: On some Older Tertiary fossils of uncertain age from the Murray Desert. *Transactions of the Royal Society of South Australia*, vol. 23, p. 102–111.
- Tenison-Woods, J. E., 1879: On some Tertiary fossils from Muddy Creek, Western Victoria. *Proceedings of the Linnean Society of New South Wales*, vol. 3, p. 222–240.
- Tian, J., Zhao, Q. H., Wang, P. X., Li, Q. Y. and Cheng, X. R., 2008: Astronomically modulated Neogene sediment records from the South China Sea. *Paleoceanography*, vol. 23, doi:10.1029/2007PA001552 PA3210.
- Tojo, B. and Masuda, F., 1999: Tidal growth patterns and growth curves of the Miocene potamidid gastropod *Vicarya yokoyamai*. *Paleontological Research*, vol. 3, p. 193–201.
- Tsuda, K., Itoigawa, J. and Yamanoi, T., 1986: Mangrove swamp fauna and flora in the Middle Miocene of Japan. *Palaeontological Society of Japan, Special Paper*, no. 29, p. 129–134.
- Vermeij, G. J., 2011: Shifting sources of productivity in the coastal marine tropics during the Cenozoic era. *Proceedings of the Royal Society B*, vol. 278, p. 2362–2368.
- Vermeij, G. J., 2012: The evolution of gigantism on temperate shores. *Biological Journal of the Linnean Society*, vol. 106, p. 776–793.
- Vredenburg, E., 1928: Descriptions of Mollusca from the post-Eocene Tertiary formation of north-western India: Gastropoda (in part) and Lamellibranchiata. *Memoirs of the Geological Survey of India*, vol. 50, p. 351–507.
- Wenz, W., 1940: Gastropoda. Teil 4. Prosobranchia. In, Schindewolf, O. H. ed., *Handbuch der Paläozoologie*, Band 6, Teil 1, p. 721–960. Gebrüder Borntraeger, Berlin.
- Williams, H. H. and Richards, D. H. H., 1968: Observations on *Pseudanakis rotundata* (Rudolphi, 1819) Mozgovoi 1950, a common but little known nematode parasite of *Raia radiata* Donovan in the northern North Sea. *Journal of Helminthology*, vol. 42, p. 199–220.