

## **Pteruthiidae and Erpornithidae (Aves: Corvides): two new family-group names for babbler-like outgroups of the vireos (Vireonidae)**

Authors: Sangster, George, Cibois, Alice, and Reddy, Sushma

Source: Bulletin of the British Ornithologists' Club, 142(2) : 239-243

Published By: British Ornithologists' Club

URL: <https://doi.org/10.25226/bboc.v142i2.2022.a8>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Pteruthiidae and Erpornithidae (Aves: Corvides): two new family-group names for babbler-like outgroups of the vireos (Vireonidae)

by George Sangster, Alice Cibois & Sushma Reddy

Received 14 December 2021; revised 4 March 2022; published 3 June 2022

<http://zoobank.org/urn:lsid:zoobank.org:pub:34181EAA-CC3E-490A-889D-4F5B32044D34>

The Old World genera *Erpornis* Hodgson, 1844, and *Pteruthius* Swainson, 1832, were long considered babblers Timaliidae (Deignan 1964, Sibley & Monroe 1990, Howard & Moore 1991). This was based on gross morphology (in *Erpornis*, similarity to *Yuhina*; in *Pteruthius*, bold plumage pattern reminiscent of, e.g., the laughingthrush genus *Garrulax*) and biogeography (presence alongside many 'other' species of babblers in the Oriental region), rather than phylogenetic analysis.

In the first molecular phylogenetic analysis of babblers, Cibois *et al.* (2002) found that the species known as '*Yuhina*' *zantholeuca* was not closely related to other species of *Yuhina* but was placed in an unresolved position near the crows *Corvus* and cuckooshrikes *Coracina*. As a consequence, the species was removed from *Yuhina* and placed in the monotypic genus *Erpornis* (Cibois *et al.* 2002). *Erpornis* was subsequently found to be closely related to the New World vireos Vireonidae (Barker *et al.* 2004). Another study showed that *Pteruthius* does not belong with the babblers but is sister to the *Erpornis*-Vireonidae clade (Reddy & Cracraft 2007). These findings were corroborated by Jönsson *et al.* (2016), Oliveros *et al.* (2019) and Stervander *et al.* (2020).

Following the discovery that *Erpornis* and *Pteruthius* represent successive sister groups of the vireos, they were lumped within an expanded Vireonidae in several taxonomies (Dickinson & Christidis 2014, Winkler *et al.* 2015, Fjeldså *et al.* 2020b). Placing *Erpornis* and *Pteruthius* in Vireonidae was probably necessitated by the lack of available family-group names for these two taxa. However, a good reason to refrain from placing *Erpornis* and *Pteruthius* into Vireonidae is that it changes the meaning of Vireonidae and affects several aspects of the systematics and biogeography of the family. For example, the time to their most recent common ancestor (MRCA) is shifted at least ten million years backwards (Stervander *et al.* 2020: fig. A2.1), the geographic origin of the MRCA shifts from the Americas to Asia, and the synapomorphies and diagnostic character states of 'Vireonidae' are modified. One can argue that these changes were unnecessary and potentially confusing. Given the long stability of the meaning of the name Vireonidae (e.g. Sharpe 1903, Wetmore 1930, Mayr & Amadon 1951, Deignan 1964, Wolters 1980, Sibley & Monroe 1990, Dickinson 2003), the alternative taxonomic solution, to exclude *Erpornis* and *Pteruthius* and place these in new family taxa, should be proposed. For example, Slager *et al.* (2014), who were aware of the close relationship of *Erpornis* and *Pteruthius* to the vireos, used the name Vireonidae in its traditional sense, i.e. for the New World taxa alone, excluding *Pteruthius* and *Erpornis*.

Placing *Erpornis*, *Pteruthius* and Vireonidae in separate family taxa is in line with the current classification of Passeriformes, as reviewed in Fjeldså *et al.* (2020a). Fjeldså *et al.* (2020b) followed the idea, within Passeriformes, of temporal banding (Avise & Johns 1999), in which the classification at higher ranks (i.e. families and genera) should correspond to groups of similar age, based on calibrated phylogenies. They did not, however, establish a strict cut-off line (as had Jönsson *et al.* 2016 for Corvides), knowing that confidence intervals of nodes are often wide and can vary among analyses (e.g. Oliveros *et al.* 2019, Stervander

TABLE 1  
Divergence times of *Pteruthius* and *Erpornis* from Vireonidae calculated in various dated phylogenies.

| <i>Pteruthius</i> | <i>Erpornis</i> | Source                          |
|-------------------|-----------------|---------------------------------|
| 27.5–27.7 Mya     | 22.4–22.5 Mya   | Reddy & Cracraft (2007)         |
| 23 Mya            | 21 Mya          | Jönsson <i>et al.</i> (2016)    |
| 15 Mya            | -               | Moyle <i>et al.</i> (2016)      |
| 17 Mya            | 12.5 Mya        | Oliveros <i>et al.</i> (2019)   |
| 27.5 Mya          | 21 Mya          | Stervander <i>et al.</i> (2020) |

TABLE 2  
Taxa treated as families (*sensu* Fjeldså *et al.* 2020b) that were inferred to be of equal or younger age than *Pteruthius*. Taxa marked with an asterisk were inferred to be younger than both *Pteruthius* and *Erpornis*.

| Chronogram                      | Taxa  |
|---------------------------------|---|
| Oliveros <i>et al.</i> (2019)   | Dendrocolaptidae, Xenopidae, Furnariidae, Malaconotidae, Pityriasisidae, Aegithinidae, Platysteiridae, Vangidae, Lamprolidae, Rhipiduridae, Ifridae, Paradisaeidae, Corvidae, Platylphidae, Laniidae, Paradoxornithidae, Sylviidae, Timaliidae, Leiothrichidae, Pellorneidae, Buphagidae, Sturnidae, Mimidae, Polioptilidae, Troglodytidae, Viduidae, Estrildidae, Rhodinocichlidae, Plectrophenacidae, Emberizidae, Cardinalidae, Mitrospingidae*, Thraupidae*, Passerellidae, Parulidae*, Icteridae*, Calyptophilidae* and Phaenicophilidae*.   |
| Stervander <i>et al.</i> (2020) | Furnariidae, Dendrocolaptidae, Xenopidae, Platyrhynchidae, Tachuridae, Rhynchocyclidae, Chaetopidae, Eupetidae, Dicuridae, Lamprolidae*, Rhipiduridae*, Corcoracidae*, Platylphidae*, Paradisaeidae*, Laniidae*, Corvidae*, Melampittidae, Ifridae*, Monarchidae*, Machaerirhynchidae, Artamidae*, Cracticidae*, Rhagologidae, Aegithinidae, Pityriasisidae*, Malaconotidae*, Platysteiridae*, Vangidae*, Sylviidae, Zosteropidae, Timaliidae, Pellorneidae, Leiothrichidae, Polioptilidae, Troglodytidae, Buphagidae, Sturnidae, Mimidae, Urocynchramidae, Ploceidae, Amblyospizidae, Viduidae*, Estrildidae*, Passeridae, Fringillidae, Plectrophenacidae*, Rhodinocichlidae*, Emberizidae*, Calyptophilidae*, Mitrospingidae*, Thraupidae*, Cardinalidae*, Passerellidae*, Phaenicophilidae*, Icteridae* and Parulidae*. |

*et al.* 2020). Most taxa treated as families by Fjeldså *et al.* (2020b), however, diverged around 20 Mya, a date similar to estimates for the *Erpornis* lineage and the *Pteruthius* clade (Table 1). In the chronogram of Oliveros *et al.* (2019), 38 taxa treated as families (*sensu* Fjeldså *et al.* 2020b) are of equal or younger age than *Pteruthius* and six of these are also of equal or younger age than *Erpornis* (Table 2). Similarly, in the unconstrained chronogram of Stervander *et al.* (2020), 56 taxa treated as families (*sensu* Fjeldså *et al.* 2020b) are of equal or younger age than *Pteruthius* and 28 of these are also of equal or younger age than *Erpornis* (Table 2).

Because no family-group names are available for *Erpornis* and *Pteruthius*, we propose:

**Erpornithidae new family**

**Type genus** *Erpornis* Hodgson, 1844

**Diagnosis** Small passerines (body length 12.0–13.5 cm) with rather plain plumage and distinct yellow vent and tail. Differs from all species of Vireonidae by combination of (i) short erect crest, (ii) uniform head without pale supercilium behind the eye or pale eye-ring, (iii) lack of wingbars, (iv) plain tertials and (v) pale pink legs.

Differs from all species of Pteruthiidae by combination of (i) slender bill, (i) short erect crest, (iii) lack of sexual dimorphism in plumage, (iv) greenish head and upperparts, (v) lack of a dark eyestripe and (vi) uniform greenish wings without wingbars.

**Included taxa** *Erpornis zantholeuca* Blyth, 1844.



## Pteruthiidae new family

**Type genus** *Pteruthius* Swainson, 1832

**Diagnosis** Small passerines (body length 11.5–21.0 cm) with short stout bill and colourful plumage (except the Green Shrike-Babbler *P. xanthochlorus* complex). Differs from Vireonidae by (i) sexual dimorphism in plumage and (ii) head pattern, with black head (in male Black-headed Shrike-Babbler *P. rufiventer*), black head with white supercilium (males of the Pied Shrike-Babbler *P. flaviscapis* complex), ear-coverts bordered by dark crescent and reddish throat (Black-eared Shrike-Babbler *P. melanotis* complex), dark lores, pale supercilium behind eye and reddish throat (Chestnut-fronted Shrike-Babbler *P. aenobarbus* complex), or plain greyish head with dark lores but no pale supercilium (*P. xanthochlorus* complex).

Differs from *Erpornis* by combination of (i) stout bill, (ii) lack of crest, (iii) sexual dimorphism in plumage, (iv) black or well-marked head in most species (except the *P. xanthochlorus* complex), and (v) wings either black with yellowish or chestnut tertials (in *P. flaviscapis* complex), black or dark brown with white to rufous-buff wingbars (*P. melanotis* and *P. aenobarbus* complexes), or greenish with paler wingbars (*P. xanthochlorus* complex).

**Included taxa** A single genus (*Pteruthius*) with nine or 19 species, depending on whether divergence in songs (Rheindt & Eaton 2009; Gill *et al.* 2021) or diagnosability of plumage (Reddy 2008), respectively, is used as the taxonomic criterion for species limits.

## Discussion

An alternative solution would be to treat Erpornithidae as a subfamily of Vireonidae, based on (i) the view that placement of the Asian *Erpornis* with the New World vireos conveys useful biogeographic information because it would emphasise the Asian origin of Vireonidae, (ii) the fact that in the phylogeny of Oliveros *et al.* (2019) only a small number of families are younger than Erpornithidae, and these are all in the nine-primaried New World oscines, and (iii) the age of Erpornithidae is younger than a large number of nodes within families in the phylogeny of Oliveros *et al.* (2019), which means that if temporal banding is applied to these groups, many new families should be named.

Ranking is obviously subjective, and therefore disagreements about the ‘appropriate’ rank cannot be solved empirically. In this case, ranking *Erpornis* as a subfamily and placing it in Vireonidae has two negative consequences: it results in an unnecessary change of the name of the vireos (‘Vireoninae’) and attaches the name ‘Vireonidae’ to a different, larger, older and geographically much more widespread group. We are not convinced by the arguments in favour of merging *Erpornis* with the vireos in a single family. First, merging the Asian *Erpornis* with the New World vireos in itself would not convey any information about the geographic origin of the vireos. Rather, the origin of a particular group, i.e. the direction of past geographic movement, is inferred from mapping the geographic range of multiple groups on a phylogeny. Second, whereas in the phylogeny of Oliveros *et al.* (2019) only six families were younger than Erpornithidae, and these were all in the nine-primaried New World oscines, in the phylogeny of Stervander *et al.* (2020) no fewer than 28 were of equal or younger age than *Erpornis* (Table 2), and these were not restricted to the nine-primaried New World oscines. Several nine-primaried New World oscine taxa were ranked as families by Barker *et al.* (2013) to keep the names of several major previously recognised groups stable (e.g. Thraupidae, Cardinalidae, Icteridae, Parulidae). We agree with this approach because it serves nomenclatural stability, even if it results in relatively ‘young’ family-level taxa. We have applied this approach to Erpornithidae

and Pteruthiidae, so that Vireonidae can be maintained for the vireos. Third, the fact that several lineages in other families are older than Erpornithidae does not imply that these should also be ranked as families. Only when temporal banding is enforced with a strict threshold will there be a need to change the rank of these taxa, and we believe this is not necessary. Ranking *Erpornis* as a family is in line with the current classification of Passeriformes, as reviewed in Fjeldså *et al.* (2020a), which is only loosely based on temporal banding.

### Acknowledgements

We thank Don Roberson for his encouragement to write this note, and two anonymous referees for their insightful comments.

### References

- Avise, J. C. & Johns, G. C. 1999. Proposal for a standardized temporal scheme of biological classification for extant species. *Proc. Natl. Acad. Sci. USA* 96: 7358–7363.
- Barker, F. K., Cibois, A., Schikler, P., Feinstein, J. & Cracraft, J. 2004. Phylogeny and diversification of the largest avian radiation. *Proc. Natl. Acad. Sci. USA* 101: 11040–11045.
- Barker, F. K., Burns, K. J., Klicka, J., Lanyon, S. M. & Lovette, I. J. 2013. Going to extremes: contrasting rates of diversification in a recent radiation of New World passerine birds. *Syst. Biol.* 62: 298–320.
- Bock, W. J. 1994. History and nomenclature of avian family-group names. *Bull. Amer. Mus. Nat. Hist.* 222: 1–281.
- Cibois, A., Kalyakin, M. V., Han, L.-X. & Pasquet, E. 2002. Molecular phylogenetics of babblers (Timaliidae): revaluation of the genera *Yuhina* and *Stachyris*. *J. Avian Biol.* 33: 380–390.
- Deignan, H. G. 1964. Timaliinae, Panurinae, Picathartinae. Pp. 240–442 in Mayr, E. & Paynter, R. A. (eds.) *Check-list of birds of the world*, vol. 10. Mus. Comp. Zool., Cambridge, MA.
- Dickinson, E. C. (ed.) 2003. *The Howard and Moore complete checklist of the birds of the world*. Third edn. Christopher Helm, London.
- Dickinson, E. C. & Christidis, L. (eds.) 2014. *The Howard and Moore complete checklist of the birds of the world*, vol. 2. Fourth edn. Aves Press, Eastbourne.
- Fjeldså, J., Christidis, L. & Ericson, P. G. P. (eds.) 2020a. *The largest avian radiation*. Lynx Edicions, Barcelona.
- Fjeldså, J., Christidis, L., Ericson, P. G. P., Stervander, M., Ohlson, J. I. & Alström, P. 2020b. An updated classification of passerine birds. Pp. 45–63 in Fjeldså, J., Christidis, L. & Ericson, P. G. P. (eds.) *The largest avian radiation*. Lynx Edicions, Barcelona.
- Gill, F., Donsker, D. & Rasmussen, P. (eds.) 2021. IOC world bird list (v11.2). doi: 10.14344/IOC.ML.11.2.
- Howard, R. & Moore, A. 1991. *A complete checklist of the birds of the world*. Second edn. Academic Press, London.
- Jönsson, K. A., Fabre, P.-H., Kennedy, J. D., Holt, B. G., Borregaard, M. K., Rahbek, C. & Fjeldså, J. 2016. A supermatrix phylogeny of corvid passerine birds (Aves: Corvidae). *Mol. Phyl. & Evol.* 94: 87–94.
- Mayr, E. & Amadon, D. 1951. A classification of recent birds. *Amer. Mus. Novit.* 1496: 1–42.
- Moyle, R. G., Oliveros, C. H., Andersen, M. J., Hosner, P. A., Benz, B. W., Manthey, J. D., Travers, S. L., Brown, R. M. & Faircloth, B. C. 2016. Tectonic collision and uplift of Wallacea triggered the global songbird radiation. *Nat. Comm.* 7: 12709.
- Oliveros, C. H., Field, D. J., Ksepka, D. T., Barker, F. K., Aleixo, A., Andersen, M. J., Alström, P., Benz, B. W., Braun, E. L., Braun, M. J., Bravo, G. A., Brumfield, R. T., Chesser, R. T., Claramunt, S., Cracraft, J., Cuervo, A. M., Derryberry, E. P., Glenn, T. C., Harvey, M. G., Hosner, P. A., Joseph, L., Kimball, R. T., Mack, A. L., Miskelly, C. M., Peterson, A. T., Robbins, M. B., Sheldon, F. H., Silveira, L. F., Smith, B. T., White, N. D., Moyle, R. G. & Faircloth, B. C. 2019. Earth history and the passerine superradiation. *Proc. Natl. Acad. Sci. USA* 116: 7916–7925.
- Reddy, S. 2008. Systematics and biogeography of the shrike-babblers (*Pteruthius*): species limits, molecular phylogenetics, and diversification patterns across southern Asia. *Mol. Phyl. & Evol.* 47: 54–72.
- Reddy, S. & Cracraft, J. 2007. Old World shrike-babblers (*Pteruthius*) belong with New World vireos (Vireonidae). *Mol. Phyl. & Evol.* 44: 1352–1357.
- Rheindt, F. E. & Eaton, J. A. 2009. Species limits in *Pteruthius* (Aves: Corvidae) shrike-babblers: a comparison between the Biological and Phylogenetic Species Concepts. *Zootaxa* 2301: 29–54.
- Sharpe, R. B. 1903. *A hand-list of the genera and species of birds*, vol. 4. Trustees of the Brit. Mus., London.
- Sibley, C. G. & Monroe, B. L. 1990. *Distribution and taxonomy of birds of the world*. Yale Univ. Press, New Haven, CT & London.
- Slager, D. L., Battey, C. J., Bryson, R. W., Voelker, G. & Klicka, J. 2014. A multilocus phylogeny of a major New World avian radiation: the Vireonidae. *Mol. Phyl. & Evol.* 80: 95–104.

- Stervander, M., Fjeldså, J., Christidis, L., Ericson, P. G. P., Ohlson, J. I. & Alström, P. 2020. An updated chronology of passerine birds. Pp. 387–396 in Fjeldså, J., Christidis, L. & Ericson, P. G. P. (eds.) *The largest avian radiation*. Lynx Edicions, Barcelona.
- Winkler, D. W., Billerman, S. M. & Lovette, I. J. 2015. *Bird families of the world*. Lynx Edicions, Barcelona.
- Wolters, H. E. 1980. *Die Vogelarten der Erde*. Lief. V. Paul Parey, Hamburg.

*Addresses:* George Sangster, Naturalis Biodiversity Center, Darwinweg 2, PO Box 9517, 2300 RA Leiden, the Netherlands, e-mail: g.sangster@planet.nl. Alice Cibois, Natural History Museum of Geneva, CP 6434, CH 1211 Geneva, Switzerland, e-mail: alice.cibois@ville-ge.ch. Sushma Reddy, Bell Museum of Natural History and Dept. of Fisheries, Wildlife, Conservation Biology, Univ. of Minnesota, Twin Cities, 2003 Upper Buford Circle, Saint Paul, MN 55108, USA, e-mail: sreddy@umn.edu

