

## **A checklist of lichens from Iraq, with four new records of lichens from Iraq**

Authors: Karim, Shram H., Salih, Salah A., and Al-Zubaidy, Adel A.

Source: Lindbergia, 2021(1)

Published By: Dutch Bryological and Lichenological Society and Nordic Bryological Society

URL: <https://doi.org/10.25227/linbg.01140>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# A checklist of lichens from Iraq, with four new records of lichens from Iraq

Shram H. Karim, Salah A. Salih and Adel A. Al-Zubaidy

S. H. Karim and S. A. Salih ✉ (salah.salh@spu.edu.iq), Dept of Natural Resources and Environmental Science, Sulaimani Polytechnic Univ., Sulaymaniyah, Iraq. – A. A. Al-Zubaidy, Horticulture and Landscape Design, Sulaimani Polytechnic Univ., Sulaymaniyah, Iraq.

The lichen species of Iraq are poorly known and no accurate checklist has been produced so far. Here we carried out an extensive review of existing literature and carried out additional field work to expand the knowledge of lichens in Iraq. We present the first checklist of lichens in Iraq which comprises a total of 236 species. The field survey identified four lichen species as new to Iraq. Although this checklist is not a full checklist of the lichens of Iraq it represents the most complete list to date. It is likely that increased sampling will lead to further new records of lichen species in Iraq.

Keywords: checklist, Halabja province, Iraq, lichens, new records

The lichen species of Iraq have been poorly documented for a long time and an accurate checklist to illustrate lichen flora of Iraq has been lacking. This is somewhat due to the fact that the exploration of the lichen flora has relied principally on the occasional visits of scientists visiting from abroad. The first recorded lichen species in Iraq were reported by Steiner (1921) who reported the lichen specimens collected by Handel-Mazzetti during his expedition to Mesopotamia, Kurdistan, Syria and Prinkipo. The majority of lichen species reported here was collected in the Mosul area. Later in 1969, 34 further lichen species were collected from Iraq by Schubert (1973). A study of biological soil crusts of the Middle East by Galun and Garty (2001) reported several species, one of which was *Diploschistes diacapsis*, that was previously reported to Iraq by Reichert (1940); and seven more terrestrial lichens (*Catapyrenium squamulosum*, *Collema crispum*, *Diploschistes diacapsis*, *Fulgensia bracteata*, *Psora decipiens*, *Squamarina lentigera* and *Toninia sedifolia*) studied in the north of Mesopotamia between Baghdad and Fallfar were also reported by Schubert (1973). Two further species of lichen were identified in Iraq by Şenkardeşler et al. (2014). A further contribution was made by Almola et al. (2017), with 37 lichens reported for first time from Iraq. Finally, another annotated list of lichens listed 30 lichen taxa as supplementary data (Aziz and Qadir 2016). Since a full

revision up-to-date list of lichen for Iraq is still lacking, a checklist is provided here. However, it must be indicated that the current paper also does not produce a complete account of lichen species. The checklist is a compilation of data from the previous studies listed above in combination with new fieldwork. Through carrying out new fieldwork we have added four new lichen records to Iraq. In this paper we describe the four new species reported in Iraq and provide a comprehensive checklist of lichen taxa known to occur in Iraq comprising 236 species in total.

## Material and methods

Lichen specimens were collected from Halabja province and Byara district located in the north of Iraq. During the survey lichen specimens were preserved in a paper bags with a piece of substrate to protect them until observations, identification and photographing the specimens could be carried out. Morphological characters were examined on dry material under a dissecting microscope. Then, we used the keys exists in the books and journals of lichens from Turkey and Iran (Davis 1970, Hale 1979, Goward et al. 1994 Brodo et al. 2001, Dobson 2005, Smith 2009, Coppins and Dobson 2012). The specimens are deposited in the herbarium of Sulaimani polytechnic university in the college of applied science in Halabja province.

In addition, all scientific publications containing information about the lichens of the Iraq have been critically investigated and utilized for compilation of the checklist. As far as practicable, original sources were checked in order

---

This work is licensed under the terms of a Creative Commons Attribution 4.0 International License (CC-BY) <<http://creativecommons.org/licenses/by/4.0/>>. The license permits use, distribution and reproduction in any medium, provided the original work is properly cited.




Figure 1. Collection localities of the survey Halabja province and Byara district (Iraq).

to remove those records which have arisen through misunderstanding of the original reference. The scientific publications used to show names of all lichens reported from Iraq are Steiner (1921), Reichert (1940), Schubert (1973), Şenkardeşler et al. (2014), Aziz and Qadir (2016) and Almola et al. (2017).

## Results

Through carrying out a literature review of previous lichen studies we found a total of 232 lichen species in Iraq. By carrying out additional fieldwork we found four new species of lichen occurring in Iraq, *Candelaria concolor*, *Dermatocarpon vellereum*, *Diploicia canescens* and *Lichinella nigritella*. In addition, we confirmed 17 of the species that were identified as occurring in Iraq from previous studies.

The present checklist below contains 236 species of lichen from Iraq. The list of lichens presented is a combination of published literature data with those species reported in the current paper. Full details of the newly recorded species are provided below. The species marked by an asterisk (\*) in the below list are new reports for Iraq and the species are marked with # also found in the study locations.

### The list of lichens from Iraq

*Acarospora cervina* A. Massal.  
*Acarospora fuscata* (Schrader) Arnold  
*Acarospora glaucocarpa* (Ach.) Körb.  
*Acarospora hiliaris* (Dufour) Hue  
*Acarospora impressula* Th. Fr.  
*Acarospora nodulosa* (Dufour) Hue

*Acarospora strigata* (Nyl.) Jatta  
*Amphoridium calcisedum* (DC.) Servit  
*Anaptychia desertorum* (Rupr.) Poelt #  
*Aspicilia caesiocinerea* (Nyl. ex Malbr.) Arnold  
*Aspicilia calcarea* (L.) Mudd #  
*Aspicilia candida* (Anzi) Hue  
*Aspicilia cinerea* (L.) Körb. #  
*Aspicilia contorta* (Hoffm.) Krempelh.  
*Aspicilia farinosa* Flörke) Flagey  
*Aspicilia ferruginea* (J. Steiner) Szatala  
*Aspicilia leprosenes* (Sandst.) Hav.  
*Aspicilia microspora* var. *insensibilis* Szatala  
*Aspicilia subpercaena* Szatala  
*Bellemerea aplina* (Sommerf.) Clauzade & Cl. Roux  
*Bellemerea cinereorufescens* (Ach.) Clauzade & Cl. Roux  
*Biatora vernalis* (L.) Fr.  
*Buellia epipolia* (Ach.) Mong.  
*Buellia spuria* (Schaer.) Anzi  
*Caloplaca vitellinula* (Nyl.) H. Olivier  
*Caloplaca aegyptiaca* (Müll. Arg.) J. Steiner  
*Caloplaca arenaria* (Pers.) Müll. Arg.  
*Caloplaca aurantia* (Pers.) Hellbom  
*Caloplaca biatorina* (A. Massal.) J. Steiner  
*Caloplaca bolacina* (Tuck.) H. Magn.  
*Caloplaca chalybaea* (Fr.) Müll. Arg.  
*Caloplaca citrina* (Hoffm.) Th. Fr. #  
*Caloplaca deceptoria* (Flagey) J. Steiner  
*Caloplaca aegyptiaca* (Müll. Arg.) J. Steiner  
*Caloplaca erythrocarpa* (Pers.) Zwackh  
*Caloplaca feracissima* H. Magn.  
*Caloplaca flavorubescens* (Hudson) J. R. Laundon  
*Caloplaca fulgens* Körb.  
*Caloplaca holocarpa* (Hoffm. ex Ach.) M. Wade

*Caloplaca ignea* Arup  
*Caloplaca inspersa* (J. Steiner) J. Steiner  
*Caloplaca lithophila* H. Magn.  
*Caloplaca murorum* (Hoffm.) Th. Fr.  
*Caloplaca ochracea* (Schaer.) Flagey  
*Caloplaca paepalostoma* (Anzi) Jatta  
*Caloplaca pelloidella* (Nyl.) Hasse  
*Caloplaca polycarpoides* (J. Steiner) M. Steiner & Poelt  
*Caloplaca pyracea* (Ach.) Zwackh  
*Caloplaca thallincola* (Wedd.) Du Rietz  
*Caloplaca variabilis* (Pers.) Müll. Arg.  
*Caloplaca velana* (A. Massal.) Du Rietz  
*Caloplaca verruculifera* (Vain.) Zahlbr.  
***Candelaria cornicolor* (Dickson) Stein \***  
*Candelariella aurella* (Hoffm.) Zahlbr. #  
*Candelariella vitellina* (Hoffm.) Müll. Arg.  
*Candelariella xanthostigma* (Ach.) Lettau  
*Catapyrenium squamulosum* (Ach.) Breuss  
*Coccocarpia erythroxyli* (Sprengel) Swinscow & Krog  
*Collema crispum* (Hudson) Weber ex F.H. Wigg.  
*Collema cristatum* (L.) F. H. Wigg.  
*Collema flaccidum* (Ach.) Ach.  
*Collema fuscovirens* (With.) J. R. Laundon  
*Collema ligerinum* (Hy) Harm.  
*Collema polycarpon* Hoffm.  
*Collema tenax* (Sw.) Ach.  
*Dermatocarpon hepaticum* (Ach.) Th. Fr.  
*Dermatocarpon miniatum* (L.) W. Mann  
*Dermatocarpon monstrosus* (Schaer.) Vain.  
***Dermatocarpon vellereum* Zschacke \***  
*Dimelaena oreina* (Ach.) Norman  
***Diploicia canescens* (Dickson) A. Massal. \***  
*Diploschistes steppicus* Reichert  
*Diploschistes caesioplumbeus* (Nyl.) Vain.  
*Diploschistes diacapsis* (Ach.) Lumbsch  
*Diploschistes muscorum* (Scop.) R. Sant.  
*Diploschistes ocellatus* (Fr.) Norman  
*Diploschistes scruposus* (Schreber) Norman  
*Diploschistes steppicus* Reichert  
*Diplotomma bedinii* (H. Magn.) P. Clerc & Cl. Roux  
*Dirina catalinae* Hasse  
*Endocarpon pusillum* Hedwig  
*Fulgensia bracteata* (Hoffm.) Räsänen  
*Fulgensia fulgens* (Sw.) Elenkin  
*Fulgensia schistidii* (Anzi) Poelt  
*Fulgensia subbracteata* (Nyl.) Poelt  
*Graphina peplophora* M. Wirth & Hale  
*Graphis elegans* (Borrer ex Sm.) Ach.  
*Gypsoplaca macrophylla* (Zahlbr.) Timdal  
*Haematomma ventosum* (L.) A. Massal. #  
*Heterodermia crocea* R.C. Harris  
*Heterodermia diademata* (Taylor) D. D. Awasthi  
*Hypogymnia austerodes* (Nyl.) Räsänen #  
*Hypotrachyna livida* (Taylor) Hale  
*Kaernefeltia merrillii* (Du Rietz) Thell & Goward  
*Lecanactis abietina* (Ach.) Körb.  
*Lecanora dispersa* (Pers.) Sommerf.  
*Lecanora hybocarpa* (Tuck.) Brodo  
*Lecanora mutabilis* (Ach.) Nyl.  
*Lecanora xylophila* Hue  
*Lecania koerberiana* J. Lahm  
*Lecania subcaesia* (Nyl.) B. de Lesd.  
*Lecanora albescens* (Hoffm.) Branth & Rostrup #  
*Lecanora argopholis* (Ach.) Ach.  
*Lecanora cinereofusca* var. *appalachensis* Brodo  
*Lecanora circinata* (Pers.) Ach.  
*Lecanora circumborealis* Brodo & Vitik.  
*Lecanora dispersa* (Pers.) Sommerf.  
*Lecanora garovaglii* (Körb.) Zahlbr.  
*Lecanora hoffmannii* (Ach.) Müll. Arg.  
*Lecanora Kurdistanica* J. Steiner  
*Lecanora lentigera* (Weber) Ach.  
*Lecanora muralis* (Schreber) Rabenh. #  
*Lecanora novomexicana* H. Magn.  
*Lecanora polytropa* (Hoffm.) Rabenh.  
*Lecanora pruinosa* Chaub.  
*Lecanora versicolor* (Pers.) Ach.  
*Lecidea atrobrunnea* ((Raymond ex Lam. & DC.) Schaer.  
*Lecidea cyathoides* (Ach.) Ach.  
*Lecidea deeptoria* Nyl.  
*Lecidea fuscoatra* (L.) Ach. #  
*Lecidella stigmathea* (Ach.) Hertel & Leuckert  
*Lepraria incana* (L.) Ach. #  
*Lepraria vouauxii* (Hue) R.C. Harris  
*Leptotrema wightii* (Taylor) Müll. Arg.  
*Lichinella cribellifera* (Nyl.) Henssen  
***Lichinella nigritella* (Lettau) P.P. Moreno & Egea \***  
*Lobothallia praevalida* (Nyl.) Hafellner  
*Megaspora verrucosa* (Ach.) Hafellner & V. Wirth  
*Melanelia disjuncta* (Erichsen) Essl.  
*Melanelia glabra* (Schaer.) Essl.  
*Melanelia panniformis* (Nyl.) Essl.  
*Mycoblastus affinis* (Schaer.) Schauer  
*Neofuscelia pulla* (Ach.) Essl.  
*Normandina pulchella* (Borrer) Nyl.  
*Ochrolechia frigida* (Sw.) Lynge  
*Ochrolechia pallescens* (L.) Körb.  
*Ochrolechia tartarea* (L.) A. Massal.  
*Orphniospora moriopsis* (A. Massal.) D. Hawksw.  
*Pachyospora verrucosa* (Ach.) Massal.  
*Pannaria rubiginosa* (Thunb.) Delise  
*Pannaria conoplea* (Ach.) Bory  
*Parmelia acetabulum* (Neck.) Duby  
*Parmelia caperata* (L.) Ach.  
*Parmelia conspersa* Ach.  
*Parmelia quercina* (Willd.) Vain. #  
*Parmelia saxatilis* (L.) Ach.  
*Parmelia tinctina* Maheu & A. Gillet  
*Parmelina quercina* (Willd.) Hale  
*Parmelina tiliacea* (Hoffm.) Hale  
*Parmotrema chinense* (Osbeck) Hale & Ahti  
*Parmotrema perforatum* (Jacq.) A. Massal.  
*Peltula obscurans* (Nyl.) Gyelnik  
*Peltula richardsii* (Herre) Wetmore  
*Pertusaria aspergilla* (Ach.) J.R. Laundon  
*Pertusaria flavicunda* Tuck.  
*Pertusaria lactea* (L.) Arnold #  
*Pertusaria paratuberculifera* Dibben  
*Pertusaria rubefacta* Erichsen  
*Pertusaria subambigens* Dibben  
*Pertusaria subpertusa* Brodo  
*Pertusaria texana* Müll. Arg.



*Pertusaria trachythallina* Erichsen  
*Pertusaria velata* (Turner) Nyl.  
*Phaeophyscia orbicularis* (Necker) Moberg  
*Physcia aipolia* (Ehrh. ex Humb.) Furnr. #  
*Physcia biziana* (A. Massal.) Zahlbr. #  
*Physcia caesia* (Hoffm.) Furnr.  
*Physcia perisidiosa* Erichsen  
*Physcia pulverulenta* (Schreb.) Hampe ex Fürnr.  
*Physcia semipinnata* (J. F. Gmelin) Moberg  
*Physcia thomsoniana* Essl.  
*Physconia distorta* (With.) J. R. Laundon  
*Physconia distorta* (With.) J. R. Laundon  
*Placidium lacinulatum* (Ach.) Breuss  
*Placocarpus schaereri* (Fr.) Breuss  
*Porpidia flavocaerulescens* (Hornem.) Hertel & A. J. Schwab  
*Psora decipiens* (Hedwig) Hoffm.  
*Psora lurida* (With.) DC.  
*Psora vallesiaca* (Schaer.) Timdal  
*Pyrenula nitida* (Weigel) Ach.  
*Pyrenula pseudobufonia* (Rehm) R.C. Harris  
*Rhizocarpon chioneum* (Norman) Th. Fr.  
*Rhizocarpon chionophilum* Th. Fr.  
*Rhizocarpon disporum* (Naeg. ex Hepp) Müll.Arg.  
*Rhizocarpon geographicum* (L.) DC.  
*Rhizocarpon hochstetteri* (Körb.) Vain.  
*Rhizocarpon macrosporium* Räsänen  
*Rhizocarpon obscuratum* (Ach.) A. Massal.  
*Rhizocarpon reductum* Th. Fr.  
*Rhizocarpon richardii* (Lamy ex Nyl.) Zahlbr.  
*Rhizocarpon umbilicatum* (Ramond) Flagey  
*Rhizoplaca melanophthalma* (DC.) Leuckert & Poelt  
*Rhizoplaca marginalis* (Hasse) W.A. Weber  
*Rhizoplaca subdiscrepans* (Nyl.) R. Sant.  
*Rinodina ascociscana* (Tuck.) Tuck.  
*Rinodina atrocinerea* (Fr.) Körb.  
*Rinodina bischoffii* (Hepp) A. Massal.  
*Rinodina immersa* (Körb.) Zahlbr.  
*Rinodina bolanderi* H. Magn.  
*Rinodina populicola* H. Magn.  
*Rinodina biechoffii* (Hepp) A. Massal.  
*Roccellina conformis* Tehler  
*Sarcogyne regularis* Körb.  
*Squamarina cartilaginea* (With.) P. James  
*Squamarina crassa* (Huds.) Poelt  
*Squamarina lentigera* (Weber) Poelt  
*Squamarina stella-petraea* Poelt  
*Staurothele diffractella* (Nyl.) Tuck.  
*Tephromela grumosa* (Pers.) Hafellner & Cl. Roux #  
*Thelidium pyrenophorum* (Ach.) Mudd  
*Thelomma californicum* (Tuck.) Tibell  
*Toninia alutacea* (Anzi) Jatta  
*Toninia candida* (Weber) Th. Fr.  
*Toninia sedifolia* (Scop.) Timdal  
*Toninia coeruleonigricans* (Lightf.) Th. Fr.  
*Trapelia glebulosa* (Sm.) J. R. Laundon  
*Trapelia placodioides* Coppins & P. James #  
*Trapeliopsis wallrothii* (Flörke) Hertel & Gotth. Schneider  
*Verrucaria amphibia* Clemente  
*Verrucaria attica* J. Steiner  
*Verrucaria baldensis* A. Massal. #  
*Verrucaria beltraminiana* (A. Massal.) Trevis.

*Verrucaria calcivora* Nyl.  
*Verrucaria calciseda* DC.  
*Verrucaria dolomitica* (A. Massal.) Kremp.  
*Verrucaria glaucina* Ach.  
*Verrucaria macrostoma* Dufour ex DC.  
*Verrucaria marmorea* (Scop.) Arnold  
*Verrucaria maura* Wahlenb.  
*Verrucaria nigrescens* Pers.  
*Verrucaria rupestris* Schrader  
*Verrucaria tristis* (A. Massal.) Kremp.  
*Xanthoparmelia plittii* (Gyelnik) Hale  
*Xanthoria elegans* (Link) Th. Fr.  
*Xanthoria parietina* (L.) Th. Fr.  
*Xanthomendoza fallax* (Hepp ex Arn.) Søchting, Kärnefelt & S.Y. Kondr.  
*Xanthomendoza fulva* (Hoffm.) Søchting, Kärnefelt & S.Y. Kondr.

## New record species for Iraq

### *Candelaria concolor* (Dickson) Stein

Found on tree barks. The lichen thallus is typically forming small, irregularly spreading discrete suborbicular, sometimes coalescing or in scattered fragments. The thallus appressed, lobes flattened, finely divided, surface flat or some wavy, often somewhat raised and fan-like, the margins distinctly overlapping, entire. The color of thallus lemon yellow to mustard yellow, paling to yellow green in shade and almost white underside. The rhizines scattered, white to pinkish. The species is distinguished by its microfoliose thallus which form distinct rosettes usually with moss habitat on the bark.

Note: the most conspicuous features for the *Candelaria concolor* species were seen on the lower surface, which is white, smooth and does not show any arachnoid structures as in *Candelaria pacifica*. Poelt (1974) described the lower cortex of *Candelaria concolor* as a 3–4 cell thick paraplectenchyma, built of numerous hyphae and with bundles of white rhizines. Furthermore, the production of ascomata could be observed in several specimens having polyspored asci. Above all, the lower surface gives the impression of a compact cortex, in contrast to the arachnoid lower surface in *Candelaria pacifica*. This species was distributed in Byara.

### *Dermatocarpon vellereum* Zschacke

Found on huge stones. The lichen thallus usually monophyllous, foliose, umbilicate, saxicolous, leathery, upper side light brownish to brownish red, white to dark pruinose, lower side black, with dense, thick stumpy, coralloid rhizinomorphs.

Note: widespread in subtropical to lower temperate regions China and eastern Europe, India, Nepal. This species is distributed also in Iraq. This species is located in Byara and Sargat, and it was common in study locations.

### *Diploicia canescens* (Dickson) A. Massal.

Found on stones. The thallus of lichen is crustose, placodioid, continuous, forming rosettes, effigurate lobes: radiating, but sometimes irregular, discrete or confluent; lobe tips: usually rotund or truncate. The surface is white, grey-white or pale blue-grey, plane to convex or concave at the periphery.

Note: this species it has two subspecies which are *Diploicia canescens* subsp. *australasica* and *Diploicia canescens* subsp.

*Canescens*. Moon et al. (2014) described the *Diploicia canescens* subsp. *australasica* is mainly restricted to the Southern Hemisphere with its center of distribution in Australia and New Zealand, whereas *Diploicia canescens* subsp. *canescens* is widely distributed in the Northern Hemisphere (Bratt 1984, Elix et al. 1988, Dalby 1994, Calatayud et al. 1995, Sipman 2002). The distribution of subsp. *australasica* now extends to Korea. According to Ryan et al. (2004), *Diploicia canescens* is distributed widely in the world including temperate Europe, Mediterranean region, Africa, Hawaii, Japan, Australasia, and western North America. This species also found around Halabja city.

#### ***Lichinella nigritella* (Lettau) P.P. Moreno & Egea**

Found on rocks. The thallus of lichen is black; the lobes of thallus are repeatedly branched and ascending with only the marginal lobes adpressed. The lobe shape is rather variable depending on the stage of branching. Because the lobe presence of numerous globose to scale-like isidia, there surface is rough. The broad, marginal lobes are often ridged.

Note: *Lichinella nigritella* is closely related to *Lichinella cribellifera* which is similar in overall appearance. However, *Lichinella nigritella* in the more narrow and more erect lobes which are deeply branched from the beginning, usually bearing numerous, small, globose or scalelike isidia. Schultz (2005) reports *Lichinella cribellifera* to be restricted to acidic rock, whereas *Lichinella nigritella* was found on acidic as well as calcareous rocks in North America. Also he said both species are widely distributed in the study region, but *Lichinella nigritella* appears to be more common.

For separation from *Lichinella cribellifera* that mature thallus, broad marginal lobe with ascending margin and numerous granules on the surface and the thallus consisting of small rosette with rounded, sparsely branched lobes and numerous, globose isidia on the lobe surface. We found both species in the study area, but *Lichinella cribellifera* is reported to Iraq previously. *Lichinella nigritella* is also distributed around Halabja city.

## **Discussion**

The known species diversity of Iraq at the moment includes 236 species of lichens. Although not all of provinces of Iraq were studied equally, some features of its diversity can be discussed. According to the limited data, the most lichens diverse in Iraq are north of Iraq (Kurdistan region). This seems plausible, since this mountainous region has the greatest diversity of natural conditions, including excellent substrate diversity, with a long geological history. The north of Iraq is also a diverse mountain territory. It is expected that the known lichen flora of this region will increase further through further targeted surveying. As presence of lichen species have still not yet been completely reported in all Iraq districts, our knowledge of lichens stimulated us to do this literature review on lichen species of Iraq, and to conduct surveys in Halabja province and Byara. Therefore, this comprehensive checklist has been made. This checklist is a guide to the presence of lichens in Iraq. The aim of making the list was to follow development records as to their origin and to correct mistaken data that caused by misidentification

and incorrect synonymy as far as possible. The old names are revised also into up-to date nomenclature. In our list we have gathered only records that were reliable or at least probably correct.

The increasing importance of lichens in the management and conservation of vegetation systems in Iraq has generated considerable local demand for a checklist of Iraq. However, it must be indicated that the current paper also does not produce a complete account of lichen species. The checklist is a blend of data from recent studies and old studied data gathered.

**Acknowledgements** – We are grateful to Dr. Alison Eyres from the United Kingdom for the technical and linguistic support, and valuable comments. We would like also to thank college of applied science at Halabja province for their kind help to use laboratory equipments such microscopes and other tools.

## **References**

- Almola, Z. S., Al-Ni'ma, B. A. and Ramadan, N. A. 2017. Lichen diversity in Amadiya and Rowanduz districts in Iraq. – Bangla. J. Plant Taxon. 24: 23–32.
- Aziz, F. H. and Qadir, S. B. 2016. Common and new records of lichens from Iraqi Kurdistan Region. – Iraq. Bull. Iraq Nat. Hist. Mus. 14: 51–68.
- Bratt, C. C. 1984. *Diploicia canescens* (Dicks.) Mass. new to North America. – Bryologist 87: 160–161.
- Brodo, I. M., Sharnoff, S. D. and Sharnoff, S. 2001. Lichens of north America. – Yale Univ. Press.
- Calatayud, V., Atienza, V. and Barreno, E. 1995. Lichenicolous fungi from the Iberian Peninsula and the Canary Islands. – Mycotaxon 55: 363–382.
- Coppins, S. and Dobson, F. S. 2012. Lichens. An illustrated guide to the British and Irish species. – Lichenologist 44: 563.
- Dalby, D. H. 1994. *Diploicia canescens* (Dicks.) A. Massal. in Shetland. – Bot. J. Scotland 47: 123–128.
- Davis, P. H. 1970. Flora of Turkey and the East Aegean Islands, Vol. 3.
- Dobson, F. S. 2005. Lichens: an illustrated guide to the British and Irish species. – Richmond Publishing.
- Elix, J. A., Jenkins, G. A. and Lumbsch, H. T. 1988. Chemical variation in the lichen genus *Diploicia* (Ascomycotina). – Mycotaxon 33: 457–466.
- Galun, M. and Garty, J. 2001. Biological soil crusts of the Middle East. – In: Biological soil crusts: structure, function and management. Springer, pp. 95–106.
- Goward, T., McCune, B. and Meidinger, D. 1994. The lichens of British Columbia. Illustrated keys. Part 1: 1–181.
- Hale, M. E. 1979. How to know the lichens. – WC Brown Co.
- Moon, K. H., Aptroot, A., Elix, J. A. et al. 2014. *Diploicia canescens* subsp. *australasica* Caliciaceae found in Korea. *Diploicia canescens* subsp. *australasica* Caliciaceae found in Korea. – 89: 51–53.
- Poelt, J. 1974. Zur Kenntnis der Flechtenfamilie Candelariaceae. Ein Beitrag mit besonderer Berücksichtigung einiger südamerikanischer Arten. – Phytion 16: Fasc. 1–4, 189–210.
- Reichert, I. 1940. A new species of *Diploschistes* from oriental steppes and its phytogeographical significance. – Palestine J. Bot. Rehovot Ser. 3: 162–182.
- Ryan, B. D., Lumbsch, H. T., Messuti, M. I. et al. 2004. Lichen flora of the Greater Sonoran Desert region. – Lichens Unlimited, p. 357.
- Schubert, R. 1973. Notizen zur Flechtenflora des nördlichen Mesopotamien (Irak) Mit 2 Abbildungen. – Feddes Rep. 83: 585–589.

- Schultz, M. 2005. An overview of *Lichinella* in the southwestern United States and northwestern Mexico, and the new species *Lichinella granulosa*. – *Bryologist* 108: 567–590.
- Şenkardeşler, A., Lőkös, L. and Farkas, E. 2014. Lectotypification of names of lichen taxa described by Ödön Szatala. – *Taxon* 63: 139–145.
- Sipman, H. J. M. 2002. Lichens of mainland Yemen. – *Willdenowia* 32: 127–135.
- Smith, C. W. 2009. Lichens of Great Britain and Ireland. – British Lichen Society.
- Steiner, J. 1921. Lichenes aus Mesopotamien und Kurdistan sowie Syrien und Prinkipo. – *Ann. Naturhistorischen Museums Wien*, pp. 1–68.