

Euro Med-Checklist Notulae, 7

Authors: Raab-Straube, Eckhard Von, and Raus (ed.), Thomas

Source: Willdenowia, 47(1) : 89-96

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.47.47112>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Notulae ad floram euro-mediterraneam pertinentes No. 36

ECKHARD VON RAAB-STRAUBE¹ & THOMAS RAUS¹ (ed.)

Euro+Med-Checklist Notulae, 7

Version of record first published online on 23 March 2017 ahead of inclusion in April 2017 issue.

Abstract: This is the seventh of a series of miscellaneous contributions, by various authors, where hitherto unpublished data relevant to both the Med-Checklist and the Euro+Med (or Sisyphus) projects are presented. This instalment deals with the families *Amaranthaceae*, *Compositae*, *Cruciferae*, *Cucurbitaceae*, *Orchidaceae*, *Orobanchaceae* and *Solanaceae*. It includes new country and area records, taxonomic and distributional considerations for taxa in *Amaranthus*, *Bidens*, *Datura*, *Echinocystis*, *Erigeron*, *Orobanche*, *Platanthera*, *Senecio* and *Tauscheria*, and the validation of four new combinations in *Phelipanche*.

Key words: Euro+Med PlantBase, Med-Checklist, Europe, Mediterranean, vascular plants, distribution, taxonomy, new record, new combination

Article history: Contributions received 15 December 2016 to 28 February 2017; peer-review completed 10 March 2017; received in revised form 13 March 2017; accepted for publication 14 March 2017.

Citation: Raab-Straube E. von & Raus Th. (ed.) 2017: Euro+Med-Checklist Notulae, 7 [Notulae ad floram euro-mediterraneam pertinentes No. 36]. – Willdenowia 47: 89–96. doi: <https://doi.org/10.3372/wi.47.47112>

Notice

A succinct description of the Euro+Med project, with a list of recognized territories and their abbreviations, and the conventions used to indicate the status and presence of taxa, can be found in the introduction to the first instalment of the Euro+Med Notulae (Greuter & Raab-Straube 2005: 223–226) and on the Euro+Med PlantBase website (Euro+Med 2006+). For the previous instalment of the Euro+Med-Checklist Notulae, see Raab-Straube & Raus (2016).

The following have contributed entries to the present instalment: N. M. G. Ardenghi, K. P. Buttler, G. Domina, E. Eleftheriadou, R. El Mokni, Th. Giannakis, Ya. M. Golovanov, P. Hein, D. Iamónico, M. S. Knyazev, A. Yu.

Korolyuk, E. von Raab-Straube, S. Rätzel, Th. Raus, J. Svensson, K. Theodoropoulos, R. Tsandekidis, S. Tsiftsis and S. M. Yamalov.

Amaranthaceae

Amaranthus emarginatus subsp. *pseudogracilis* (Thell.) Hügin

A Cr: Greece, Crete: Nomos of Chania, Eparchia of Apokoronos, Georgioupoli, 35°21'55"N, 24°15'40"E, 1 m, sandy bank of a freshwater stream by the harbour, 2 Jun 2015, Svensson (B). – A xenophyte of tropical origin not previously reported from the Cretan area (Strid 2016b: 46, map 151). The subspecies, easily

¹ Botanischer Garten und Botanisches Museum Berlin, Freie Universität Berlin, Königin-Luise-Str. 6–8, 14195 Berlin, Germany; *e-mail: e.raab-straub@bgbm.org (author for correspondence), t.raus@bgbm.org

recognizable by its ascending growth form and elongated terminal part of the inflorescence, is new for the whole of Greece. Hitherto known Greek occurrences refer to *Amaranthus emarginatus* subsp. *emarginatus* (for details, see Raus 1997: 145). The infraspecific taxonomy of *A. emarginatus* (Hügin 1987) was left unmentioned by Dimopoulos & al. (2013) and was even unknown to Iamonico (2015).

Th. Raus & J. Svensson

Compositae (Asteraceae)

Bidens pilosa L.

A **Cr**: Greece, Crete: Nomos of Chania, Eparchia of Kitionia, Agia, 35°28'20"N, 23°55'53"E, 40 m, wet ditch by roadside, 2 Jun 2015, *Svensson* (B). – First record for the Cretan area of this xenophyte of North American origin. In Greece otherwise known as an established weed of ruderal places in N Peloponnisos (*Garnweidner* 1989: 168; *Chronopoulos & Christodoulakis* 1996: 166) and on the island of Rodos (*Galanos* 2015: 270; *Kleinstauber & al.* 2016: 154).

Th. Raus & J. Svensson

Erigeron sumatrensis Retz. (= *Conyza albida* Spreng.)

N **Tn**: Tunisia: Bizerta, 37°16'65.0"N, 09°52'13.9"E, 3 m, harbour construction site, 27 Oct 2010, *El Mokni* (herb. El Mokni); Monastir, 35°45'50.4"N, 10°49'53.4"E, 11 m, public gardens, 7 Dec 2016, *El Mokni* (HFLA, herb. El Mokni). – *Erigeron sumatrensis* is a species presumably native to South America and widely distributed as a weed in tropical and subtropical regions worldwide (*Chen & Brouillet* 2011). This species is currently recorded in N Africa in all countries except Libya and Egypt (see, e.g., *Greuter* 2006+; *SANBI* 2012). However, concerning Tunisia, no status of naturalization appears to have been defined (see also *Le Floch & al.* 2010) and the Euro+Med PlantBase (*Greuter* 2006+) reports "Alien (status unknown)". As a consequence, a specification about the alien status of *E. sumatrensis* in Tunisia is needed. We found many populations identifiable as *E. sumatrensis* in the localities of Ain Draham, Beja, Bizerta, Boussalem, Bulla regia, Farnana, Hammamet, Jendouba, Monastir, Nefza and Tabarka, where they grow in human-made habitats (roadsides, cultivated lands, public gardens, walls) and occupy areas of 5–100 ha. All these populations were observed since the year 2004, so the species can be considered as naturalized in Tunisia according to the definitions by *Pyšek & al.* (2002).

D. Iamonico & R. El Mokni

Senecio noëanus Rupr.

+ **Rf(E)**: Russia: Orenburg region, Sol'-Iletsk district, Subural plateau, Verkhnechekhendinsky cretaceous mountains, 10 km W of Troitsk village, 50.683463°N, 54.470749°E, 6 Jun 2016, *Golovanov* (UFA; det. M. S. Knyazev); Akbulaksky district, Subural plateau, Korsak-Bas mountain, 6 km NE of abandoned village of Chagan, 50.702620°N, 55.764822°E, clay habitat at foot of mountain, 26 May 2016, *Korolyuk & al.* (NS, UFA; det. M. S. Knyazev). – In Russia, this species is widespread in the Lower Volga region, the Saratov region (*Davidenko & al.* 2016) and the Caucasus, where it grows on sandy and pebbly shores of rivers and lakes, in steppes and semi-deserts and also on saline soils. Outside Russia, it occurs in Asia Minor, Iran and C Asia (*Konechnaya* 1994). *Senecio noëanus* is new for the flora of the Orenburg region and the east of the European part of Russia. The nearest localities are reported in the Aktyubinsk region of Kazakhstan (*Aypeisova* 2012).

Ya. M. Golovanov, S. M. Yamalov,
A. Yu. Korolyuk & M. S. Knyazev

Cruciferae (Brassicaceae)

Tauscheria lasiocarpa DC.

+ **Rf(E)**: Russia: Orenburg region, Akbulaksky district, Subural plateau, Korsak-Bas mountain, 6 km NE of abandoned village of Chagan, 50.702685°N, 55.764307°E, clay and broken stone mountain slopes, 26 May 2016, *Korolyuk & al.* (UFA; det. M. S. Knyazev). – *Tauscheria lasiocarpa* occurs in Russia along the Lower Volga (Volgograd region), and also in the south of W Siberia, in steppes and on slopes with saline and clay habitats. Outside Russia, it occurs in Afghanistan, China, Iraq, Kazakhstan, Kyrgyzstan, Mongolia, Pakistan, Tajikistan, Turkmenistan and Uzbekistan (*Kotov* 1979; *Dorofeyev* 2002). Although given for the Orenburg region (*Ryabinina & Knyazev* 2009), the species was not substantiated by herbarium material, and the collection cited above is the first secure record in S Ural and the east of the European part of Russia. The nearest localities are reported in the Aktyubinsk region of Kazakhstan (*Aypeisova* 2012).

Ya. M. Golovanov, S. M. Yamalov,
A. Yu. Korolyuk & M. S. Knyazev

Cucurbitaceae

Echinocystis lobata (Michx.) Torr. & A. Gray

N **Gr**: Greece: C Macedonia, Nomos of Thessaloniki, Axios delta, 40°36'11.9"N, 22°42'46.7"E, 3 m,

climbing on *Rubus* spp. and on other woody species, 30 Jul 2016, *Giannakis* (TAUF); Axios delta, 40°37'19.2"N, 22°42'41.5"E, 5 m, on bank, climbing on *Rubus* spp. and on other woody species, 5 Aug 2016, *Giannakis* (TAUF); *ibid.*, 40°37'14.4"N, 22°42'41.8"E, 7 m, on bank, 27 Aug 2016, *Giannakis* (TAUF); Axios river, 40°37'14.9"N, 22°42'26.6"E, 10 m, climbing on *Rubus* spp. and on other woody species, 17 Sep 2016, *Giannakis* (TAUF); Axios river, 40°37'25.6"N, 22°42'39.3"E, 10 m, climbing on *Rubus* spp. and on other woody species, 11 Nov 2016, *Eleftheriadou & al.* (TAUF). – *Echinocystis lobata* is a xenophyte originating from the E part of North America (SE Canada to NE USA, see map in Slavík & Lhotská 1967: 259). It was introduced in the late 19th and early 20th centuries as an ornamental and medical plant to many countries in Europe and planted in several botanical gardens (DAISIE 2008). There has been increasing invasion during the last 20 years along the main rivers and floodplains from W to E Europe up to the Urals and beyond throughout Siberia (Cze-repanov 1995: 196; DAISIE 2008). Nowadays it is considered as fully naturalized in C and SE Europe (Slavík & Lhotská 1967; Tutin 1968; Vasić 2005; Lambdon & al. 2008+). In Bulgaria, it occurs only along the Danube River in the N part of the country (Petrova & al. 2013) and has not yet been recorded from Albania (Barina & al. 2014) or the Former Yugoslav Republic of Makedonija. There are no previous published records from Greece either (Arianoutsou & al. 2010; Dimopoulos & al. 2013). The cited populations in the Axios delta contain at least 1000 individuals. *Echinocystis lobata* is a fast-growing annual vine that can climb up to 12 m. It often completely covers woody plants of the area, such as *Alnus glutinosa* (L.) Gaertn., *Rubus* spp., *Salix* spp. and *Ulmus* spp.

Th. Giannakis, E. Eleftheriadou,
K. Theodoropoulos, S. Tsiftsis & R. Tsandekidis

Orchidaceae

Platanthera fornicata (Bab.) Buttler

+ **Gr:** Greece: Epirus, Nomos of Ioannina, Eparchia of Konitsa, S of Aetomilitsa, 40°14'32"N, 20°52'24"E, 800 m, open deciduous *Quercus* forest on sandstone and flysch with abundant *Cephalanthera rubra* (L.) Rich., *Epipactis atrorubens* (Hoffm.) Besser, *Ophrys scolopax* subsp. *cornuta* (Steven) E. G. Camus, *Orchis simia* Lam., *Orchis purpurea* Huds. and *Platanthera chlorantha* (Custer) Rchb., 21 May 1983, *Willing* 1237 (B as *P. bifolia*); *ibid.*, N

of Drosopiji, 40°13'31"N, 20°54'57"E, 740 m, extensive meadows on sandstone and flysch, 21 May 1983, *Willing* 1243 (B as *P. bifolia*); *ibid.*, S of Vourbani, 40°14'18"N, 20°47'32"E, 880 m, marly sandstone slope at entrance of village, 19 May 1983, *Willing* 1214 (B as *P. bifolia*); Thessaly, Nomos & Eparchia of Trikkala, 1.9 km SE of Elati, 39°29'02"N, 21°32'50"E, 780 m, grassy orchard, 29 May 1987, *Willing* 2583 (B as *P. bifolia*); W Makedonia, Nomos & Eparchia of Florina, 3.5 km SW of Drosopigi, 40°40'07"N, 21°26'20"E, 1080–1150 m, slope with meadows and *Fagus* forest on schist, 31 May 1984, *Willing* 1498a (B as *P. bifolia*); *ibid.*, Nomos & Eparchia of Kozani, Mt Titaros, SW of Grammeni Oxa, 40°10'30"N, 22°05'30"E, 1400–1600 m, mountain meadow with rivulets, 18 Jun 1989, *Willing* 6689 & 6690 (B as *P. bifolia*); *ibid.*, Nomos of Imathia, Eparchia of Naousa, 4.3 km NW of Naousa, 40°39'18"N, 22°01'40"E, 750 m, meadow in deciduous forest, 1 Jun 1985, *Willing* 1773 (B as *P. bifolia*); *ibid.*, W of Rodochori, 41°40'51"N, 21°59'52"E, 720 m, swamp meadow, 4 Jun 1984, *Willing* 1531 (B as *P. bifolia*); *ibid.*, C Makedonia, Nomos of Pella, Eparchia of Edessa, 4.7 km E of Ano Grammatiko, 40°42'23"N, 21°57'01"E, 1090 m, swamp meadows and pastures, 4 Jun 1984, *Willing* 1534 (B as *P. bifolia*); *ibid.*, 2 km SE of Ano Grammatiko, 40°41'08"N, 21°57'24"E, 1260 m, *Pteridium* heathland in *Fagus* forest, 5 Jun 1984, *Willing* 1543 (B as *P. bifolia*); *ibid.*, Eparchia of Almopia, 2.8 km NW of Pefkoto, 41°04'56"N, 22°04'13"E, 1110 m, *Fagus* forest, 26 May 1986, *Willing* 2054 (B as *P. bifolia*); *ibid.*, NNW of Livadhia, 41°01'39"N, 22°16'20"E, 1170 m, overgrown valley floodplain, 28 May 1986, *Willing* 2108 (B as *P. bifolia*); *ibid.*, Tzena Mts, S slopes of chain between peaks Tzena and Pinovo, above Notia, 41°07'50"N, 22°10'30"E, 950–1000 m, in narrow valley with beech wood on schist, 21 Jun 1992, *Greuter & al.* 23710 (B as *P. bifolia*); *ibid.*, Nomos of Kilikis, Eparchia of Peonia, Mt Paikon 3.3 km S of Livadhia, 40°58'30"N, 22°17'30"E, 1280 m, slope with *Juniperus* sp. and *Asphodeline lutea* (L.) Rchb., 10 Jun 1990, *Willing* 9859 (B as *P. bifolia*); *ibid.*, 2 km WNW of Livadhia, 41°00'N, 22°16'E, 1150 m, small wet gully on deforested slope, 11 Jun 1990, *Willing* 9984 (B as *P. bifolia*); *ibid.*, E Makedonia, Nomos & Eparchia of Drama: W Rhodopi Mts, 0.3 km S of Skaloti, 41°24'30"N, 24°16'30"E, 970 m, rocky slope with open low deciduous forest, 13 Jun 1990, *Willing* 10166 (B as *P. bifolia*); *ibid.*, 0.9 km S of Skaloti, 41°24'N, 24°16'E, 950 m, slope in *Pinus* for-

est, 17 Jun 1992, *Willing 18895* (B as *P. bifolia*); *ibid.*, 3.3 km S of Kallikarpo, 41°23'N, 24°13'E, 600 m, meadow in *Quercus* scrub, 22 Jun 1992, *Willing 19793* (B as *P. bifolia*). – Buttler (2012a, 2012b) recently reappraised and corroborated the taxonomic findings of Babington (1836) and Müller (1868) that the name *Platanthera bifolia* (s.l.) designates at least two taxa, *P. bifolia* (L.) Rich. s.str. and *P. fornicata*, which deserve specific rank based on combined differences in morphology, chorology, ecology and flowering time. The former seems to be confined to atlantic-subatlantic parts of W and C Europe including a disjunct distribution area in the Alps, whereas the latter is widespread in Europe extending eastwards to temperate C Siberia. The ecological and phenological discontinuities largely exclude hybridization in areas of sympatric occurrence. The revision by K. P. Buttler of recent collections of this group from Greece kept at B revealed that the majority of specimens (i.e. 17 out of 18) belong to *P. fornicata*. This does not automatically mean the exclusion of *P. bifolia* s.str. from the flora of Greece. There is one single specimen in B (viz. Greece (NC), C Makedonia, Nomos of Kilkis, Eparchia of Peonia, Mt Paikon 2.2 km WNW of Livadhi, 41°00'30"N, 22°16'00"E, 1180 m, rivulet with scattered wet meadow patches in dry grassland, 28 Jun 1989, *Willing 7441*) that exhibits the floral dimensions of *P. bifolia* s.str. but deviates in habit from NW European specimens of that taxon examined by Buttler (2011a). Whether such populations represent another, as yet undescribed taxon within the *P. bifolia* complex is so far unknown and should be subject to further taxonomic research. Similar forms that, for the time being, were included with doubt in *P. bifolia* s.str. by Buttler (2012a, 2012b) have been reported from the Carpathian Mts in Romania (Schur 1866: 646, as *P. bifolia* var. *obtusifolia* Schur). The localities of the revised material refer to the floristic regions of Southern Pindos (SPi), Northern Pindos (NPi), North Central (NC) and North East (NE) as circumscribed in Dimopoulos & al. (2013: 29, fig. 2). As a consequence, published records of *P. bifolia* (s.l.) from other parts of Greece, viz. the Ionian islands (Boissier 1882: 82–83; Kapteyn den Boumeester & Willing 1988), Sterea Ellas (Willing & Willing 1983: 410) and East Central (Biel & Rudolph 1992: 48) are in need of revision by re-examination (or collection, respectively) of relevant voucher material. To facilitate such efforts, a determination key for species of the *P. bifolia* aggregate (translated from Buttler 2012a: 96) is given here:

1. Flowers small: spur 12–20(–23) mm long, labellum 6–10.5(–12) mm long, anthers <1 mm apart, parallel or downwards slightly connivent; pollinia on very short, <1 mm-long stalks. Plants relatively low-growing, 10–25(–35) cm tall, inflorescence usually dense, in lateral view without gaps between flowers. Flowering time 2–3 weeks earlier than in *P. fornicata* in areas of sympatric occurrence *Platanthera bifolia*
- Flowers large: spur (18–)25–41 mm long, labellum (9.5–)11–16 mm long, anthers 1–1.5 mm apart, parallel; pollinia on distinct, >1 mm-long stalks. Plants relatively high-growing, (25–)30–90 cm tall, inflorescence usually lax, in lateral view with pronounced gaps between flowers. In overlapping range of floral characters, large-flowered and low-growing or small-flowered and high-growing plants belong here *Platanthera fornicata*

Within *Platanthera fornicata*, a putative geographical differentiation of a (sub-)meridional subspecies (*P. fornicata* subsp. *atropatanica* (B. Baumann & al.) Buttler based on *P. bifolia* subsp. *atropatanica* B. Baumann & al. described from Azerbaijan; Baumann & al. 2003) and a temperate-boreal subspecies (*P. fornicata* subsp. *fornicata*) needs further, corroborating taxonomic investigation of additional, more comprehensive material from Europe and SW Asia (Buttler 2012a: 96, 103).

K. P. Buttler & Th. Raus

Orobanchaceae

Orobanche grenieri F. W. Schultz – Fig. 1

+ **Ar:** Armenia: Ararat Marz, near road from Garni to Nubarashen, little pass c. 1 km N of Azat Reservoir, 40°05'24"N, 48°37'10"E, c. 1200 m, semi-desert, moving clay, open soil with very sparse vegetation cover, parasitic on *Lactuca takhtadzhianii* Sosn. (root attachment verified, matrix nova), 10 Jul 2013, *Gabrielian & al. 2013-24* (B 10 0576237, ERE). – After the recent findings of *Orobanche grenieri* in Azerbaijan (Rätzel & al. 2015) and Georgia (Piwowarczyk & al. 2015), this is the first confirmed record of this species for the flora of Armenia. Former records of the closely related *O. cernua* Loebl., well-known from several regions of Armenia (Tsaturyan & Grigoryan 1987), may also belong here.

S. Rätzel, P. Hein &
E. von Raab-Straube

In the last few years, plant scientists, in following Schneeweiss & al. (2004a, 2004b), have transferred the species of *Orobanche* L. previously referred to *O. sect. Trionychon* Wallr. to the genus *Phelipanche* Pomel. Also, during the 10th World Congress on Parasitic Plants




Fig. 1. A & B: *Orobanche grenieri* – Armenia: Ararat province, between Garni and Nubarashen, pass N of Azat Reservoir, 10 Jul 2013, photograph by E. von Raab-Straube.

held in Kuşadası, Turkey, in June 2009, it was agreed to adopt this taxonomic separation into two distinct genera (Joel 2009). The main diagnostic differences between *Orobanche* and *Phelipanche* (Teryokhin 1997) are: flowers with 2 adnate bracteoles at base of calyx in *Phelipanche*, versus bracteoles missing in *Orobanche*; ripe capsule opening apically with style usually shedding in *Phelipanche*, versus opening laterally with persistent style in *Orobanche*; flowering stem usually branched in *Phelipanche*, versus usually unbranched in *Orobanche*; basic chromosome number $x = 12$ and chromosomes medium-sized, submetacentric to acrocentric in *Phelipanche*, versus $x = 19$ and chromosomes small, mostly metacentric to submetacentric in *Orobanche* (Schneeweiss & al. 2004b).

Several studies suggest deviating relationships between the taxa belonging to *Orobanche* sect. *Orobanche* and *O.* sect. *Trionychon* (Román & al. 2003; Manen & al. 2004; Park & al. 2007). Up to now, the genus *Orobanche* has been considered in a wide sense (Domina 2009;

Domina & Raimondo 2009; Domina & al. 2013; Domina & Danin 2014), including the two traditional sections of *Orobanche* as circumscribed by Beck von Mannagetta (1930). However, providing the taxonomic practice accepted in the Euro+Med PlantBase (Euro+Med 2006+) and the up-to-date and critically evaluated consensus on the taxonomy of the species concerned, it is preferable to follow the taxonomic delimitation that has become current in this group. This treatment has the practical convenience of distinguishing morphologically well-characterized taxa at the generic level. The four new combinations proposed below result from this choice.

G. Domina

***Phelipanche ballii* (Maire) Domina, *comb. nov.* $\equiv$ *Orobanche purpurea* var. *ballii* Maire in Mém. Soc. Sci. Nat. Maroc 21–22: 12. 1930 $\equiv$ *Phelipanche purpurea* subsp. *ballii* (Maire) Carlón & al. in Doc. Jard. Bot. Atlántico 6: 104. 2008 $\equiv$ *Orobanche ballii* (Maire) Domina in Pl. Biosyst. 147: 763. 2013.**

Phelipanche cohenii (Domina & Danin) Domina, **comb. nov.** $\equiv$ *Orobanche cohenii* Domina & Danin in Fl. Medit. 24: 63. 2014.

Phelipanche daninii (Domina & Raimondo) Domina, **comb. nov.** $\equiv$ *Orobanche daninii* Domina & Raimondo in Fl. Medit. 19: 185. 2009.

Phelipanche tricholoba (Reut.) Domina, **comb. nov.** $\equiv$ *Phelypaea tricholoba* Reut. in Candolle, Prodr. 11: 10. 1847 $\equiv$ *Orobanche tricholoba* (Reut.) Domina in Willdenowia 39: 331. 2010.

Solanaceae

Datura wrightii Regel

A **Tn**: Tunisia: Gouv. Bizerte, Cap Serrat, E slope and around houses on beach, 37°13'07–19"N, 09°13'22–26"E, 0–50 m, ruderal, perennial, old fruits, 30 Mar 2014, *OPTIMA Iter Mediterraneum XII, Tunisia, 2014 0930* (PAL-GR 061982 as *D. metel* L.). – The plant was originally misidentified as *Datura metel* L. (Greuter & Domina 2015), which is easily recognizable by the capsules covered with blunt and short tubercles. *Datura wrightii*, provided with spiny capsules and stems covered with short eglandular indumentum, has recently been discovered for the first time in N Africa in Morocco (Verloove in Sukhorukov & al. 2016). This type of indumentum, along with size of stigmas and corollas, enables its distinction from the similar *D. inoxia* Mill. (the discussed specimen is available at: http://147.163.105.223/zoomify/view_img.asp?ic=61982_GR). N. M. G. Ardenghi

Acknowledgements

P. Hein and E. v. Raab-Straube cordially thank Mariam Aghababayan and Eleonora Gabrielian for their invaluable help in organizing a common field trip to Armenia in 2013. Financial support from the Verein der Freunde des Botanischen Gartens und Botanischen Museums Berlin-Dahlem e. V. is gratefully acknowledged. The field trip was carried out in the framework of the project “Developing tools for conserving the plant diversity of the Transcaucasus” funded by VolkswagenStiftung (reference AZ85021).

References

Arianoutsou M., Bazos I., Delipetrou P., Kokkoris Y. 2010: The alien flora of Greece: taxonomy, life traits and habitat preferences. – Biol. Invasions **12**: 3525–3549.
Aypeisova S. A. 2012: Konspekt floryi Aktyubinskogo floristicheskogo okruga. – Aktobe: Aktyubinskij Go-

sudarstvennyj Universitet Im. K. Zhubanova, Ministerstvo obrazovaniya i nauki respubliki Kazakhstan.
Babington C. C. 1836: On several new or imperfectly understood British and European plants. – Transact. Linn. Soc. London **17**: 451–464.
Barina Z., Rakaj M., Somogyi G., Erös-Honti Z. & Pifko D. 2014: The alien flora of Albania: history, current status and future trends. – Weed Res. **54**: 196–215.
Baumann B., Baumann H., Lorenz R. M. & Peter R. 2003: Beiträge zur Orchideenflora des östlichen Transkaukasus und Talysch (Aserbaidshan). – J. Eur. Orch. **35**: 163–231.
Beck von Mannagetta G. 1930: *Orobanchaceae*. – In: Engler A. (ed.), Das Pflanzenreich **96**. – Leipzig: Engelmann.
Biel B. & Rudolph F. 1992: Orchideenkartierung auf der Pelion-Halbinsel, Griechenland. – Ber. Arbeitskreis. Heimische Orchid. **9**: 31–63.
Boissier E. 1882–1884: Flora orientalis **5** [pp. 1–428 (1882), pp. 429–868 (1884)]. – Genève, Bâle & Lyon: H. Georg.
Buttler K. P. 2012a [“2011”]: Revision von *Platanthera bifolia* sensu lato. Taxonomisch-nomenklatorische Neubewertung des Formenkreises um die Weiße Waldhyazinthe. – Jahresber. Wetterauischen Ges. Gesamte Naturk. **159–161** (Festschrift): 93–108.
Buttler K. P. 2012b: Revision von *Platanthera bifolia* sensu lato. Taxonomisch-nomenklatorische Neubewertung des Formenkreises um die Weiße Waldhyazinthe. Ergänzungen. – Published at: http://www.wetterauischegesellschaft.de/images/veroeffentlichungen/buttler_2011_revision_von_platanthera_bifolia_sensu_lato.pdf [accessed 24 Feb 2017].
Chen Y. & Brouillet L. 2011: *Erigeron* L. – Pp. 634–650 in: Wu Z. Y., Raven P. H. & Hong D. Y. (ed.), Flora of China **20–21**. – Beijing: Science Press; St. Louis: Missouri Botanical Garden Press.
Chronopoulos G. & Christodoulakis D. 1996: Contribution to the urban ecology of Greece: The flora of the city of Patras and the surrounding area. – Bot. Helv. **106**: 159–176.
Czerepanov S. K. 1995: Vascular plants of Russia and adjacent states (the former USSR). – Cambridge: Cambridge University Press.
DAISIE [Delivering Alien Invasive Species Inventories for Europe] 2008+: Species factsheet: *Echinocystis lobata*. – Published at: http://www.europe-aliens.org/pdf/Echinocystis_lobata.pdf [accessed 15 Jan 2017].
Davidenko O. N., Grebenyuk S. I. & Nevsky S. A. 2016: Sbory redkikh vidov rastenij v povolzh'e. Collecting rare plant species in the Volga region. – Nauchnaya Zhizn' **2016(2)**: 211–219.
Dimopoulos P., Raus Th., Bergmeier E., Constantinidis Th., Iatrou G., Kokkini S., Strid A. & Tsanoudakis D. 2013: Vascular plants of Greece: An annotated checklist. – Berlin: Botanic Garden and Botanical Museum

- Berlin-Dahlem; Athens: Hellenic Botanical Society. – Englera **31**.
- Domina G. 2009: *Orobanchaceae*. – P. 331 in: Greuter W. & Raab-Straube E. von (ed.), Euro+Med Notulae, 4. – Willdenowia **39**: 327–333.
- Domina G. & Danin A. 2014: *Orobanche cohenii* (*Orobanchaceae*) a new species from Israel. – Fl. Medit. **24**: 63–69.
- Domina G., Greuter W., Marino P. & Schäfer P. A. 2013: Types of names of *Orobanche* taxa described from North Africa. – Pl. Biosyst. **147**: 758–766.
- Domina G. & Raimondo F. M. 2009: A new species of *Orobanche* (*Orobanchaceae*) from Israel. – Fl. Medit. **19**: 185–188.
- Dorofeyev V. I. 2002: Krestocvetnye (*Cruciferae* Juss.) Evropejskoj Rossii. – Turczaninowia **5**(3): 5–114.
- Euro+Med 2006+ [continuously updated]: Euro+Med PlantBase – the information resource for Euro-Mediterranean plant diversity. – Published at <http://ww2.bgbm.org/EuroPlusMed> [accessed 21 Feb 2017].
- Galanos C. J. 2015: The alien flora of terrestrial and marine ecosystems of Rodos island (SE Aegean, Greece). – Willdenowia **45**: 261–278.
- Garnweidner E. 1989: Florenliste der Exkursion der Bayerischen Botanischen Gesellschaft nach Kreta. – Ber. Bayer. Bot. Ges. **60**: 157–168.
- Greuter W. 2006+ [continuously updated]: *Erigeron sumatrensis* Retz. – Euro+Med PlantBase – the information resource for Euro-Mediterranean plant diversity. – Published at: http://euromed.luomus.fi/euromed_map.php?taxon=283047&size=medium [accessed 13 Dec 2016].
- Greuter W. & Domina G. 2015: 151. Checklist of the vascular plants collected during the 12th “Iter Mediterraneum” in Tunisia, 24 March – 4 April 2014. – Bocconea **27**: 21–61.
- Greuter W. & Raab-Straube E. von (ed.) 2005: Euro+Med Notulae, 1. – Willdenowia **35**: 223–239.
- Hügin G. 1987: Einige Bemerkungen zu wenig bekannten *Amaranthus*- Sippen (*Amaranthaceae*) Mitteleuropas. – Willdenowia **16**: 453–478.
- Iamónico D. 2015: *Amaranthaceae*. – In: Euro+Med PlantBase – the information resource for Euro-Mediterranean plant diversity. – Published at <http://ww2.bgbm.org/EuroPlusMed/PTaxonDetail.asp?NameCache=Amaranthaceae&PTReffk=7300000> [accessed 24 Feb 2017].
- Joel D. M. 2009: The new nomenclature of *Orobanche* and *Phelipanche*. – Weed Res. **49**: 6–7.
- Kapteyn den Boumeester D. & Willing E. 1988: Aktuelle Verbreitung der Orchideen auf Kerkira (Korfu/Griechenland). – Ber. Arbeitskreis. Heimische Orchid., Beiheft **2**: 4–128.
- Kleinsteuber A., Ristow M. & Hassler M. (ed.) 2016: Flora von Rhodos und Chalki. Band 1: Allgemeiner Teil. Spezieller Teil: *Polypodiopsida*, *Equisetopsida* und *Lycopodiopsida*. *Pinopsida* und *Gnetopsida*. *Magnoliopsida* (Familien A–F). – Karlsruhe: Naturwissenschaftlicher Verlag A. Kleinsteuber.
- Konechnaya G. Yu. 1994: Krestovnik – *Senecio* L. – Pp. 52–63 in: Tzvelev N. N. (ed.), Flora Evropejskoj Chasti SSSR **7**. – Sankt-Peterburg: „Nauka“.
- Kotov M. I. 1979: Tausheriya – *Tauscheria* Fisch. ex DC. – P. 72 in: Fedorov A. A. (ed.), Flora Evropejskoj Chasti SSSR **4**. – Leningrad: “Nauka”, Leningradskoe Otdelenie.
- Lambdon P. W., Pyšek P., Basnou C., Hejda M., Ariannoutsou M., Essl F., Jarošík V., Pergl J., Winter M., Anastasiu P., Andriopoulos P., Bazos I., Brundu G., Celesti-Grappo L., Chassot P., Delipetrou P., Josefsson M., Kark S., Klotz S., Kokkoris Y., Kühn I., Marchante H., Perglová I., Pino J., Vilà M., Zikos A., Roy D. & Hulme P. E. 2008: Alien flora of Europe: species diversity, temporal trends, geographical patterns and research needs. – Preslia **80**: 101–149.
- Le Floch E., Boulos L. & Vela E. 2010: Catalogue synonymique commenté de la Flore de Tunisie. – Tunis: Ministère de l’Environnement et du Développement durable.
- Manen J. F., Habashi C., Jeanmonod D., Park J. M. & Schneeweiss G. M. 2004: Phylogeny and intraspecific variability of holoparasitic *Orobanche* (*Orobanchaceae*) inferred from plastid *rbcL* sequences. – Molec. Phylogen. Evol. **33**: 482–500.
- Müller H. 1868: Beobachtungen an westfälischen Orchideen. – Verh. Naturhist. Vereines Preuss. Rheinl. Westphalens **25**: 1–62 & Tab. I–II.
- Park J. M., Manen J. F. & Schneeweiss G. M. 2007: Horizontal gene transfer of a plastid gene in the non-photosynthetic flowering plants *Orobanche* and *Phelipanche* (*Orobanchaceae*). – Molec. Phylogen. Evol. **43**: 974–985.
- Petrova A., Vladimirov V. & Georgiev V. 2013: Invasive alien species of vascular plants in Bulgaria. – Sofia: Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences.
- Piowarczyk R., Kwolek D., Denysenko M., Cygan M., Góralski G., Ślesak H., Tuleja M. & Joachimiak A. J. 2015: *Orobanche grenieri* (*Orobanchaceae*), a southwestern European species newly found in Asia. – Ann. Bot. Fenn. **52**: 411–418.
- Pyšek P., Richardson D. M., Rejmanek M., Webster G. L., Williamson M. & Kirschner J. 2002: Alien plants in checklists and floras: towards better communication between taxonomists and ecologists. – Taxon **53**: 131–143.
- Raab-Straube E. von & Raus Th. (ed.) 2016: Euro+Med-Checklist Notulae, 6 [Notulae ad floram euro-mediterraneam pertinentes 35]. – Willdenowia **46**: 423–442.
- Rätzel S., Ristow M. & Uhlich H. 2015: *Orobanche grenieri* F. W. Schultz. – Pp. 456–457 in: Raab-Straube E. von & Raus Th. (ed.), Euro+Med-Checklist Notulae, 5 [Notulae ad floram euro-mediterraneam pertinentes 34]. – Willdenowia **45**: 449–464.

- Raus Th. 1997: *Amaranthus* L. – Pp. 138–146 in: Strid A. & Tan K. (ed.), *Flora hellenica* **1**. – Königstein: Koeltz Scientific Books.
- Román B., Alfaro C., Torres A., Satovic Z., Pujadas A. & Rubiales D. 2003: Genetic relationships among *Orobanch* species as revealed by RAPD analysis. – *Ann. Bot.* **91**: 637–642.
- Ryabinina Z. N. & Knyazev M. S. 2009: The identification manual of vascular plants of the Orenburg region. – Moscow: Scientific Press Ltd KMK.
- SANBI 2012: *Erigeron sumatrensis* Retz. – In: African Plant Database (version 3.4.0). Conservatoire et Jardin botaniques de la Ville de Genève and South African National Biodiversity Institute, Pretoria. – Published at: <http://www.ville-ge.ch/musinfo/bd/cjb/africa/details.php?langue=en&id=101799> [accessed 13 Dec 2016].
- Schneeweiss G. M., Colwell A. E., Park J. M., Jang C. & Stuessy T. 2004a: Phylogeny of holoparasitic *Orobanch* (*Orobanchaceae*) inferred from nuclear ITS sequences. – *Molec. Phylogen. Evol.* **30**: 465–478.
- Schneeweiss G. M., Palomeque T., Colwell A. E. & Weiss-Schneeweiss H. 2004b: Chromosome numbers and karyotype evolution in holoparasitic *Orobanch* (*Orobanchaceae*) and related genera. – *Amer. J. Bot.* **91**: 439–448.
- Schur J. F. 1866: *Enumeratio plantarum Transsilvaniae exhibens: stirpes phanerogamas sponte crescentes atque frequentius cultas, cryptogamas vasculares, characeas, etiam muscos hepaticasque*. – Vindobonae: G. Braumüller.
- Slavík B. & Lhotská M. 1967: Chorologie und Verbreitungsbiologie von *Echinocystis lobata* (Michx.) Torr. et Gray mit besonderer Berücksichtigung ihres Vorkommens in der Tschechoslowakei. – *Folia Geobot. Phytotax.* **2**: 255–282.
- Sukhorukov A. P., Martín-Bravo S., Verloove F., Maroyi A., Iamónico D., Catarino L., El Mokni R., Daniel T. F., Belyaeva I. V. & Kushunina M. 2016: Chorological and taxonomic notes on African plants. – *Bot. Lett.* **163**: 417–428.
- Teryokhin E. S. 1997: *Weed Broomrapes: Systematics, Ontogenesis, Biology, Evolution*. – Landshut: Aufstiegs-Verlag.
- Tsaturyan T. G. & Grigoryan S. B. 1987: *Orobanchaceae*. – Pp. 361–384 in: Takhtajan A. L. (ed.), *Flora Armenii* **8. Verbenaceae – Lentibulariaceae**. – Yerevan: Izdatel'stvo Akademii Nauk Armjanskoy SSR.
- Vasić O. 2005: *Echinocystis lobata* (Michx.) Torrey et Gray in Serbia. – *Acta Bot. Croat.* **64**: 369–373.
- Willing B. & Willing E. 1983: Beitrag zu Verbreitung der Orchideen Ätoliens und Akarnaniens sowie der Insel Lefkas (NW-Griechenland). – *Mitteilungsbl. Arbeitskreis Heimische Orchid. Baden-Württemberg* **15**: 351–413.

Willdenowia

Open-access online edition www.bioone.org/loi/will  BioOne

Online ISSN 1868-6397 · Print ISSN 0511-9618 · Impact factor 0.500

Published by the Botanic Garden and Botanical Museum Berlin, Freie Universität Berlin

© 2017 The Authors · This open-access article is distributed under the CC BY 4.0 licence