

Development and Characterization of Microsatellite Loci For the Endangered Scrub Lupine, *Lupinus aridorum* (Fabaceae)

Authors: Ricono, Angela, Bupp, Glen, Peterson, Cheryl, Nunziata, Schyler O., Lance, Stacey L., et al.

Source: Applications in Plant Sciences, 3(4)

Published By: Botanical Society of America

URL: <https://doi.org/10.3732/apps.1500013>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

PRIMER NOTE

DEVELOPMENT AND CHARACTERIZATION OF MICROSATELLITE LOCI FOR THE ENDANGERED SCRUB LUPINE, *LUPINUS ARIDORUM* (FABACEAE)¹

ANGELA RICONO², GLEN BUPP³, CHERYL PETERSON³, SCHYLER O. NUNZIATA⁴, STACEY L. LANCE⁴,
AND CHRISTIN L. PRUETT^{2,5}

²Department of Biological Sciences, Florida Institute of Technology, Melbourne, Florida 32901 USA; ³Rare Plant Conservation Program, Bok Tower Gardens, Lake Wales, Florida 33853 USA; and ⁴Savannah River Ecology Laboratory, University of Georgia, Aiken, South Carolina 29802 USA

- *Premise of the study:* Microsatellite primers were developed in scrub lupine (*Lupinus aridorum*, Fabaceae), an endemic species to Florida that is listed as endangered in the United States, to assess connectivity among populations, identify hybrids, and examine genetic diversity.
- *Methods and Results:* We isolated and characterized 12 microsatellite loci polymorphic in scrub lupine or in closely related species (i.e., sky-blue lupine [*L. diffusus*] and Gulf Coast lupine [*L. westianus*]). Loci showed low to moderate polymorphism, ranging from two to 14 alleles per locus and 0.01 to 0.86 observed heterozygosity.
- *Conclusions:* These loci are the first developed for Florida species of lupine and will be used to determine differentiation among species and to aid in conservation of the endangered scrub lupine.

Key words: Fabaceae; Florida; *Lupinus*; microsatellite; PCR primers.

Scrub lupine (*Lupinus aridorum* McFarlin ex Beckner) is an endangered plant restricted to the Winter Haven and Mount Dora ridge systems in central Florida that has declined primarily due to habitat loss (USFWS, 1987). Polymorphic genetic markers are needed to answer questions about genetic diversity and connectivity among *L. aridorum* populations, genetic relatedness among Florida lupine species including the closely related Gulf Coast lupine (*L. westianus* Small), and hybridization between sympatric populations of *L. aridorum* and sky-blue lupine (*L. diffusus* Nutt.; Bupp, 2013).

METHODS AND RESULTS

Total genomic DNA was extracted from leaf samples of two individuals from *L. aridorum* populations using a QIAamp DNA Mini Kit (QIAGEN, Valencia, California, USA). Using a Covaris S220 (Woburn, Massachusetts, USA), we prepared an Illumina paired-end shotgun library by shearing 1 µg of DNA as described in the Illumina TruSeq DNA Library Kit (Illumina, San Diego, California, USA) and using a multiplex identifier adapter index. Illumina sequencing, with 100-bp paired-end reads, was conducted on a HiSeq 2000 (Illumina). We used the program PAL_FINDER_v0.02.03 (Castoe et al., 2012) to examine five million reads and extracted the reads that contained di-, tri-,

tetra-, penta-, and hexanucleotide microsatellites. Positive reads were batched to the program Primer3 (version 2.0.0; Rozen and Skaletsky, 1999) for primer design. We selected loci for primer sequences that only occurred one time in the five million reads to avoid problems with copy number of the sequence in the genome. Ninety loci of the 1740 loci that met this criterion were chosen. One primer from each pair was modified on the 5' end with an engineered sequence (CAG tag 5'-CAGTCGGGCGTCATCA-3') to enable use of a third primer in the PCR that was fluorescently labeled with one of three dyes (6-FAM, NED, or VIC; Applied Biosystems, Culver City, California, USA).

Primer pairs were tested for amplification and polymorphism using DNA obtained from four individuals. Amplifications were in 20-µL volumes (250 µg/mL bovine serum albumin [BSA], 2 µL 10× Buffer B, 25 mM MgCl₂, 5 µM unlabeled primer, 0.5 µM tag-labeled primer, 5 µM universal dye-labeled primer, 2.5 mM dNTPs, 0.5 units *Taq* DNA polymerase [Fisher Scientific, Pittsburgh, Pennsylvania, USA], and 20 ng DNA template) using a Bio-Rad MyCycler (Hercules, California, USA) thermal cycler. We used touchdown cycling conditions to amplify DNA and to attach the universal dye-labeled primer. Parameters consisted of an initial denaturation step of 2 min 30 s at 95°C; followed by 20 cycles of 95°C for 20 s, 65–50°C annealing temperature for 20 s (decreasing 0.5°C per cycle), and extension step of 72°C for 30 s; followed by 15 cycles of 95°C for 20 s, 55–45°C for 20 s, and 72°C for 30 s. Cycles were followed with a final extension step of 72°C for 10 min. Amplifications were run on an ABI3730XL sequencer (Applied Biosystems).

Twelve of the tested primer pairs amplified high-quality PCR product that exhibited polymorphism in *L. aridorum*, *L. diffusus*, or *L. westianus* (Table 1). We then assessed the variability at these loci using 19–22 individuals of *L. aridorum*, 9–22 *L. diffusus*, and 12–20 *L. westianus* (Table 2). Alleles were scored using GeneMapper software (Applied Biosystems). We evaluated the number of alleles per locus, observed heterozygosity, and expected heterozygosity and tested for Hardy–Weinberg equilibrium (HWE) and linkage equilibrium using Arlequin version 3.5 (Excoffier et al., 2005). Sequences of raw paired-end reads are available in the SRA database of the National Center for Biotechnology Information (bioproject no. PRJNA274660) and as Appendix S1. Vouchers of leaves collected for this study were deposited at Bok Tower Gardens, Lake Wales, Florida (Table 2).

We found that the number of alleles per locus ranged from one to seven for *L. aridorum*, one to nine for *L. westianus*, and one to 14 for *L. diffusus*. For

¹Manuscript received 11 February 2015; revision accepted 5 March 2015.

This work was supported by the Florida Institute of Technology, Bok Tower Gardens, U.S. Fish and Wildlife Service, Florida Forest Service, Florida Department of Agriculture and Consumer Services, and by the U.S. Department of Energy (award no. DE-FC09-07SR22506 to the University of Georgia Research Foundation).

⁵Author for correspondence: cpruett@fit.edu

doi:10.3732/apps.1500013

TABLE 1. Characteristics of 12 microsatellite loci in *Lupinus aridorum*, *L. westianus*, and *L. diffusus*.

Locus	Primer sequences (5'–3')	Repeat motif	Allele size range (bp) ^a	T _a (°C)
Luar3	F: *AGAATATAAAGGTTTACAAGGGC R: AAGCAGTTGTTGACTACACAGATACG	(AAAG) ₉	212–226	61/51
Luar11	F: *GGATTAGATTGCTACTATTGGCCC R: CAGTCCGACCAGATAGTTTAACCG	(AACT) ₈	142–193	54/44
Luar12	F: *TGGCAGGGAAGGGAAGTAGG R: CAGTCGCATGAGAGCGGG	(ATC) ₂₀	242–248	54/44
Luar15	F: *GCACATGGCTTAACCAACTTCC R: CTAAGCGTGCGCATGTGG	(ATAGG) ₁₁	151–161	65/55
Luar31	F: *TTTCATACCCCTGCGTTTCC R: GTGCTGGAACAGGTAGTGGC	(ATAGG) ₁₂	259–269	54/44
Luar48	F: *GCGTCGTTTGACATTGACG R: GGGAATTGAGAATAAAGAGGG	(AAAG) ₉	140–166	54/44
Luar69	F: *TCACCATTATCCACATTACC R: GTGGCCCATTCAGTTCC	(AAC) ₁₅	236–242	65/55
Luar71	F: *CCGTATCCATATCCACTTTCCC R: GTTTAGTTTCATTGTGCAACCCGC	(TTC) ₁₂	318–324	55/45
Luar73	F: *TGACAGCTAGAGGTTTCAAGGC R: GTTTAGCTTCTCTTCCACGCAAGC	(TTC) ₁₁	147–159	55/45
Luar74	F: *CTTCTCACCTCATTTCGAATTC R: GTTTGGGAACCCATTATTCGAGG	(TTC) ₁₃	238–250	55/45
Luar84	F: *TTCTATCTTATCCTATCTTATTGTGTCC R: GTTTCAGGATATGTTCCGACGGG	(ATAGG) ₉	118–138	60/50
Luar89	F: *ATGTATGAACAACACACGGG R: GTTTAGTTGATCAAATGGCGGAGG	(AACCT) ₁₀	408–418	60/50

Note: T_a = annealing temperature for touchdown protocol.

^aRange of alleles in base pairs including CAG tag (location indicated with asterisk on primer sequence).

polymorphic loci, observed heterozygosity for *L. aridorum* ranged from 0.01 to 0.68, 0.15 to 0.86 for *L. westianus*, and 0.01 to 0.78 for *L. diffusus*. Expected heterozygosity for *L. aridorum* was between 0.08 and 0.73, 0.19 and 0.83 for *L. westianus*, and 0.09 and 0.90 for *L. diffusus*. All loci were in linkage equilibrium but several loci were out of HWE after correction for multiple tests (Table 2). Deviations from HWE are expected because samples were collected from several

areas and individuals might be able to self-fertilize (Bupp, 2013; Bupp, personal communication).

CONCLUSIONS

These loci are the first to be developed for Florida lupine species and are polymorphic in at least three of the five species of North American unifoliate lupines (Dunn, 1971), *L. aridorum*, *L. diffusus*, and *L. westianus*. Although microsatellite primers have been developed for other, distantly related species of lupines (e.g., Gonzalez et al., 2010), cross-species amplification of *L. aridorum* with these primers was unlikely because North American unifoliate lupines represent a genetically divergent lineage that is distinctly different from other lupines (Mahé et al., 2011).

Lupinus aridorum is a critically endangered species, and a clear understanding of the effects of habitat fragmentation on the genetic diversity of populations and connectivity among populations is needed to inform conservation efforts. In addition, many of these loci are polymorphic in *L. westianus*, a threatened species in the state of Florida (Wunderlin and Hansen, 2008). Conservation genetic studies will aid in management of these rare lupines.

LITERATURE CITED

- BUPP, G. 2013. Cytogenetic and population genetic analysis of the endangered scrub lupine (*Lupinus aridorum*). Master's Thesis, Florida Institute of Technology, Melbourne, Florida, USA.
- CASTOE, T. A., A. W. POOLE, A. P. J. DE KONING, K. L. JONES, D. F. TOMBACK, S. J. OYLER-McCANCE, J. A. FIKE, ET AL. 2012. Rapid microsatellite identification from Illumina paired-end genomic sequencing in two birds and a snake. *PLoS ONE* 7: e30953.
- DUNN, D. B. 1971. A case of long range dispersal and "rapid speciation" in *Lupinus*. *Transactions of the Missouri Academy of Science* 5: 26–38.

TABLE 2. Genetic properties of 12 microsatellites developed for *Lupinus aridorum* with cross-species amplification in *L. westianus* and *L. diffusus*.^a

Locus	<i>L. aridorum</i> (n = 19–22)			<i>L. westianus</i> (n = 10–20)			<i>L. diffusus</i> (n = 9–22)		
	A	H _o	H _e ^b	A	H _o	H _e ^b	A	H _o	H _e ^b
Luar3	2	0.08	0.22	2	0.16	0.33	5	0.14	0.25
Luar11	1	—	—	5	0.15	0.24	1	—	—
Luar12	2	0.38	0.31	3	0.20	0.19	2	0.04	0.12
Luar15	5	0.68	0.51	1	—	—	1	—	—
Luar31	2	0.01	0.08	1	—	—	2	0.01	0.09
Luar48	4	0.08	0.26*	6	0.52	0.68	14	0.70	0.90
Luar69	5	0.55	0.73	7	0.41	0.83*	8	0.62	0.87
Luar71	4	0.40	0.55	5	0.68	0.69	8	0.38	0.83*
Luar73	7	0.60	0.66	9	0.86	0.82	8	0.68	0.89
Luar74	6	0.52	0.70	5	0.68	0.67	6	0.78	0.85
Luar84	3	0.40	0.63	5	0.76	0.68	7	0.50	0.72
Luar89	1	—	—	1	—	—	3	0.20	0.65*

Note: A = number of alleles sampled; H_e = expected heterozygosity; H_o = observed heterozygosity; n = number of individuals sampled.

^aGeographic locations for samples are: *Lupinus aridorum* (Orange County, Pines of Wekiva); *L. westianus* (Walton County, Point Washington State Forest); and *L. diffusus* (Brevard County: 28.0176°N, 80.6000°W; 28.7711°N, 80.7829°W; 28.5054°N, 80.6600°W; Hillsborough County: 27.5953°N, 82.2366°W; Polk County: 27.9385°N, 81.5738°W). All samples are from Florida, and vouchers are deposited at Bok Tower Gardens, Lake Wales, Florida (accession no. La61069s).

^bLoci that are out of Hardy–Weinberg equilibrium are indicated with an asterisk after Bonferroni correction for multiple tests ($P < 0.004$).

- EXCOFFIER, L., G. LAVAL, AND S. SCHNEIDER. 2005. Arlequin ver. 3.0: An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1: 47–50.
- GONZALEZ, P. B. P., S. C. K. STRAUB, J. J. DOYLE, P. E. M. ORTEGA, H. E. S. GARRIDO, AND I. J. M. BUTLER. 2010. Development of microsatellite markers in *Lupinus luteus* (Fabaceae) and cross-species amplification in other lupine species. *American Journal of Botany* 97: e72–e74.
- MAHÉ, F., D. MARKOVA, R. PASQUET, M.-T. MISSET, AND A. AÏNOUCHE. 2011. Isolation, phylogeny and evolution of the *SmyRK* gene in the legume genus *Lupinus* L. *Molecular Phylogenetics and Evolution* 60: 49–61.
- ROZEN, S., AND H. SKALETISKY. 1999. Primer3 on the WWW for general users and for biologist programmers. In S. Misener and S. A. Krawetz [eds.], *Methods in molecular biology*, vol. 132: Bioinformatics methods and protocols, 365–386. Humana Press, Totowa, New Jersey, USA.
- USFWS. 1987. Endangered and threatened wildlife and plants: Endangered status for *Lupinus aridorum* (scrub lupine). *Federal Register* 52: 11172–11175.
- WUNDERLIN, R. P., AND B. F. HANSEN. 2008. Atlas of Florida vascular plants. Institute for Systematic Botany, University of South Florida, Tampa, Florida, USA.