



## **Nest Site Selection of Montagu's Harrier *Circus pygargus* Breeding in Natural Habitats in Eastern Poland**

Author: Wiącek, Jarosław

Source: *Ardea*, 97(1) : 117-119

Published By: Netherlands Ornithologists' Union

URL: <https://doi.org/10.5253/078.097.0114>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Nest site selection of Montagu's Harrier *Circus pygargus* breeding in natural habitats in eastern Poland

Jarosław Wiącek<sup>1</sup>

Wiącek J. 2009. Nest site selection of Montagu's Harrier *Circus pygargus* breeding in natural habitats in eastern Poland. *Ardea* 97(1): 117–119.



Nest site selection of the ground-nesting Montagu's Harrier was studied in a natural habitat in eastern Poland, the Chełm Calcareous Marshes (850 ha). All nests in the study area were placed in the sedge community *Cladietum marisci*. The water at nest sites was deeper, and the vegetation was higher, denser and broader-leaved than found at random points. Distance to the nearest meadows did not differ between nest sites and random points; all nests were placed over water, though. These data suggest a preference for nesting in wet places with older and higher vegetations growing in high density.

Key words: *Circus pygargus*, nest site selection, natural habitat, Poland

<sup>1</sup>Department of Nature Conservation, Biology Institute, Curie-Skłodowska University, Akademicka 19 Street, 20-033 Lublin, Poland (wiacek@hektor.umcs.lublin.pl)

Marshes and grasslands are typical breeding areas of Montagu's Harriers (Simmons 2000). The primeval breeding habitat of Montagu's Harrier used to be steppe and marshland, but cereal crops and alfalfa fields have become main breeding habitats in western Europe in the second half of the 20th century (Arroyo *et al.* 2004). However, in Poland most harriers still nest in their natural habitat (Tomiałojć & Stawarczyk 2003), as in my study area in Chełm's Calcareous Marshes (Wiącek 2008).


Vegetation height is a main criterion determining the nest location in cereal crops (Arroyo *et al.* 2003), as it is in the choice of the exact location of nests within a vegetation patch in natural shrubland in Castellón in Spain (Limiñana *et al.* 2006). In this study, I evaluate the influence of vegetation height, depth of water, density of the vegetation near the nest and leaf structure on the nest site selection of Montagu's Harriers.

## Study area and methods

The data were collected in the calcareous marshes of the Chełm region near the Polish-Ukrainian border in the nature reserves of 'Bagno Serebryskie' and 'Roskosz' in 2007 (Fig. 1A). The combined surface area encom-

passed 850 ha (51°10'N, 23°37'E). The marshlands are dominated by the sedge community *Cladietum marisci*. For a more detailed description of the study area, see Wiącek & Niedźwiedź (2005).

The field work was based on the methods described by Tyler *et al.* (1998), with modifications when necessary. Vegetation height, depth of water and vegetation density were measured at 19 observed nests from 28 pairs breeding in the area and at 25 random control points. Vegetation heights were measured (above water level) during the incubation period between 15 May and 14 June. Mean vegetation height was calculated from eight measurements with an accuracy of 1 cm at a distance of 1 m from the nest. The same focal points were used to measure water depth. Width of the sedge leaves was measured at eight points around the nest with an accuracy of 1 mm, at 0.5 m above the water level (dominant plant species around the nests). The vegetation density was measured near the nest (at a distance of 0.5 m from the nest) in plots of 0.1 m<sup>2</sup> at a height of 0.5 m above the water level. The distance of the nest to nearby meadows was measured with a measuring tape to the nearest meter. The random control points were placed irrespective of nests in semi-



**Figure 1.** A. Special bird protection area near Chelm in eastern Poland, showing study sites. B. Nests and random points in the 'Bagno Serebryskie' nature reserve.

colonies and randomly across the study area (Fig. 1B). Percentage plant cover near nests and random points was estimated in five categories: + single plants, 1–25%, 26–50%, 51–75%, 76–100%. Analyses were made with nonparametric statistics (Mann–Whitney test). Analyses were performed with Statistica 6.1. Data are presented as means and medians.

**Results**

Harriers built their nests mainly in Saw Sedge *Cladium mariscus* dominated vegetations (Table 1). Nests were situated in vegetations standing in deeper water than found at random points. Harriers also preferred higher,

**Table 1.** Vegetation found near Montagu's Harrier nests ( $n = 19$ ) and at random points ( $n = 25$ ). (+ = single plants).

Plant species	Plant cover (%)	
	near nests	random points
<i>Cladium mariscus</i>	76–100	70–100
<i>Phragmites communis</i>	5–25	0–30
<i>Filipendula ulmaria</i>	+	
<i>Frangula alnus</i>	+	+
<i>Lysimachia vulgaris</i>	+	+
<i>Lythrum salicaria</i>	+	
<i>Galium uliginosum</i>	+	+
<i>Thelypteris palustris</i>	2–15	+
<i>Cirsium palustris</i>	+	
<i>Lathyrus paluster</i>	+	
<i>Potentilla erecta</i>	+	
<i>Campanula patula</i>	+	+
<i>Salix cinerea</i>	+	+
<i>Salix rosmarinifolia</i>	+	+
<i>Origanum vulgare</i>	+	
<i>Convolvulus arvensis</i>		+

denser and broader-leaved vegetation (Table 2). Birds built nests near meadows but differences with random points were not statistically significant.

**Discussion**

Montagu's Harriers preferred high vegetations with older, broad-leaved sedges. Vegetation height at nests in the study area was higher than found at nests in natural habitats in France and Kazakhstan or in cereal crops in western Europe (Arroyo *et al.* 2004). The opposite was found in Castellón province in Spain, where the dense Mediterranean shrubland necessitated nest-building in small patches of shorter vegetation to allow

**Table 2.** Variables measured at nests of Montagu's Harriers and random points.

	Nests ( $n = 19$ )			Random points ( $n = 25$ )			Mann–Whitney test	
	Average $\pm$ SD	Median	Range	Average $\pm$ SD	Median	Range	Z	P
Water depth (cm)	18.5 $\pm$ 13.1	19	1–40	15.1 $\pm$ 12.8	8	0.5–36	2.96	0.003
Vegetation height (cm)	112 $\pm$ 11.4	115	86–122	104.2 $\pm$ 11.6	104	81–120	2.51	0.01
Vegetation density (in 0.1 m <sup>2</sup> )	128 $\pm$ 26.3	118	95–192	67 $\pm$ 13.4	63	48–95	5.61	0.001
Width of leaves (mm)	12.3 $\pm$ 0.7	12	11–13.5	9.2 $\pm$ 1.1	9	7–11.5	5.53	0.001
Distance to meadows (m)	80.4 $\pm$ 50.8	63	11–165	60.4 $\pm$ 28.7	65	15–125	0.17	0.86

for safe take-offs and landings (Limiñana *et al.* 2006). High vegetations may deter predators from approaching the nest, and from spotting the nest from the air. Local predators known to occur in eastern Poland include Red Fox *Vulpes vulpes*, American Mink *Mustela vison*, Eurasian Marsh Harrier *Circus aeruginosus*, Hooded Crow *Corvus cornix* and Common Magpie *Pica pica* (Kitowski 1998, Polak 2007, Wiącek unpubl. data). All nests in the study area were built over water, and water depth was found to influence nest site selection. Water probably increases nest safety against mammals predators, although absence of water does not deter Montagu's Harriers from nesting in dry habitats (Arroyo *et al.* 2004). Most nests in the study area were situated within 100 m from the border of the marsh (Table 2). This trend to build nests near the meadow's border could be the result of a trade-off between better take-off conditions for fledglings during food transfers and risks from ground predators. Young birds from nests situated near meadows start hunting earlier, possibly because of a higher risk from ground predators (Krogulec 1992, Wiącek 2006a).

The older patches of vegetation, i.e. the nest sites preferred by Montagu's Harriers, are illegally burned sometimes. Burning has a strong, negative impact on breeding numbers of this species in the study area (Wiącek 2006b). After burning, it takes almost five years before the habitat again attracts breeding Montagu's Harriers, probably because the regrowth is at first slender-leaved and has a low density, offering little protection to the nest against predators. A similar situation was described in the Bittern *Botaurus stellaris* where density and age of the vegetation near the nest were found to be important factors limiting nest safety and breeding success (Polak 2007).

## References

- Arroyo B.E., Bretagnolle V. & Garcia J.T. 2003. Land use, agricultural practices and conservation of Montagu's Harrier. In: Thompson D.B.A., Redpath S.M. & Marquiss M. (eds) Raptors in a changing environment. JNCC, Peterborough, pp. 449–463.
- Arroyo B.E., Garcia J.T. & Bretagnolle V. 2004. Montagu's Harrier *Circus pygargus*. The Birds of Western Palearctic Update, Vol. 6: 41–55.
- Kitowski I. 1998. Group mobbing on birds and foxes by the Montagu's Harrier *Circus pygargus* in the period of fledgling emancipation. Notatki Ornitologiczne 39: 211–217 (In Polish)
- Krogulec J. 1992. Breeding ecology of Montagu's Harrier *Circus pygargus* on calcareous marshes near Chelm. Ph.D Thesis University of Maria Curie-Skłodowska, Lublin. (In Polish)
- Limiñana R., Soutullo A., Urios V. & Surroca M. 2006. Vegetation height selection in Montagu's Harriers *Circus pygargus* breeding in a natural habitat. Ardea 94: 280–284.
- Polak M. 2007. Nest-site selection and nest predation in the Great Bittern *Botaurus stellaris* population in eastern Poland. Ardea 95: 31–38.
- Tomiałojć L. & Stawarczyk T. 2003. Avifauna of Poland – distribution, abundance and changes. Polskie Towarzystwo Przyjaciół Przyrody 'pro Natura', Wrocław. (in Polish)
- Tyler G.A., Smith K.W. & Burges D.J. 1998. Reedbed management and breeding Bitterns *Botaurus stellaris* in the UK. Biol.Conserv. 86: 257–266.
- Wiącek J. 2006. Pair formation in the Montagu's Harrier. Berkut 15: 151–158.
- Wiącek J. 2006a. Food transfer in Montagu's Harrier *Circus pygargus* during the courtship. Acta Ornithol. 41: 88–91.
- Wiącek J. 2006b. The fluctuation in the number of Montagu's Harrier *Circus pygargus* in the context of the change in the protection status and utilization methods on the territory of Chelm's Calcareous Peat Bogs. Polish J. Environ. Stud. 15: 737–741.
- Wiącek J. 2008. Benefits and costs of semi-colonial breeding in the Montagu's Harrier *Circus pygargus*. Belg. J. Zool. 138: 36–40.
- Wiącek J., Niedźwiedz M. 2005. The food of the Montagu's Harrier during pre-laying period. Berkut 14: 189–192.

## Samenvatting

Tegenwoordig broeden in West-Europa de meeste Grauwe Kiekendieven *Circus pygargus* in landbouwgewassen. In Polen is dat nog niet het geval. Daar zijn natuurlijke habitats het meest in gebruik als broedplaats, zo ook in het onderhavige onderzoek in een moeras in Oost-Polen. Een vergelijking tussen variabelen gemeten bij nesten en op willekeurige punten in het moeras leerde dat Grauwe Kiekendieven een voorkeur aan de dag legden voor zeggenvegetaties in water, waarbij de vegetatie dichter, hoger en ouder (dat laatste afgeleid uit de bredere bladeren van zeggen) was dan op de willekeurig gekozen punten. Deze voorkeur voor de oudere vegetatie zou een anti-predatiestrategie kunnen zijn. (RGB)

Corresponding editor: Rob G. Bijlsma

Received 9 April 2008; accepted 3 February 2009

# ARDEA

TIJDSCHRIFT DER NEDERLANDSE ORNITHOLOGISCHE UNIE (NOU)

*ARDEA* is the scientific journal of the Netherlands Ornithologists' Union (NOU), published bi-annually in spring and autumn. Next to the regular issues, special issues are produced frequently. The NOU was founded in 1901 as a non-profit ornithological society, composed of persons interested in field ornithology, ecology and biology of birds. All members of the NOU receive *ARDEA* and *LIMOSA* and are invited to attend scientific meetings held two or three times per year.

## NETHERLANDS ORNITHOLOGISTS' UNION (NOU)

**Chairman** – J.M. Tinbergen, Animal Ecology Group, University of Groningen, P.O. Box 14, 9750 AA Haren, The Netherlands

**Secretary** – P.J. van den Hout, Royal Netherlands Institute for Sea Research (NIOZ), P.O. Box 59, 1790 AB Den Burg, Texel, The Netherlands (hout@nioz.nl)

**Treasurer** – E.C. Smith, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (ekko.diny@planet.nl)

**Further board members** – E. Boerma, G.J. Gerritsen, J. Komdeur, J. Ouweland, G.L. Ouweneel, J.J. de Vries

**Membership NOU** – The 2009 membership fee for persons with a postal address in The Netherlands is €42 (or €25 for persons <25 years old at the end of the year). Family members (€9 per year) do not receive journals. Foreign membership amounts to €54 (Europe), or €65 (rest of the world). Payments to Postbank account 285522 in the name of Nederlandse Ornithologische Unie, 8897HZ Oosterend-Terschelling, The Netherlands (BIC: PSTBNL21 and IBAN: NL85 PSTB 0000 285522). Payment by creditcard is possible.

Correspondence concerning membership, payment alternatives and change of address should be sent to: J.J. de Vries, Oosterend 10 b, 8897 HZ Oosterend-Terschelling, The Netherlands (jacobird@xs4all.nl).

**Research grants** – The NOU supports ornithological research and scientific publications through its Huib Kluijver Fund and the 'Stichting Vogeltrekstation'. Applications for grants can be addressed to the NOU Secretary. Donations to either fund are welcomed by the NOU treasurer.

**Internet** – [www.nou.nu](http://www.nou.nu)

## ARDEA

**Editors of *ARDEA*** – Rob G. Bijlsma, Wapse (Editor in chief); Christiaan Both, Groningen; Niels J. Dingemanse, Groningen; Dik Heg, Bern; Ken Kraaijeveld, Leiden; Jouke Prop, Ezinge (Technical editor); Julia Stahl, Oldenburg; B. Irene Tieleman, Groningen; Yvonne I. Verkuil, Groningen

**Dissertation reviews** – Popko Wiersma, Groningen

**Editorial address** – Jouke Prop, Allersmaweg 56, 9891 TD Ezinge, The Netherlands (ardea.nou@planet.nl)

**Graphics** – Dick Visser, Haren

**Artwork** – Jos Zwarts, Bunnik

**Internet** – [www.ARDEAJournal.nl](http://www.ARDEAJournal.nl)

**Subscription *ARDEA*** – Separate subscription to *Ardea* is possible. The 2009 subscription rates are €35 (The Netherlands), €41 (Europe), and €49 (rest of the world). Institutional subscription rates are €52, €67, and €76, respectively). Payments to Postbank account 125347, in the name of Nederlandse Ornithologische Unie, Ir. van Stuivenbergweg 4, 6644 AB Ewijk, The Netherlands (BIC: PSTBNL21 and IBAN: NL65 PSTB 0000 125347). Correspondence concerning subscription, change of address, and orders for back volumes to: J.J. de Vries, Oosterend 10 b, 8897 HZ Oosterend-Terschelling, The Netherlands (jacobird@xs4all.nl).

**Exchange of publications** – All periodicals sent in exchange for *ARDEA* should be addressed to: Library of the Netherlands Ornithologists' Union (NOU), c/o Tineke Prins, Institute of Systematics and Population Biology, Zoological Museum, P.O. Box 94766, 1090 GT Amsterdam, The Netherlands.

**Books for review** – should be addressed to: *ARDEA* Secretariat, c/o J. Prop, Animal Ecology Group, Biological Centre, P.O. Box 14, 9750 AA Haren, The Netherlands. After review, the books will be deposited in the NOU Library in Haren.

**NOU Library (journals)** – Mauritskade 57, Amsterdam, Mo–Fr 10:00–16:00 (to check in advance by telephone + 31 20 525 6614).

**NOU Library (books)** – Library Biological Centre, Kerklaan 30, Haren (G.), Mo–Thu 09:00–17:00 (to check at [www.rug.nl/bibliotheek/locaties/bibfwn/index](http://www.rug.nl/bibliotheek/locaties/bibfwn/index)).

© Nederlandse Ornithologische Unie (NOU), 2009

Layout by Dick Visser, Haren, The Netherlands

Printed by Van Denderen, Groningen, The Netherlands, April 2009

Downloaded From: <https://complete.bioone.org/journals/Ardea> on 08 May 2025

Terms of Use: <https://complete.bioone.org/terms-of-use>