



Movements of Immature European Honey Buzzards *Pernis apivorus* in Tropical Africa

Authors: Strandberg, Roine, Hake, Mikael, Klaassen, Raymond H.G.,
and Alerstam, Thomas

Source: Ardea, 100(2) : 157-162

Published By: Netherlands Ornithologists' Union

URL: <https://doi.org/10.5253/078.100.0207>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Movements of immature European Honey Buzzards *Pernis apivorus* in tropical Africa

Roine Strandberg^{1,*}, Mikael Hake², Raymond H.G. Klaassen³ & Thomas Alerstam¹

Strandberg R., Hake M., Klaassen R.H.G. & Alerstam T. 2012. Movements of immature European Honey Buzzards *Pernis apivorus* in tropical Africa. *Ardea* 100: 157–162.



Immature European Honey Buzzards *Pernis apivorus* are believed to remain in tropical Africa during the first years of their lives. We studied their movements during this period with the aid of satellite telemetry. After crossing the Sahara Desert on autumn migration, all six tracked young buzzards stopped at relatively northerly latitudes, between 9.9–13.6°N. Of the five individuals that continued transmitting, four made south-directed movements, mainly in November, to areas located further south or east within latitudes 1.7–9.8°N. Three young buzzards were tracked for more than three months in tropical Africa, and these individuals continued to perform extensive movements within the tropics throughout the tracking period. They travelled between 2,430 and 3,990 km (minimum distances) during 13 to 14 months, in which they visited several sites. In contrast, adult birds migrate directly to their wintering sites where they remain stationary within restricted territories. The mobile life of young Honey Buzzards during the period prior to their first northbound migration may be associated with responses to seasonal weather changes in the tropics and prospecting behaviour. These movements may also reflect intraspecific competition which might be catalyzed by forest degradation and fragmentation.

Key words: European Honey Buzzard, satellite tracking, nomadic movements, tropical Africa

¹Department of Biology, Lund University, Ecology Building, SE-223 62 Lund, Sweden; ²Varpmossevägen 8, SE-436 39 Askim, Sweden; ³Dutch Montagu's Harrier foundation and Animal Ecology Group, Centre for Ecological and Evolutionary Studies, P.O. Box 11103, 9700 CC Groningen, The Netherlands; *corresponding author (Roine.strandberg@biol.lu.se)

Our knowledge about the ecology of long-distance migrating birds in the tropics is limited, despite the fact that this period is of the utmost importance to understand population changes (for a review see Zwarts *et al.* 2009). Of particular interest is the ranging behaviour of immature birds of long-lived species, such as raptors, which spend up to several years in the non-breeding range before returning to their breeding grounds (e.g. Mellone *et al.* 2011 and references therein). The development of satellite transmitters small enough to be carried by birds has made it possible to track individuals throughout the annual cycle. The focus of satellite tracking studies has hitherto been the process of animal migration. However, there are also several studies concerning the wintering behaviour of tracked migrants, such as age differences in ranging behaviour of wintering Gyrfalcons *Falco rusticolus* (Burnham &

Newton 2011) and young Egyptian Vultures *Neophron percnopterus* (Meyburg *et al.* 2004), flexibility and site fidelity in wintering areas of White Storks *Ciconia ciconia* (Berthold *et al.* 2001, Berthold *et al.* 2004), within-Africa movements of immature Short-toed Eagles *Circus gallicus* (Mellone *et al.* 2011), and habitat-related winter-movements of Montagu's Harriers *Circus pygargus* (Trierweiler *et al.* 2012).

In this study we investigate the movements of young European Honey Buzzards *Pernis apivorus* after reaching tropical Africa south of the Sahara. Honey Buzzards normally start breeding when they are three years or older (Roberts 2011; R.G. Bijlsma pers. comm.) and remain in the wintering areas until they are two years old (Gamauf & Friedl 2011). Thus, information about the whereabouts and habits of young Honey Buzzards during their first visit to the wintering

areas may be important from a conservation perspective, as survival and carryover effects are, to a large extent, influenced by what the birds encounter at the wintering grounds in Africa.

METHODS

From 1997 to 2005 we attached satellite transmitters (PTT-100s, Microwave Telemetry Inc.) as backpacks to seven adult and eight juvenile Honey Buzzards at nest sites in southwestern Sweden between 57–59°N and 12–13°E (autumn migrations described in Hake *et al.* 2003). Juvenile birds included siblings from two different nests; two young (No. 2 and 3) were tagged at the same nest in 2000, and three young (No. 4, 5 and 6) were tagged at another nest during consecutive seasons (No. 4 born in 2004; No. 5 and 6 born in 2005).

To study post-migratory behaviour, we used positions from the period after the birds had completed autumn migration and reached tropical Africa south of the Sahara. For defining stationary periods, transit travels and post-juvenile movements (see definitions below) we mainly used the more precise ARGOS-class 1–3 positions (<http://www.argos-system.org/manual>). Class 0 positions were only used if there were two or

more locations separated by less than 10 km at roost sites or by less than 25 km during longer stationary periods (e.g. when only receiving a few class 0 positions from an area during a period of one month).

We classified movements as (see Table 1):

(1) Autumn migration movements – a southward directed movement from the nesting area to the first stopover area south of the Sahara, where the birds stopped for at least 7 days (one week) in a restricted area (moving <10 km from the location on the first day).

(2) Transit movements – a directed flight away from the first stopover area south of the Sahara to an area where the bird stopped >30 days (one month) before making nomadic movements. All observed transit movements consisted of daily flights of >50 km/day.

(3) Post-juvenile movements – movements during the period in tropical Africa prior to the first northbound migration, typically back and forth to one or several staging sites where the bird was stationary for longer periods. These movements were recorded during several days and over distances of hundreds of km. Transmission from juveniles No. 1, 2 and 3 ended during their first winter (October–January), while the other three young were tracked during 14–19 months after arriving in tropical West Africa (Table 1).

Table 1. Timing of movements (shaded rows), flight distances and staging areas for young Honey Buzzards in tropical Africa. Transit period: the period for the first directed movement to the major staging area. Number of positions used for calculating transit and post-juvenile movements is divided into class 3, 2, 1, and 0 positions. Note that individuals 2 and 3 stopped transmitting during their first winter in Africa (= post-juvenile movements unknown). Tracks of individuals 2–6 are shown in Figure 1.

Individual	1	2	3	4	5	6
Arrival date	7 Oct 1998	14 Oct 2000	15 Oct 2000	9 Oct 2004	4 Oct 2005	12 Oct 2005
Arrival site (coordinates)	NW Nigeria (12.1N, 6.2E)	SW Mali (11.0N, 7.8W)	NW Nigeria (13.6N, 5.8E)	S Burkina Faso (9.9N, 3.3W)	N Ivory Coast (10.3N, 6.4W)	N Ivory Coast (10.3N, 6.6W)
Transit period	-	2–13 Nov	23 Oct–14 Nov	-	7 Nov–4 Dec	4–28 Nov
Distance (km)	-	951	928	-	3699	1889
Number of positions	-	2,1,2,0	1,3,2,0	-	8,7,2,1	2,7,1,1
Staging area (coordinates)	-	SW Ivory Coast (5.2N, 6.2W)	C Togo (7.7N, 0.9E)	Ivory Coast-W Ghana (6.4–9.8N, 2.3–6.3W)	D.R. Congo-C.A. Republic (1.7–8.0N, 15.4–22.1E)	Benin-Togo-Ghana (6.7–9.3N, 1.3W–2.7E)
Start date post-juv movement	-	-	-	14 Nov 2004	23 Feb 2006	26 Mar 2006
Distance (km)	-	-	-	2430	3990	2457
Number of positions	-	-	-	2,6,7,4	12,7,2,3	7,7,1,0
Last message (coordinates)	23 Oct 1998 (12.1N, 6.4E)	31 Dec 2000 (5.2N, 6.2W)	8 Jan 2001 (7.7N, 0.9E)	19 Dec 2005 (9.0N, 4.8W)	23 Mar 2007 (5.5N, 20.9E)	10 May 2007 (6.8N, 2.6E)




Figure 1. Post-juvenile movements made by young Honey Buzzards in tropical Africa with NDVI for November 2005 (representing the vegetation index at the time when adults are stationary in the area while juveniles perform their transit movements). Individuals are presented in Table 1. The stars indicate wintering sites for tracked adult Honey Buzzards from Sweden (Hake *et al.* 2003).

RESULTS

Six out of seven adult Honey Buzzards reached their wintering grounds. These birds travelled directly to wintering sites at latitudes 4.4–8.8°N. The four individuals that were tracked during 1 to 4 months after arrival remained stationary within areas less than 53 km² according to high quality positions (wintering sites shown in Figure 1; see also Hake *et al.* 2003).

Out of eight juvenile birds, six arrived south of the Sahara. The two juveniles that did not reach tropical Africa stopped transmitting during autumn migration in Denmark and the Mauritanian desert, respectively. All six successful juveniles stopped in the Guinea savanna zone, between latitudes 9.9–13.6°N, where they arrived between 4 and 15 October. These sites ranged from Mali in the west to Nigeria in the east (Table 1). Juvenile No. 1 stopped transmitting from the latter site by 23 October.

Of the remaining five birds, four individuals (No. 2, 3, 5 and 6) initiated transit movements between 23 October and 7 November and arrived at their main wintering sites between 13 November and 4 December. These movements were first directed to the south, after which the birds turned eastwards as soon as they reached the coast (Figure 1). The fifth individual (No. 4) started to move on 14 November 2004 from its first

post-migratory site, but this movement was not considered to be a true transit movement, as the bird did not continue to a staging area further south. Instead, this and subsequent movements strongly resembled the post-juvenile movements observed for the other juveniles (see below), in which the bird regularly returned to latitudes close to the post-migratory area (reaching northern Ivory Coast). This individual was tracked until 19 December 2005, when transmission ceased 13 months after the first post-juvenile movement within the tropics.

The siblings from 2005 (individuals No. 5 and 6) were also tracked for extensive periods after their November transit movement to their main wintering area. These individuals initiated nomadic movements two and a half and four months after arrival to the main staging areas, respectively. These birds continued to make such post-juvenile movements until the last contacts with these birds in March and May 2007; 13 months after their first post-juvenile movements had started.

Post-juvenile movements were recorded mainly in February–May and October–December (Figure 2). As the solar-panel powered satellite transmitters seem to work inefficiently inside forest with closed canopy, longer periods without contact with the birds suggest stationary behaviour (Figure 2). Concerning the three




Figure 2. The monthly distribution of distance flown by immature Honey Buzzards throughout the transmission period. The different types of movement are defined in the methods section. During the periods of migration, transit movements and post-juvenile movements, the numbers of recorded birds were 6, 4, and 3; the numbers of positions were 61, 39, and 59, respectively. Bars show the monthly proportion of recorded flight distance for each category of movement (y-axis on left-hand side). The total number of days with positions received from the transmitters during the post-juvenile period are given by the dashed line (y-axis on right-hand side).

juveniles transmitting for a longer period of time in Africa (No. 4, 5 and 6), very few positions were obtained during their first summer in Africa. For individual No. 4 only one position was obtained between the end of April and the end of October, for individual No. 5 positions were obtained during only one transmission cycle (8 h) between mid-May and September, and for individual No. 6 no positions were obtained from the beginning of June until the end of March.

DISCUSSION

After completing autumn migration and arriving in tropical West Africa, young Honey Buzzards differed in their behaviour from adults in two main ways: (1) the young birds used post-migratory stopover sites in the Guinea savanna zone during the first 1–3 months after their Sahara crossing, after which they continued southwards by transit movements, and (2) the young birds made extensive nomadic movements during their stay in Africa. In contrast, adult Honey Buzzards seem to travel directly to well-defined wintering sites where they remain stationary (Hake *et al.* 2003; also shown by Meyburg & Meyburg 2011). In a Dutch tracking study (www.uva-bits.nl) most adults showed similar behaviour during the winter. However, some adults

used several well-defined home ranges far apart (from West Africa to Equatorial Guinea) during the same winter (Jan van Diermen, Willem van Manen & Willem Bouten unpubl.), i.e. similar to the juveniles tracked in this study. This is interesting, but further studies are required to investigate whether this is a common behaviour among some adults or if such movements may be triggered by deteriorating conditions in their regular wintering area due to e.g. weather conditions or disturbance.

The use of post-migratory stopover sites has earlier been found to occur among both adult and juvenile Marsh Harriers *Circus aeruginosus* and Montagu's Harriers, in which birds seem to explore temporal feeding opportunities at the northern edge of the Sahel (Strandberg *et al.* 2008, Trierweiler *et al.* 2012). In the Honey Buzzard, sub-Saharan stopovers were used only by juvenile birds. The juveniles have no previous experience of migratory journeys, and it seems reasonable that they, after crossing the Sahara, stop as soon as they find a suitable feeding habitat, as their energy reserves may be exhausted. The timing of movements to more southern sites suggested that transit movements may have been associated with the start of the dry season, causing reduced foraging opportunities in the northern staging areas.

Transit movements were directed south-southwest, in accordance with the general migration direction, but the birds moved eastwards as soon as they reached the coastline. Similarly, all four juveniles (out of seven birds) that reached the southern coastline of West Africa in an Austrian tracking study changed their orientation from south(-west) to east as soon as the coast was reached (Gamauf & Friedl 2011). Thus, it seems a general behaviour of immature inexperienced Honey Buzzards to continue their travels in an eastward direction when confronted with the coast, which will enhance their chances to find suitable wintering habitat as tropical forest is more abundant in Central Africa. Interestingly, such eastward movement is against the prevailing easterly winds (Kemp *et al.* 2010), suggesting that this change in orientation is an innate response rather than an effect of wind drift.

All three immature birds that were tracked for longer periods showed extensive irregular movements which occurred during late winter to spring and during late autumn to early winter. Most of these post-juvenile movements occurred at the time when adult birds migrate. However, considerable amounts of post-juvenile movements were also observed during winter, when the adults, at least the majority, are strictly stationary. During the breeding period of Honey

Buzzards (June–August) relatively few observations of post-juvenile movements were obtained, but these positions suggest that the birds were stationary in Africa during the northern summer. This implies that post-juvenile movements are regulated by an internal annual clock, related to periods of migratory restlessness. Alternatively, these movements may be a response to changing environmental conditions, and thus related to local dry and wet seasons. Periods of intensive rainfall may provide unfavourable feeding conditions for Honey Buzzards, while dry seasons might promote nomadic behaviour to explore better feeding opportunities. In this respect, it is interesting to note that the irregular post-juvenile movements took place between seemingly similar habitats (tropical forest), in contrast to the transit flights when the birds moved from open, woody savanna into tropical forest habitat. Another explanation is that the post-juvenile movements are related to the fact that the adult's northbound migration to the breeding areas means that there will suddenly be an increase in feeding areas available during a period of four months (when the adults have left Africa and migrated to temperate breeding grounds), which probably are of higher quality provided that adults are dominant over juveniles. Hence, immature birds may move to occupy the adult's deserted winter territories in spring and move away when the adults return during autumn.

The age-related behavioural differences in tropical Africa, with immatures moving around while most adults remain stationary in small territories, might also be related to intraspecific competition, where juveniles are forced to search for suitable habitats and unoccupied territories as they arrive later to the wintering area. The degradation and fragmentation of tropical forests might catalyze this behaviour as relatively little suitable habitat where individuals can stay for longer periods might be left (see also Bijlsma 2002). Although we have no idea if carrying capacity is currently reached for Honey Buzzards in Africa, the ongoing rapid loss of natural forests in Africa, the preferred wintering habitat of our satellite transmitter birds, is unlikely to be beneficial for Honey Buzzards in general.

Interestingly, the young Honey Buzzard tracked for two years (No. 6) did not show any tendency to move northwards during its second summer in Africa, indicating that at least some Honey Buzzards are not returning to the breeding area until their third spring season as 4-cy birds. That some Honey Buzzards already return to Europe in their second spring (as a 3-cy bird) is shown by an individual tracked from Austria (Gamauf & Friedl 2011).

ACKNOWLEDGEMENTS

The study was approved by the Ethical Committees in Uppsala (C92/6, C179/9) and Malmö/Lund (204-06), with permission to capture the birds from the Swedish Environmental Protection Agency (412-429-05 Nf). We are especially grateful to Bo Kanje, Sonja Kanje, Nils Kjellén, Jan Bergkvist and Leif Danielsson for field assistance. Thanks to Rob G. Bijlsma for constructive comments very much improving the manuscript. We are also grateful for excellent support from Microwave Telemetry Inc. We wish to acknowledge use of the Maptool program for graphics in this paper (www.seaturtle.org). This research was financed through grants from the Swedish Research Council.

REFERENCES

- Alerstam T., Hake M. & Kjellen N. 2006. Temporal and spatial patterns of repeated migratory journeys by ospreys. *Anim. Behav.* 71: 555–566.
- Berthold P., van den Bossche W., Fiedler W., Kaatz C., Kaatz M., Leshem Y., Nowak E. & Querner U. 2001. Detection of a new important staging and wintering area of the white stork *Ciconia ciconia* by satellite tracking. *Ibis* 143: 450–455.
- Berthold P., Kaatz M. & Querner U. 2004. Long-term satellite tracking of white stork (*Ciconia ciconia*) migration: constancy versus variability. *J. Ornithol.* 145: 356–359.
- Bijlsma R.G. 2002. Life-history traits of honey buzzards (*Pernis apivorus*) in Africa. *Vogelwarte* 41: 240–248.
- Burnham K.K. & Newton I. 2011. Seasonal movements of gyrfalcons *Falco rusticolus* include extensive periods at sea. *Ibis* 153: 468–484.
- Gamauf A. & Friedl C. 2011. Zug- und Überwinterungsstrategien junger Wespenbussarde *Pernis apivorus*. *Vogelwarte* 49: 290–291.
- Hake M., Kjellén N. & Alerstam T. 2003. Age-dependent migration strategy in honey buzzards *Pernis apivorus* tracked by satellite. *Oikos* 103: 385–396.
- Kemp M.U., Shamoun-Baranes J., van Gasteren H., Bouten W. & van Loon E.E. 2010. Can wind help explain seasonal differences in avian migration speed? *Journal of Avian Biology* 41: 672–677.
- Mellone U., Yáñez B., Limiñana R., Muños A.R., Gonzáles J.M., Urios V. & Ferrer M. 2011. Summer staging areas of non-breeding short-toed snake eagles *Circaetus gallicus*. *Bird Study* 58: 516–521.
- Meyburg B.U., Max Gallardo M., Meyburg C. & Dimitrova E. 2004. Migrations and sojourn in Africa of Egyptian vultures (*Neophron percnopterus*) tracked by satellite. *J. Ornithol.* 145: 273–280.
- Meyburg B.-U. & Meyburg C. 2011. Satellitentelemetrische Untersuchungen an adulten deutschen Wespenbussarden. *Vogelwarte* 49: 280–281.
- Roberts S. 2011. Honey Buzzards in Wales 2011. *Boda Wennol* 2011: 12–13.
- Strandberg R., Klaassen R.H.G., Hake M., Olofsson P., Thorup K. & Alerstam T. 2008. Complex timing of marsh harrier *Circus aeruginosus* migration due to pre- and post-migratory movements. *Ardea* 96: 159–171.

- Trierweiler C., Mullié W.C., Drent R.H., Exo K.M., Komdeur J., Bairlein F., Harouna A., de Bakker M. & Koks B.J. 2012. A Palaearctic migratory raptor species tracks shifting prey availability within its wintering range in Sahel. *J. Anim. Ecol.* doi: 10.1111/j.1365-2656.2012.02036.x
- Zwarts L., Bijlsma R.G., van der Kamp J. & Wymenga E. 2009. *Living on the edge: wetlands and birds in a changing Sahel.* KNNV Publishing, Zeist.

SAMENVATTING

Om de bewegingen van Wespddieven *Pernis apivorus* gedurende hun eerste levensjaren te volgen, werden zes jonge vogels op het nest voorzien van een satellietzender. Na het oversteken van de Sahara in oktober verbleven de vogels enige tijd in West-Afrika, ruim boven de evenaar op 9,9–13,6° noorderbreedte. In

november trokken vier van de vijf vogels met een nog functionerende zender verder in zuidelijke en zuidoostelijke richting tot op 1,7–9,8° noorderbreedte. Drie Wespddieven waren in tropisch Afrika gedurende lange tijd te volgen. In een periode van 13 tot 14 maanden legden ze 2.430–3.990 km af, waarbij meerdere gebieden werden bezocht. Volwassen vogels laten een ander trekpatroon zien door zonder tussenstops naar het overwinteringsgebied te vliegen. Daar blijft hun actieradius vervolgens beperkt tot een klein gebied. De trekbewegingen van jonge Wespddieven in tropisch Afrika hebben mogelijk te maken met weersomstandigheden of exploratief gedrag. Het is ook mogelijk dat jonge Wespddieven voedselconcurrentie met oude vogels uit de weg gaan, al dan niet onder invloed van aantasting van hun leefgebied. (JP)

Corresponding editor: Jouke Prop

Received 29 September 2012; accepted 30 October 2012