



## **Plumage Aberration in Northern Goshawk *Accipiter gentilis***

Authors: Maniarski, Roman, and Ciach, Michał

Source: *Ardea*, 100(2) : 211-213

Published By: Netherlands Ornithologists' Union

URL: <https://doi.org/10.5253/078.100.0214>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Plumage aberration in Northern Goshawk *Accipiter gentilis*

Roman Maniarski<sup>1</sup> & Michał Ciach<sup>2,\*</sup>



Maniarski R. & Ciach M. 2012. Plumage aberration in Northern Goshawk *Accipiter gentilis*. *Ardea* 100: 211–213.

The subspecies of Northern Goshawk *Accipiter gentilis* present clinal plumage variation. However, aberrant plumages are exceptional. On 2 February 2012, in Sędziszów (southern Poland), a dead Northern Goshawk with aberrant plumage was found. The bird was mostly beige but had retained the banding pattern in the flight feathers and streaking on the breast and underwing coverts. Birds in the described aberrant plumage might be confused with the Eastern Palearctic subspecies *A. g. albidus*.

Key words: Northern Goshawk, *Accipiter gentilis*, aberrant plumage, brown mutation

<sup>1</sup>The Wildlife Research and Conservation Society, ul. Sienkiewicza 68, 25-501 Kielce, Poland; <sup>2</sup>Department of Zoology and Wildlife Management, Forest Biodiversity Institute, Faculty of Forestry, University of Agriculture, al. 29 Listopada 46, 31-425 Kraków, Poland;

\*corresponding author (michal.ciach@ur.krakow.pl)

Plumage aberrations have been recorded in many bird species (Sage 1962). These can be restricted to single or multiple feathers, or it may concern the whole body, which then radically changes the appearance of an individual bird (e.g. Graham *et al.* 2005, Vittery 2005, Ciach & Ściborski 2008, Ciach 2009). Plumage aberrations in birds are rarely recorded, given the fact that they usually attract the observer's attention. In most birds of prey, records of abnormal coloration are exceptional (Ciach *et al.* 2011).

On 2 February 2012, a second calendar-year female Northern Goshawk *Accipiter gentilis* with atypical coloration collided with a building in Sędziszów, a town in southern Poland bordered by farmland and abandoned fields (50°34'N, 20°03'E). The plumage was beige (Figure 1). The upperwing coverts were a rather uniform beige except for the marginal coverts which were darker. Scapulars were uniform beige-white. The uppertail coverts and rump had a brownish tinge. The head was beige. Underparts were beige with a carrot-orange tinge, and a distinct brownish streaking on the breast. The underwing coverts, which were darker than the upperwing, showed a dark streaking pattern, and along with the undertail coverts were carrot-orange tinged. All primaries and secondaries were light coffee-coloured and had a brownish banding. The rectrices on the upper and under side had a distinct brown banding. As a result of the coloration, the bird appeared darker

from below. Bare parts were unchanged: yellow legs with blackish claws, grey bill with blackish tip and yellow cere. Body measurements (see Svensson 1992) were: wing (maximum wing chord) – 350 mm; tail – 256 mm; bill length – 32.3 mm; and tarsus – 92 mm. The female was sexed by dissection. The specimen is preserved in the collection of the Natural History Museum at the University of Wrocław, Poland (register number 201588).

The described Northern Goshawk presents a brown aberration. It is defined as a qualitative reduction of eumelanin (phaeomelanin not affected). As a result of an incomplete oxidation of eumelanin, black feathers become dark brown (van Grouw 2006). Those feathers, being sensitive to sunlight, bleach quickly, which explains the contrast between under and upper parts. Our bird differs from several other aberrant Northern Goshawks mentioned in the literature. Aberrant offspring of Northern Goshawk with a genetic defect were reported in the city of Hamburg, where individuals with a pale silver-blue plumage displayed severe locomotor disorders (Rutz *et al.* 2004). Clark (1999) mentioned leucistic (reported as partial albino, which is impossible: a bird is an albino, or not) and dilute plumages of Northern Goshawk, not giving, however, any details. Brown mutation has been recorded recently in Common Buzzard *Buteo buteo* in Poland (Ciach *et al.* 2011).



**Figure 1.** (A) Dorsal and (B) ventral view of a juvenile female Northern Goshawk with brown mutation (2 February 2012, Sędziszów, southern Poland).

Across Eurasia, the Northern Goshawk shows a tendency to become progressively paler from south to north, and from west to east. The dark grey plumage of European *A. g. gentilis* shifts into the paler forms of the eastern races *A. g. buteoides* and *A. g. albidus* (Cramp & Simmons 1980). The latter subspecies, occurring from north-east Siberia to Kamchatka Peninsula, is conspicuously pale (Kretchmar & Probst 2003). The nominate Northern Goshawk in aberrant plumage, as described in this note, may be confused with individuals of the Eastern Palearctic subspecies *A. g. albidus*.

We are grateful to Alojzy Przemyski, who provided the specimen, and to Ludwik Maksalon, who measured the specimen. We also thank two anonymous referees for comments on this note.

## References

- Ciach M. 2009. The first record of melanism in the Red-backed Shrike *Lanius collurio*. *Ornis Svecica* 19: 55–56.
- Ciach M. & Ściborski M. 2008. Plumage aberrations in two Wood Sandpipers *Tringa glareola*. *Wader Study Group Bull.* 115: 126.
- Ciach M., Kwarciany A. & Świtała D. 2011. Records of brown plumage aberration in the Common Buzzard *Buteo buteo*. *Ornis Svecica* 21: 119–122.
- Clark W.S. 1999. A field guide to the raptors of Europe, the Middle East, and North Africa. Oxford University Press, Oxford.
- Cramp S. & Simmons K.E.L. (eds) 1980. The Birds of the Western Palearctic. Vol. II. Oxford University Press, Oxford.
- Graham D.M., Collins P.C. & Jessop R.E. 2005. Aberrant plumages in some migratory waders in Australia. *Wader Study Group Bull.* 107: 31–35.

- Kretchmar A.V. & Probst R. 2003. Der weiße Habicht *Accipiter gentilis albidus* in Nordost-Sibirien – Portrait eines Mythos. *Limicola* 17: 289–305.
- Rutz C., Zinke A., Bartels T. & Wohlsein P. 2004. Congenital neuropathy and dilution of feather melanin in nestlings of urban-breeding Northern Goshawks (*Accipiter gentilis*). *J. Zoo Wildl. Medicine* 35: 97–103.
- Sage B.L. 1962. Albinism and melanism in birds. *British Birds* 55: 201–225.
- Svensson L. 1992. Identification Guide to European Passerines. Stockholm.
- van Grouw H. 2006. Not every white bird is an albino: sense and nonsense about colour aberrations in birds. *Dutch Birding* 28: 79–89.
- Vittery A. 2005. Food-induced erythrism in House Sparrow. *British Birds* 98: 434–435.

## Samenvatting

Afwijkingen in de kleur van het verenkleed komen bij roofvogels weinig voor. Hier wordt zo'n geval beschreven voor een eerstejaars vrouwtje Havik *Accipiter gentilis*, dat op 2 februari 2012 werd dood gevonden in het zuiden van Polen. De overheersende kleur was beige. Op de onderzijde (vooral de ondervleugeldeveren) was de kleur wat donkerder door een oranje zweem die over de veren lag. De bandering op de vleugelpennen en de lengtestreping op de borst en de ondervleugeldeveren was zichtbaar gebleven. De kleur van de naakte delen week niet af van wat gewoonlijk bij deze soort wordt aangetroffen. De overwegend bleke impressie die deze vogel bood, zou tot verwarring met de oostelijke ondersoort *A.g. albidus* kunnen leiden. (RGB)

Corresponding editor: Rob G. Bijlsma

Received 13 June 2012; accepted 12 November 2012