

## **First record of *Gracixalus quyeti* (Amphibia: Anura: Rhacophoridae) from Laos: molecular consistency versus morphological divergence between populations on western and eastern side of the Annamite Range**

Authors: Egert, Jennifer, Luu, Vinh Quang, Nguyen, Truong Quang, Le, Minh Duc, Bonkowski, Michael, et al.

Source: Revue suisse de Zoologie, 124(1) : 47-51

Published By: Muséum d'histoire naturelle, Genève

URL: <https://doi.org/10.5281/zenodo.322663>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# First record of *Gracixalus quyeti* (Amphibia: Anura: Rhacophoridae) from Laos: molecular consistency versus morphological divergence between populations on western and eastern side of the Annamite Range

Jennifer Eger<sup>1,7</sup>, Vinh Quang Luu<sup>1,2,7</sup>, Truong Quang Nguyen<sup>3</sup>, Minh Duc Le<sup>4,5,6</sup>, Michael Bonkowski<sup>1</sup> & Thomas Ziegler<sup>1,7\*</sup>

<sup>1</sup> Institute of Zoology, Department of Terrestrial Ecology, University of Cologne, Zùlpicher Strasse 47b, D-50674 Cologne, Germany. E-mail: m.bonkowski@uni-koeln.de

<sup>2</sup> Department of Wildlife, Faculty of Natural Resources and Environmental Management, Vietnam National University of Forestry, Xuan Mai, Chuong My, Hanoi, Vietnam. E-mail: qvinhfuv@yahoo.com.au

<sup>3</sup> Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Hanoi, Vietnam. E-mail: nqt2@yahoo.com

<sup>4</sup> Faculty of Environmental Sciences, Hanoi University of Science, Vietnam National University, 334 Nguyen Trai Road, Hanoi, Vietnam. E-mail: le.duc.minh@hus.edu.vn

<sup>5</sup> Centre for Natural Resources and Environmental Studies, Hanoi National University, 19 Le Thanh Tong, Hanoi, Vietnam

<sup>6</sup> Department of Herpetology, American Museum of Natural History, Central Park West at 79th Street, New York, New York 10024, USA

<sup>7</sup> AG Zoologischer Garten Köln, Riehler Strasse 173, D-50735 Cologne, Germany.

\* Corresponding author, E-mail: ziegler@koelnerzoo.de

**Abstract:** We report the first country record of the poorly known *Gracixalus quyeti* from Laos based on a recently collected specimen from Khammouane Province, central Laos. While the genetic analysis revealed nearly identical sequences, we found some differences in body ratios and color patterns among the specimen from Laos and the type series from the eastern side of the Annamite Range in Vietnam.

**Keywords:** *Gracixalus quyeti* - biogeography - distribution - new record - Annamite Range - Laos.

## INTRODUCTION

Although the knowledge of amphibian diversity in Laos has strikingly increased within the last two decades, distribution patterns and natural history of many species are still poorly known (Luu *et al.*, 2014). Information about the geographic distributions of species is essential for understanding their evolution and ecology and is furthermore crucially required for effective conservation (Nguyen *et al.*, 2014; Rowley *et al.*, 2015). The rhacophorid genus *Gracixalus* currently contains 11 species whose ranges are restricted to lowland and montane forests in China, Thailand, Vietnam and Laos (Nguyen *et al.*, 2008; Rowley *et al.*, 2011; Matsui *et al.*, 2015; Frost, 2016). Luu *et al.* (2014) recorded *G. supercornutus* (Orlov, Ho & Nguyen, 2004) from Laos, which also represented the first record of the genus *Gracixalus* from

this country. However, relatively little is known about this genus, and distribution ranges of species within the genus are largely unknown (Luu *et al.*, 2014).

In this study, we report *Gracixalus quyeti*, a recently described and rare species from Phong Nha – Ke Bang National Park (NP) in central Vietnam, for the first time from Hin Nam No National Protected Area (NPA) in central Laos on the western side of the Annamite Range.

## MATERIAL AND METHODS

**Material examined:** A single female specimen of *Gracixalus quyeti* (VNUF A.2014.73) was collected by Vinh Quang Luu, Thomas Calame, Dung Van Phan and Kieusomphone Thanabuaosy during a field survey in May 2014 in Noong Ma Village (17°17.394' N, 106°09.980'

E, 592 m a.s.l., recorded by a Garmin GPSMAP 60CSx GPS receiver and recorded in datum WGS 84), within Hin Nam No NPA, Boualapha District, Khammouane Province, central Laos (Fig. 1).

**Sampling methods:** The specimen was caught by hand and anaesthetized in a closed vessel with a piece of cotton wool containing ethyl acetate. After taking photographs, the specimen was fixed in 80% ethanol and subsequently transferred into 70% ethanol for permanent storage. A tissue sample was preserved separately in 95% ethanol. The specimen has been deposited in the collection of the Vietnam National University of Forestry (VNUF), Hanoi, Vietnam.

**Molecular analysis:** Total genomic DNA was extracted from a tissue sample using a commercially available DNeasy Tissue Kit following manufacturer's instructions (QIAGEN Inc., Valencia, CA, USA). A fragment of 16S gene was amplified using the primer pair 16Sar + 16Sbr (Palumbi *et al.*, 1991). The standard PCR conditions used for 16S were: 95° C for 5 min., 40 cycles of [95° C for 30 sec., 50° C for 45 sec., 72° C

for 60 sec.] and 72° C for 6 min. All PCR products were visualized on a gel before sequencing. Successful amplifications were purified to eliminate PCR components using GeneJET™ PCR Purification kit (Fermentas, Canada). Purified PCR products were sent to Macrogen Inc. (Seoul, South Korea) for sequencing. The obtained sequence was compared to those available from other species using the BLAST search in GenBank.

**Morphological analysis:** Determination of morphological characters followed Nguyen *et al.* (2008). Measurements were taken by the first author with a digital caliper to the nearest 0.1 mm. Abbreviations were used as follows: SVL: snout-vent length; HL: head length (from the back of mandible to the tip of snout); HW: head width (across angle of jaws); MN: distance from the back of mandible to the nostril; MFE: distance from the back of mandible to the front of eye; MBE: distance from the back of mandible to the back of eye; IFE: distance between the front of eyes; IBE: distance between the back of eyes; IN: internasal distance; EN: distance from the front of eye to the nostril; EL: horizontal eye diameter; NS: distance from nostril to the tip of snout; SL: distance from the front of eye to the tip of snout; TYD: greatest tympanum diameter; TYE: distance from tympanum to the back of eye; IUE: minimum distance between upper eyelids; UEW: maximum width of upper eyelid. Forelimb: HAL: hand length (from the base of outer palmar tubercle to the tip of fourth toe); FLL: forelimb length (from the elbow to the base of outer tubercle); TFL: third finger length (from the base of the first subarticular tubercle to the tip of third toe); fd1-4: width of discs of fingers I-IV; fw1-4: width of fingers I-IV (measured at the narrowest point of the distant phalanx). Hindlimb: FL: femur length (from vent to knee); TL: tibia length; TW: tibia width; FOL: foot length (from the base of inner metatarsal tubercle to the tip of fourth toe); FTL: fourth toe length (from the base of the first subarticular tubercle to the tip of fourth toe); TFOL: distance from the base of tarsus to the tip of fourth toe; IMT: length of the inner metatarsal tubercle; ITL: inner toe length; td1-4: width of discs of toes I-IV; fw1-4: width of toes I-IV (measured at the narrowest point of the distant phalanx). Webbing: MTTF: distance from the distal edge of metatarsal tubercle to the maximum incurvation of web between third and fourth toes; TFTF: distance from the maximum incurvation of web between third and fourth toes to the tip of fourth toe; MTFF: distance from the distal edge of metatarsal tubercle to the maximum incurvation of web between fourth and fifth toes; FFTF: distance from the maximum incurvation of the web between fourth and fifth to the tip of fourth toe. Webbing formula followed Glaw & Vences (2007). Comparative data were taken from Nguyen *et al.* (2008). Institutional abbreviations are as follows:



Fig. 1. Map showing the distribution of *Gracixalus quyeti*, including the localities of the type series after Nguyen *et al.* (2008) in Quang Binh Province, Vietnam (marked with blue dots) and our first record from Khammouane Province, Laos (marked with a red dot).



ZFMK: Zoologisches Forschungsmuseum Alexander Koenig, Bonn, Germany; VNUH: Vietnam National University, Hanoi, Vietnam.

## RESULTS

**Ecology of examined specimen:** The Laotian specimen was found at night (20:30) sitting on a branch of a shrub, ca. 0.5 m above an outcrop at an elevation of 592 m above sea level. The air temperature was 27.3°C and the relative humidity was 80%. The locality was surrounded by limestone cliffs and karst vegetation, mainly consisting of species of Ebenaceae, Dracaenaceae, Arecaeae, Meliaceae, and Moraceae.

**Molecular analysis:** The obtained sequence had 656 bps. Comparative analysis of the obtained sequence with those from GenBank (ZFMK 82999: EU871429.1, VNUH 160706: EU871428.1) showed 99% similarity between the newly collected specimen from Laos (VNUF A. 2014.73) and the holotype (ZFMK 82999) as well as the paratype (VNUH 160706) of *Gracixalus quyeti* from Vietnam. Specifically, the new sequence was different from EU871429.1 in one position and EU871428.1 in two positions.

**Morphological analysis:** The morphological diagnosis coincided with the original description of *G. quyeti* from central Vietnam (Nguyen *et al.*, 2008) in the following characters: Small rhacophorid (ZFMK 82999 SVL 34 mm, VNUF A. 2014.73 SVL 31.4 mm), vomerine teeth absent. Snout rounded, longer than the diameter of eye. Nostrils closer to tip of snout than to eye. Pupil oval and horizontal. Tympanum distinct, rounded and wider than disc of finger III. Dorsal surface of head and body and upper part of flanks with small sharp tubercles. A dark pattern forming an inverse Y, triangular spot between eyes bifurcating into two bands continuing posteriorly onto the back. Brown marbling on margin of throat and throat. Webbing moderately

developed: Ii(1)-(2)IIe(0.5)-(1 2/3)IIIe(0.5)-(2)IVe(2)-(0.5)IV.

Despite the high genetic correspondence, we also noted some differences in morphology between the examined specimen from Laos and the type specimens of *G. quyeti* (Nguyen *et al.*, 2008): *Body ratios:* Head wider than long, in contrast to longer than wide in the type series. Forelimb and hand length slightly shorter, the latter approximately 27% of SVL (30% in the holotype and 28% in the paratype). Length of finger III approximately 16% of SVL and remarkably shorter than in the specimens from Vietnam (24% in the holotype and 19% in the paratype). Hind limb approximately 1.45 times longer than SVL, shorter than in specimens from Vietnam (holotype: 1.6, paratype: 1.7). Tibia 1.5 times thinner than in the holotype. Length of toe IV 23% of SVL, remarkably shorter than in the holotype (37%) and slightly shorter compared to the paratype (25%) (Tab. 1). *Coloration in life:* Dorsal surface of head and body is greenish beige to grayish light-brown with a grayish dark-brown blotching pattern as described above, in contrast to brownish to moss-green with a dark brown pattern in the adult holotype and moss-green with an indistinct pattern in the subadult paratype. Forelimb and dorsal part of hindlimbs are beige to grayish light-brown with grayish dark-brown bands, versus moss-green with dark brown bands in the holotype and the subadult paratype. The ventral surface can only be described based on the preserved specimen: Belly, chest and throat slightly white to yellowish white with brown marbling on margin of throat and throat, versus background color more yellowish in the preserved holotype and more bluish in the living paratype (Fig. 2A-B).

## DISCUSSION

The new record of *Gracixalus quyeti* from Hin Nam No NPA, Laos is approximately 80 km straight line distance apart from the type locality in Quang Binh Province,




Fig. 2. Female of *Gracixalus quyeti* from Hin Nam No NPA, central Laos. (A) Dorsolateral view. (B) Lateral view. Photos: T. Calame.

Table 1. Morphological characters of the newly collected specimen of *Gracixalus quyeti* from Laos in comparison with the type series from Vietnam (after Nguyen *et al.*, 2008; measurements in mm, abbreviations defined in the text).

	VNUF A. 2014.73	ZFMK 82999	VNUH 160706
	female	female, holotype	subadult, paratype
	Hin Nam No NPA, central Laos	Phong Nha – Ke Bang NP central Vietnam	Phong Nha – Ke Bang NP, central Vietnam
SVL	31.4	34.0	22.0
HL	10.3	12.5	8.2
HW	11.73	11.4	7.4
MN	10.3	11.2	7.4
MFE	6.96	8.4	5.4
MBE	3.86	4.5	3.3
IFE	5.45	6.9	4.3
IBE	8.5	11.0	6.8
IN	3.08	3.0	2.1
EN	2.9	3.5	2.5
EL	4.0	4.8	3.2
NS	1.34	2.1	1.3
SL	4.22	5.4	3.8
TYD	2.0	2.5	1.6
TYE	0.92	1.0	0.7
IUE	3.89	4.1	2.4
UEW	2.36	3.0	1.8
HAL	8.4	10.3	6.1
FLL	6.4	7.2	4.9
TFL	4.97	8.1	4.2
fw 1-4	0.31/0.36/0.38/0.36	0.6/0.9/1/0.9	-
fd 1-4	0.48/0.83/0.99/1	1/1.6/1.9/1.7	1.2 (fd 3)
FL	15.68	18.5	11.5
TL	17.61	20.0	11.8
TW	2.57	4.2	2.0
FOL	12.21	15.2	13.9
FTL	7.1	12.5	5.6
TFOL	20.47	24.8	8.1
IMT	0.82	1.3	0.8
ITL	3.83	4.2	1.9
MTTF	7.1	9.4	3.9
MTFF	7.3	9.8	4.7
TFTF	4.71	5.6	3.0
FFTF	4.38	5.0	2.8
tw 1-5	0.28/0.29/0.36/0.49/0.4	0.8/0.9/1.1/1.3/1.1	-
td 1-5	0.35/0.66/0.6/0.96/0.8	0.5/0.6/0.7/0.9/0.8	-

central Vietnam. It is likely that *G. quyeti* is more widespread across the Annamite's extensive limestone areas of central Vietnam and central Laos. The species might potentially be endemic to these fragmented lowland and montane forest habitats, while its occurrence seems to be relatively rare within this presumed distribution range.

All records of *G. quyeti* were derived from different elevations in the limestone area within the transition zone between the northern and central Annamite Mountains, a semiconnected array of hills and forested limestone karst outcrops. The Annamite Range generally experiences a tropical wet monsoon climate, while the eastern oceanic Phong Nha – Ke Bang NP receives more precipitation

and has lower average temperatures than Hin Nam No on the western side in the rain shadow of the Annamite Range (Timmins & Trinh 2001; Sterling *et al.*, 2006; Bain & Hurley, 2011). Taking into account the climatic differences between both sides of the Annamite Range and the lack of knowledge about the genus *Gracixalus*, further research must clarify if the shown morphological differences in coloration and body ratios may possibly reflect ecological adaptations to different environments, evolutionary driven morphs or likely just reflect variation within this poorly known species. Still, our finding elevates the number of amphibian species recorded from Laos to 100, and the number of rhacophorid species known to occur in Laos to 34. Although the species number of amphibians known from Laos has nearly doubled during the last 15 years from 58 to 100 recorded species, the amphibian species richness of Laos is still underestimated (Stuart, 1999; Teynié *et al.*, 2014; Luu *et al.*, 2014; Frost, 2016). In particular big gaps remain in our knowledge of the Annamite's amphibian ecology and distribution.

## ACKNOWLEDGEMENTS

We are grateful to S. Wayakone, S. Bounphanmy, H. Chanthavong, K. Phanvilay (NUOL, Vientiane), S. Sengchanthavong, S. Southichack, T. Homsaysombath, and Kieusomphone Thanabuaosy (Khammouane), M. de Koning and J. Foppes (Hin Nam No Project, Laos) for supporting our field research in Laos. Export of collected specimens was done due to the export permit Number 1307/14 signed by the CITES Management Authority of Lao PDR. V. Q. Luu thanks C. V. Pham, N. T. Nguyen, K. V. Phung and D. T. Bui (VFU, Hanoi) for supporting his work. We thank E. Sterling (New York) and K. Koy (Berkeley) for providing the map, T. Calame (Vientiane) for providing photographs and assistance in the field. Many thanks to O. Pauwels (Brussels) for providing comments on a previous version of the manuscript. Field work was funded by Cologne Zoo (Germany), Rufford Foundation (England), and Idea Wild (United States) for V. Q. Luu. Research of V. Q. Luu in Germany is funded by the Ministry of Education and Training of Vietnam (MOET, Project 911) and the German Academic Exchange Service (DAAD).

## REFERENCES

Bain R.H., Hurley M. 2011. A biogeographic synthesis of the amphibians and reptiles of Indochina. *Bulletin of the American Museum of Natural History* 360: 138 pp.

- Frost D. R. 2016. Amphibian Species of the World: An Online Reference, Version 6.0. American Museum of Natural History, New York, USA. Retrieved from <http://research.amnh.org/herpetology/amphibia/index.html>
- Glaw F., Vences M. 2007. A field guide to the amphibians and reptiles of Madagascar (p. 70). *Third Edition. Frosch Verlag.*
- Luu V.Q., Calame T., Nguyen T.Q., Ohler A., Bonkowski M., Ziegler T. 2014. First records of *Gracixalus supercornutus* (Orlov, Ho & Nguyen, 2004) and *Rhacophorus maximus* Günther, 1858 from Laos. *Herpetology Notes* 7: 419-423.
- Matsui M., Khonsue W., Panha S., Eto K. 2015. A New Tree Frog of the Genus *Gracixalus* from Thailand (Amphibia: Rhacophoridae). *Zoological Science* 32(2): 204-210.
- Nguyen T.Q., Hendrix R., Böhme W., Vu T.N., Ziegler T. 2008. A new species of the genus *Philautus* (Amphibia: Anura: Rhacophoridae) from the Truong Son Range, Quang Binh Province, central Vietnam. *Zootaxa* 13: 1-13.
- Nguyen T.Q., Phung T.M., Schneider N., Botov A., Tran D.T.A., Ziegler T. 2014. New records of amphibians and reptiles from southern Vietnam. *Bonn zoological Bulletin* 63(2): 148-156.
- Orlov N.L., Ho C.T., Nguyen T.Q. 2004. A new species of the genus *Philautus* from central Vietnam (Anura: Rhacophoridae). *Russian Journal of Herpetology* 11: 51-64.
- Palumbi S., Martin A., Romano S., McMillan W.O., Stice L., Grabowski G. 1991. The simple fool's guide to PCR, version 2.0. *Honolulu: Department of Zoology and Kewalo Marine Laboratory, University of Hawaii.*
- Rowley J.J.L., Dau V.Q., Nguyen T.T., Cao T.T., Nguyen S.V. 2011. A new species of *Gracixalus* (Anura: Rhacophoridae) with a hyperextended vocal repertoire from Vietnam. *Zootaxa* 38: 22-38.
- Rowley J.J.L., Tran D.T.A., Frankham G. J., Dekker A.H., Le D.T.T., Nguyen T.Q., Dau V. Q., Hoang H.D. 2015. Undiagnosed cryptic diversity in small, microendemic frogs (Lepidolalax) from the Central Highlands of Vietnam. *PloS One* 10(5): e0128382.
- Sterling E.J., Hurley M.M., Le M.D. 2006. Vietnam - A Natural History. *New Haven and London, Yale University Press*, 423 pp.
- Stuart B.L. 1999. Amphibians and Reptiles. In: Duckworth J. W., Salter R. E., Khounboline K.: A Wildlife and Habitat Survey of Hin Namno National Biodiversity Conservation Area and Adjacent Areas, Khammouane Province, Lao PDR. *Vientiane, Lao PDR: IUCN*, 124 pp.
- Teynié A., Nguyen T.Q., Lorvelec O., Lottier A., David P. 2014. Amphibiens et Reptiles du Laos: nouvelles données nationales et provinciales. *Bulletin de la Société Herpétologique de France* 151(3): 21-52.
- Timmins R.J., Trinh C.V. 2001. An assessment of the conservation importance of the Huong Son (Annamite) Forest, Ha Tinh Province, Vietnam, based on the results of a field survey for large mammals and birds. *Center for Biodiversity and Conservation, American Museum of Natural History, New York, USA*, 95 pp.