

SEVERE PARASITISM IN AN OPOSSUM

Authors: NETTLES, VICTOR F., PRESTWOOD, ANNIE K., and DAVIDSON, WILLIAM R.

Source: Journal of Wildlife Diseases, 11(3) : 419-420

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-11.3.419>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SEVERE PARASITISM IN AN OPOSSUM

VICTOR F. NETTLES, ANNIE K. PRESTWOOD and WILLIAM R. DAVIDSON,
Southeastern Cooperative Wildlife Disease Study, Department of Parasitology,
College of Veterinary Medicine, University of Georgia, Athens 30602, U.S.A.

Abstract: Chronic debilitation and anemia were observed in a free-living opossum (*Didelphis marsupialis*) heavily parasitized by *Physaloptera turgida*, *Brachylaima virginianum*, and *Cruzia americana*. Chronic interstitial pneumonia associated with *Capillaria aerophila* and a metastrongyloid nematode also was present.

HISTORY

In February, 1973, an opossum was observed repeatedly in a farmyard in Jackson County, Georgia. The animal was emaciated, weak in the hindquarters, circled, and fell frequently. Ten days after the first observation, it was captured by hand and submitted for examination.

The opossum was an adult female weighing 2.27 kg. Prior to necropsy she was comatose and had extremely depressed respiratory and heart rates. Blood examination revealed a hematocrit of 28% and a total leukocyte count of 5600 cells/mm³ consisting of 49% lymphocytes, 45% neutrophils, 4% monocytes, and 2% eosinophils. Nucleated erythrocytes, leptocytes, polychromasia, and poikilocytosis were evident in blood films.

POST MORTEM FINDINGS

At necropsy severe dehydration and emaciation were noted. Body tissues were abnormally pale, and approximately 30 ml of clear fluid was present in the abdominal cavity. The liver was pale yellow and moderately fibrotic. Histologically, occasional foci of histiocytes and neutrophils were found in portal areas. The kidneys were abnormally dark

in the cortical region, a condition which was attributed to bile pigment in numerous tubules.

The stomach was distended with 220 stomach worms, *Physaloptera turgida* Rudolphi 1819, of which a large proportion (67%) were immature stages. Numerous pinpoint ulcers were present where worms were attached to the gastric mucosa. These ulcers penetrated the glandular mucosa to the submucosal level. Focal necrosis, mild hemorrhage, moderate numbers of eosinophils, and mononuclear inflammatory cells were present at the sites of parasite attachment. The intestinal content was dark brown and semifluid, in which particles of insect exoskeleton and mollusk shell were found. Three hundred and nine intestinal flukes, *Brachylaima virginianum* Dickerson 1930, and 1989 roundworms, *Cruzia americana* Maplestone 1930, were found in the intestine and cecum.

The lung parenchyma was firm and multifocal indurated lesions occurred on the pleural surface. Four *Capillaria aerophila* (Creplin, 1839) were found in the lungs, and there were numerous eggs in the pleural lesions. In addition, approximately 20 metastrongyloid nematodes were present. These helminths resemble those of the genus *Perostrongylus* Schlegel 1934, and studies are in

This study was supported by an appropriation from the Congress of the United States. Funds were administered and research coordinated under the Federal Aid in Wildlife Restoration Act (50 Stat. 917) and through Contract No. 14-16-0008-708, Fish and Wildlife Service, U.S. Department of the Interior.

progress to further identify this nematode. Histologic examination of the lung revealed diffuse interstitial pneumonia characterized by smooth muscle hyperplasia, alveolar wall fibrosis, and focal accumulations of histiocytes.

DISCUSSION

Physaloptera embed in the gastric mucosa where they cause focal gastric ulcers^{7,8} although recent reports^{4,5} indicate that they may not feed on host tissue. Perforation of the stomach wall¹ and clinical illness characterized by profuse diarrhea, weight loss, partial gastric occlusion and hemorrhagic enteritis⁸ have

been associated with *P. turgida* infection in opossums. The intensity of infection in this animal exceeds previous reports,^{2,6} and the gastric ulcers associated with the attachment sites probably caused chronic blood loss which resulted in anemia.

Pathologic processes were not previously associated with slightly greater numbers of *B. virginianum* or *C. americana* than those reported herein,⁹ but such large numbers of helminths undoubtedly interfere with host nutrition. Additionally, pulmonary nematodes produced lung lesions that were another stress on the host. The cumulative effects of all parasites present were considered to be the cause of chronic debilitation observed in this opossum.

LITERATURE CITED

1. FELDMAN, D. B., J. A. MOORE, M. W. HARRIS and J. L. SELF. 1972. Characteristics of common helminths of the Virginia opossum (*Didelphis virginiana*) from North Carolina. *Lab. Anim. Sci.* 22: 183-189.
2. HALEY, J. S. 1938. Parasites in some wild fur bearers. *Vet. Med.* 33: 291.
3. KRUPP, J. H. 1962. Treatment of opossums with *Physaloptera* infections. *J. Am. vet. med. Ass.* 141: 369-370.
4. LINCOLN, R. C. 1972. Studies on the biology of *Physaloptera maxillaris* Molin (Nematoda: Physalopteroidea) a parasite in the stomach of skunk *Mephitis mephitis* Schreber. M.S. Thesis, The Univ. of Guelph.
5. LINCOLN, R. C. and R. C. ANDERSON. 1973. The relationship of *Physaloptera maxillaris* (Nematoda: Physalopteroidea) to skunk (*Mephitis mephitis*). *Can. J. Zool.* 51: 437-441.
6. MILLER, G. C. and R. HARKEMA. 1970. Helminths of the opossum (*Didelphis virginiana*) in North Carolina. *Proc. Helminth. Soc. Wash.* 37: 36-39.
7. SCHELL, S. C. 1952. Studies on the life cycle of *Physaloptera hispida* Schell (Nematoda: Spiruriodea) a parasite of the cotton rat (*Sigmodon hispidus littoralis* Chapman). *J. Parasit.* 38: 462-472.
8. SOULSBY, E. J. L. 1968. *Helminths, Arthropods and Protozoa of Domesticated Animals* (Mönnig). 6th ed. The Williams and Wilkins Co., Baltimore, Md.

Received for publication 29 January 1975