

NEW HOST AND DISTRIBUTION RECORDS OF ACANTHOCEPHALA FROM NORTH AMERICAN BIRDS

Authors: SCHMIDT, G. D., and NEILAND, K. A.

Source: Bulletin of the Wildlife Disease Association, 2(3) : 78

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-2.3.78>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

**NEW HOST AND DISTRIBUTION
RECORDS OF ACANTHOCEPHALA
FROM NORTH AMERICAN BIRDS**

During the past few years the authors have surveyed the helminth parasites of birds in several localities of the United States. This paper reports some host and

distribution records of *Acanthocephala* which appear to be new. New host and state records are indicated by an asterisk in the following table. The worms are retained in the authors' collections.

G. D. SCHMIDT
K. A. NEILAND

TABLE 1. *Some records of Acanthocephala from birds*

Parasite	Host and State
Polymorphidae	
<i>Polymorphus trochus</i>	<i>Fulica americana</i>
Van Cleave, 1945	Montana*
<i>Polymorphus</i> sp.	<i>Ixoreus naevius</i> *
	Oregon
<i>Arhythmorhynchus longicollis</i>	<i>Larus delawarensis</i> *
(Villot, 1875)	Oregon*
Plagiorhynchidae	
<i>Plagiorhynchus formosus</i>	<i>Ixoreus naevius</i> *
	Oregon*, Washington
Van Cleave 1918	<i>Colaptes cafer</i>
	Washington
<i>Plagiorhynchus paulus</i>	<i>Passerella iliaca</i>
	Oregon*
Van Cleave and Williams, 1951	<i>Ixoreus naevius</i> *
	Washington*
Centrorhynchidae	
<i>Sphaerirostris lancea</i>	<i>Ixoreus naevius</i>
	Washington*
Westrumb, 1821)	<i>Perisoreus obscurus</i> *
Golvan, 1960	Washington*
Gigantorhynchidae	
<i>Mediorhynchus grandis</i>	<i>Quiscalus quiscula</i>
Van Cleave, 1916	<i>Sternella neglecta</i> *
	Colorado*
<i>Mediorhynchus papillosus</i>	<i>Turdus migratorius</i> *
	Colorado*
Van Cleave, 1916	<i>Junco caniceps</i> *
	Colorado*
	<i>Penthestes gambeli</i> *
	Colorado*
<i>Mediorhynchus</i> sp.	<i>Sciurus noveboracensis</i> *
	New York

Dept. of Zoology
Colorado State College
Greeley, Colo.
Alaska Dept. Fish and Game
Fairbanks, Alaska, 24 February, 1966

IXODES TEXANUS IN EASTERN CANADA

The presence of *Ixodes texanus* Banks has been reported only twice in Ontario, all other records indicate that this tick is restricted in Canada to the west coast area (Gregson, 1956, Can. Dept. Agric. Publ. 930). Reports from the United States, however, record the distribution of *I. texanus* in Maryland, Delaware, Virginia, North Carolina, South Carolina, Georgia and Florida in the eastern area (Clifford, Anastos, and Elbe, 1961, Entomol. Soc. Amer., Miscell. Publ. 2, 215-237).

The author has made several collections of *I. texanus* from Ontario, the records of which are listed below.

Ontario.

South March, 6.II.64, ex *Me-
phitis mephitis* 1L
Cookstown, 14.II.64, ex *Pro-
cyon lotor* 1 ♀
Strathroy, 27.IV.64, ex *Pro-
cyon lotor* 30 ♀ ♀, 1 ♂, 6LL
Fletcher, 21.V.64, ex *Procyon
lotor* 1 ♀
Mactier, 5.VI.64, ex *Procyon
lotor* 3 ♀ ♀