

## **Can Androgen Deficiency Promote an Outbreak of Psoroptic Mange Mites in Male Deer?**

Author: Bubenik, George A.

Source: Journal of Wildlife Diseases, 25(4) : 639-642

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-25.4.639>

---

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](http://www.bioone.org/terms-of-use).

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## Can Androgen Deficiency Promote an Outbreak of Psoroptic Mange Mites in Male Deer?

George A. Bubenik, Department of Zoology, University of Guelph, Guelph, Ontario, Canada N1G 2W1

**ABSTRACT:** Repeated outbreaks of infection by psoroptic mange mites (*Psoroptes equi* var. *ovis*) affecting most regions of the body and legs were observed in several male white-tailed deer (*Odocoileus virginianus*) suffering from hypoandrogenism (e.g., castrates, animals treated with antiandrogens or postprime). The massive infection was characterized by a severe alopecia and skin inflammation and began usually in early winter. One or two spray treatments of a 1% solution of Lindane was usually sufficient to eliminate the clinical signs of the disease and to restore a healthy hair coat. Neither healthy male or female deer have ever exhibited any external signs of this disease nor has the parasite been detected in the scrapings of their skin. We propose a possible relationship between the hormonal status of these animals and their resistance to this parasitic mite infection.

**Key words:** *Psoroptes equi* var. *ovis*, mange mite, white-tailed deer, *Odocoileus virginianus*, male, androgens, immunology, resistance.

Infections with psoroptic mange mites has been described in domestic livestock as well as in wild animals (Sweatman, 1971). *Psoroptes cervinus* and *Psoroptes ovis* have been observed on bighorn sheep (*Ovis canadensis*) (Sweatman, 1971; Lange et al., 1980) and *Psoroptes cervinus* has been reported in wapiti (*Cervus elaphus canadensis*) (Sweatman, 1971; Thorne et al., 1982). Infection with *Psoroptes cuniculi* has been observed in wild and captive populations of white-tailed deer (*Odocoileus virginianus*) and mule deer (*Odocoileus hemionus*) (Roberts et al., 1970; Rollor et al., 1978; Strickland et al., 1981; Schmitt et al., 1982).

The clinical signs of disease caused by mites are depilation, exudation and pruritus. The feeding activity of the parasite causes the superficial layers of epidermis to break up which induces irritation and inflammation. This is accompanied by exudation of lymph and serous fluid that form layers of crusts and scabs. The acutely in-

fectected skin is red and usually loses hair over the affected areas (Sweatman, 1971). A secondary bacterial infection often causes a purulent exudate from the skin (Thorne et al., 1982).

In the wapiti, infection was observed on the neck, body and the upper legs (Sweatman, 1971); in white-tailed deer and mule deer *Psoroptes* spp. was reported only around the pedicle and in the deep regions of the auditory canal, but not in other dermal parts (Roberts et al., 1970; Kellogg et al., 1971; Strickland et al., 1981; Schmitt et al., 1982).

Because the disease erupts almost exclusively in winter (Sweatman, 1971), massive loss of hair can cause death of the host by exposure to cold (Honest and Winter, 1956; G. A. Bubenik, pers. obs.). On the other hand, ear infection, which produces a greenish-brown waxy material plugging an ear canal, can cause a pyogenic otitis (Rollor et al., 1978). Transmission of psoroptic mites by a direct body contact or from common rubbing posts has been reported by Sweatman (1971). However, an indirect transmission of mites deposited on vehicles and people is also being considered possible (Thorne et al., 1982).

During the last 10 yr, numerous infections of *Psoroptes equi* var. *ovis* have been observed in captive adult white-tailed deer bucks born and kept at the University of Guelph Deer Research Station in Cruikston Park near Cambridge (Ontario, Canada). The species of mites was identified by K. W. Wu from Agriculture Canada (Biosystematic Research Center, Ottawa, Ontario, Canada K1A 0C6). All male deer were housed individually in 5 × 30 m pens which were separated by a wall made of particle board; these were attached 20 cm from the ground to cedar posts, so nasal

contact between animals was possible. Females and fawns were kept in one large breeding enclosure (1 ha) adjacent to individual pens of males.

In contrast to previous reports describing the psoroptic mites on white-tailed deer (Roberts et al., 1970; Kellogg et al., 1971; Strickland et al., 1981), diseased skin was observed not only around the pedicle and in the auditory canal, but also on other parts of the body and legs. Signs of infection usually started in the skin right below the antler coronet (burr). This area has a higher humidity and warmer temperature than skin in other parts of the head. From there the mites spread anteriorly into frontal and nasal areas, laterally into the cheek regions and the auditory canal and posteriorly into the skin of the neck and the shoulders. Secondary infection (possibly the result of licking) developed later in the groin region from where it spread into abdominal areas.

This unusually massive infection of body areas, not described previously in white-tailed deer, was observed only in bucks with impaired production and action of androgens. The first cases were observed in two adult bucks treated with an antiandrogen cyproterone acetate (CA) (Schering AG., D-1000, Berlin 65, Federal Republic of Germany) which blocks the central as well as the peripheral action of androgens (Bubenik et al., 1975). Several years later, two other bucks treated with CA (Bubenik et al., 1987) developed a natural infection with psoroptic mites. After the termination of the treatment with CA these animals apparently became resistant again to psoroptic mites. In addition to the CA-treated animals, two castrated white-tailed deer used in several endocrine studies (Bubenik, 1983; Morris and Bubenik, 1983) were repeatedly infected in each of five consecutive winters.

Finally, two postprime bucks (aged at 10-, and at 11- to 12-yr-old, respectively), whose testosterone levels in the rutting season were well below the average concentration (Bubenik and Schams, 1986) had

to be treated for infection with psoroptic mites. Alternatively, signs of this skin disease were not observed in two castrated bucks which were treated with testosterone in the fall in order to polish and cast overgrown antlers (Bubenik, 1983). Otherwise, healthy skin in castrates occurred only during the first winter after orchiectomy which was performed late in September.

In contrast to these cases, clinical signs of the disease caused by this mite infection were not observed in any female white-tailed deer; infection was not detected in bucks treated in September with CA for only 2 to 3 wk.

In non-castrated bucks a very mild infection (usually just in the deeper parts of the auditory canal) was observed only in deer that were under severe stress because of bacterial or viral disease (Bubenik and Brownlee, 1987) or a repeated long-term experimental blood sampling (Bubenik, 1986). Healthy, vigorous bucks were never affected, despite being exposed to nasal contact with deer that were heavily infested with mites.

The exact mode of initial transfer of the parasite to our deer was never determined. Domestic livestock (cows and sheep) kept in the nearby farm might have been a source of the mites, but other origins of infection also must be considered because the experimental transfer between wild and domestic ruminants has not been successful (Hepworth and Thomas, 1962).

The initial signs of the skin disease were usually first observed in late December. Cold and humid weather with temperatures near 5 C appeared to facilitate the spread of the disease to new sides on the body. Alternatively, dry, frosty conditions seemed to reduce the spread of mites (G. A. Bubenik, pers. obs.).

As soon as the infection became obvious in the deer (e.g., alopecia, erythema and exudate) a vigorous treatment was initiated. The obvious sign of hairless patches on the skin indicates that the infection is already established in much larger sur-

rounding areas. After the first detection of clinical signs of infection (usually using binoculars) the infected buck was tranquilized with xylazine hydrochloride (Rompun, Haver-Lockhart, Bayvet Division, Mississauga, Ontario, Canada L4K 1G4) (Bubenik, 1982) and affected areas of skin were treated with 1% solution of Lindane (Pfizer, Dorval, Quebec, Canada H9R 4V2) applied first from a spray bottle and then rubbed in with rubber gloves. Usually twice as large an area was treated as appeared macroscopically to be infected.

In most cases two treatments 2 or 3 wk apart were sufficient to restore healthy skin for the rest of the year. Because of the topical use, concentration of Lindane was much higher than recommended for the whole body dip. The skin of treated deer healed relatively quickly and a new hair growth was observed about 2 weeks after the treatment. However in some severe cases, where the detection was made late, up to four treatments were necessary to eliminate the signs of the disease as determined by a careful examination of the affected areas using a magnifying glass. However, as no microscopic investigation of the skin has been performed, mites might have survived in inaccessible areas (such as the deeper parts of the auditory canal).

That sex hormones can influence the function of the immune system has been well established. Dehydroepiandrosterone was found to have anti-autoimmune properties (Schwartz et al., 1984) and estradiol ( $E_2$ ) was reported to depress the function of the thymic lymphocytes (Grossmann et al., 1983). Conversely, dihydrotestosterone (DHT) possibly enhances function of thymic lymphocytes (Grossman and Roselle, 1983). In addition, specific  $E_2$  receptors have been detected in the reticuloendothelial cell of the thymus (Grossman and Roselle, 1983).

In addition to the effect of androgens mediated by the thymus, male sexual hormones also are known as anabolic hormones which improve the utilization of

nutritional resources (Turner and Bagnara, 1974). Decrease of androgen levels may then affect the immunoresistance by a reduction of certain vital metabolic pathways.

In view of these findings, it can be speculated that the severe psoroptic infection in some captive male white-tailed deer may be a result of their hypoandrogenesis that caused either a decrease in immunoresponsiveness of the skin or in alteration of the skin ecology favoring the proliferation of the mites.

The relationship between the levels of sexual hormones and the mite infection in wild animals deserves further study. Presently, there is no explanation why female deer never exhibited any external signs of the disease (such as the hairless patches or red skin with exudate) or how the thymic stimulation could influence the resistance of the deer against these skin parasites. Further studies exploring whether "an androgen supplement" could prevent spread of this infection in a stressed population or if androgen treatment may cure already heavily infected bucks are warranted.

I would like to thank Roy C. Anderson and Tony B. Bubenik for their kind review of this manuscript as well as Ute Strelive and King W. Wu for the identification of the mite.

#### LITERATURE CITED

- BUBENIK, G. A. 1982. Chemical immobilization of captive white-tailed deer and the use of automatic blood samplers. In *Chemical immobilization of North American wildlife*, L. Nielsen et al. (eds.). Wisconsin Humane Society, Milwaukee, Wisconsin, pp. 335-354.
- . 1983. The endocrine regulation of the antler cycle. In *Antler development in Cervidae*, R. D. Brown (ed.). Caesar Kleberg Wildlife Research Institute, Kingsville, Texas, pp. 73-107.
- . 1986. Regulation of seasonal endocrine rhythms in male boreal cervids. In *Endocrine regulation as adaptive mechanism to the environment*, I. Assenmacher and J. Boissin (eds.). Centre National de la Recherche Scientifique—Centre d'Etudes Biologique des Animaux Sauvages, France, pp. 461-474.
- , AND L. BROWNLEE. 1987. Assessing health of male white-tailed deer by white blood cell

- count. *The Journal of Wildlife Management* 51: 57-58.
- , A. B. BUBENIK, G. M. BROWN, AND D. A. WILSON. 1975. The role of sex hormones in the growth of antler bone tissues. I. Endocrine and metabolic effect of antiandrogen therapy. *Journal of Experimental Zoology* 194: 349-358.
- , AND D. SCHAMS. 1986. Relationship of age to seasonal levels of LH, FSH, prolactin and testosterone in male white-tailed deer. *Comparative Biochemistry and Physiology* 83A: 179-183.
- , D. SCHAMS, AND G. COENEN. 1987. The effect of artificial periodicity and antiandrogen treatment on the antler growth and plasma levels of LH, FSH, testosterone, prolactin and alkaline phosphatase in the male white-tailed deer. *Comparative Biochemistry and Physiology* 87A: 551-559.
- GROSSMAN, C. J., AND G. A. ROSELLE. 1983. The interrelationship of the HPG-thymic axis and immune system regulation. *Journal of Steroid Biochemistry* 19: 461-467.
- , L. J. SHOLITON, AND J. A. HELMWORTH. 1983. Characteristics of the cytoplasmic and nuclear dihydrotestosterone receptors of human thymic tissue. *Steroids* 42: 11-22.
- HEPWORTH, W. G., AND G. M. THOMAS. 1962. Attempts to transfer psoroptic mites from elk to cattle and sheep. *Journal of American Medical Association* 140: 689-690.
- HONESS, R. F., AND K. B. WINTER. 1956. Diseases of wildlife in Wyoming. Bull. 9, State of Wyoming Game and Fish Commission, Cheyenne, Wyoming, pp. 1-279.
- KELLOGG, F. E., T. P. KISTNER, R. K. STRICKLAND, AND R. R. GERRISH. 1971. Arthropod parasites collected from white-tailed deer. *Journal of Medical Entomology* 8: 495-498.
- LANGE, R. E., A. V. SANDOVAL, AND W. P. MELENEY. 1980. Psoroptic scabies in bighorn sheep (*Ovis canadensis mexicana*) in New Mexico. *Journal of Wildlife Diseases* 16: 77-82.
- MORRIS, J. M., AND G. A. BUBENIK. 1983. The effects of androgens on the development of antler bone. In *Antler development in Cervidae*, R. D. Brown (ed.). Caesar Kleberg Wildlife Research Institute, Kingsville, Texas, pp. 123-141.
- ROBERTS, I. H., W. P. MELENEY, AND R. E. PILLMORE. 1970. Ear scab-mites, *Psoroptes cuniculi* (Acarina: Psoroptidae), in captive mule deer. *Journal of Parasitology* 56: 1039-1040.
- ROLLOR, E. A., V. F. NETTLES, W. R. DAVIDSON, AND R. R. GERISH. 1978. Otitis media caused by *Psoroptes cuniculi* in white-tailed deer. *Journal of American Veterinary Medical Association* 173: 1242-1243.
- SCHMITT, S. M., T. M. COOLEY, AND T. W. SCHILLHORN-VAN-VEEN. 1982. *Psoroptes cuniculi* in captive white-tailed deer in Michigan. *Journal of Wildlife Diseases* 18: 349-351.
- SCHWARTZ, A. G., J. W. NYCE, AND R. H. TANNEN. 1984. Inhibition of tumorigenesis and autoimmune development in mice by dehydroepiandrosterone. In *Altered endocrine status during aging*. A. R. Liss, New York, New York, pp. 177-184.
- STRICKLAND, R. K., R. R. GERISH, AND J. S. SMITH. 1981. Arthropods. In *Diseases and parasite of white-tailed deer*, W. R. Davidson, E. A. Hayes, V. F. Nettles, and F. E. Kellogg (eds.). Tall Timber Research Station, Tallahassee, Florida, pp. 363-389.
- SWEATMAN, G. K. 1971. Miter and pentasomes. In *Parasitic diseases of wild mammals*, J. W. Davis and R. C. Anderson (eds.). Iowa State University Press, Ames, Iowa, pp. 3-64.
- THORNE, E. T., N. KINGSTON, W. R. JOLLEY, AND R. C. BERGSTROM (editors). 1982. *Diseases of wildlife in Wyoming*, 2nd ed. Wyoming Game and Fish Commission, Cheyenne, Wyoming, pp. 1-353.
- TURNER, C. D., AND J. C. BAGNARA. 1974. *General endocrinology*, 6th ed. W. B. Saunders, Philadelphia, Pennsylvania, pp. 1-596.

Received for publication 14 September 1988.