

## A RAPID GEL DIFFUSION PRECIPITIN TEST FOR CONTAGIOUS BOVINE PLEUROPNEUMONIA 1

Author: SHIFRINE, M.

Source: Bulletin of the Wildlife Disease Association, 3(1): 36

Published By: Wildlife Disease Association

URL: https://doi.org/10.7589/0090-3558-3.1.36

The BioOne Digital Library (<u>https://bioone.org/</u>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<u>https://bioone.org/subscribe</u>), the BioOne Complete Archive (<u>https://bioone.org/archive</u>), and the BioOne eBooks program offerings ESA eBook Collection (<u>https://bioone.org/esa-ebooks</u>) and CSIRO Publishing BioSelect Collection (<u>https://bioone.org/csiro-ebooks</u>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at <u>www.bioone.org/terms-of-use</u>.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

bitats. Since it is of zoological interest to have a knowledge of the parasite fauna of wild mammals in any region, a survey of the helminth parasites of the deer mouse may provide insight to parasitism in other rodent populations as well.

Eighty-eight deer mice collected from Cache la Poudre Canyon, Larimer County, northern Colorado, during the summer of 1964 were examined for helminth parasites. Four species of nematodes were found. The names of the parasites, number of hosts infected, per cent infection and location within the host are given below.

- 1. *Physaloptera sp.* (juveniles in tissues); 2; 2.3 per cent infected; stomach (a new host record).
- 2. Mastophorus numidica (Seurat, 1915); 33; 37.5 per cent infected; stomach.
- 3. Rictularia coloradensis Hall, 1916; 2; 2.3 per cent infected; small intestine.
- 4. Trichurus perognatha Chandler, 1945; 5; 5.7 per cent infected; caecum.

W. G. DYER and O. W. OLSEN

Dept. Biology, Minot State College, Minot, North Dakota, and Dept. Zoology, Colorado State Univ., Fort Collins, Colo. 3 August, 1966

## A RAPID GEL DIFFUSION PRECIPITIN TEST FOR CONTAGIOUS BOVINE PLEUROPNEUMONIA<sup>1</sup>

During a recent outbreak of contagious bovine pleuropneumonia (CBPP) in Kenya, three tests used for diagnosis of this disease were re-evaluated. It was found that the slide agglutination serum test (SAST) in conjunction with agar gel diffusion test (AGT) for circulating antigen, detected 100% of cattle with CBPP whereas the complement fixation test failed to detect 6% of these cases (Shifrine and Gourlay, 1966, Bull. Epiz. Drs. Afr., in press).

Both AGT and SAST require little equipment and can readily be performed in the field, which is of great advantage under the present conditions in East Africa. The disadvantage of the AGT is that it requires 24 hours or more before the results of the test can be observed. Various modifications, therefore, were tried in an attempt to obtain results within a few hours. The modification that was developed is based on the quantitative gel assay of Feinberg (Feinberg, J. G., 1959, Immunology, 2: 346), using prediffused plates.

The AGT was done in petri dishes as previously described (Gourlay, R. N., 1964, Vet. Sci., 5: 473). Wells were cut in the agar with a cutter which consisted of a center well (11 mm in diameter) circled by 6 wells (6 mm in diameter); the distance between the center and peripheral wells was 3 mm. Hyperimmune bovine antisera against Mycoplasma mycoides, the causative organism of CBPP, were placed in the center well and the plate left for o ne to three days at room temperature. Sera from 73 cattle suspected of having CBPP were placed in the peripheral wells. Precipitin lines appeared after 4 to 6 hours using prediffused plates. When the conventional AGT (both reagents placed in the wells at the same time) was used, precipitin lines appeared after one to two days. Moreover, sera from 2 infected animals produced a negative test in the conventional AGT and positive test using the prediffused plates.

The use of the modified AGT for circulating antigen in conjunction with the SAST facilitates rapid field evaluation of cattle suspected of having CBPP.

M. SHIFRINE<sup>2</sup>

East Africa Vet. Res. Org.

P.O. Box 32, Kikuyu, Kenya

31 October 1966

<sup>1</sup> The research described in this paper was partly financed by the United States of America Agency for International Development, under the terms of CCTA/AID Joint Project 16 for research on contagious bovine pleuropneumonia.

<sup>2</sup> Employed by the U.S. Department of Agriculture, Agricultural Research Service, Animal Disease & Parasite Research Division, Plum Island Animal Disease Laboratory, Greenport, Long Island, New York.

36