

Postpartum Immobilization of Adult Female Moose using Xylazine, Ketamine and Yohimbine Hydrochlorides

Authors: Garner, Dale L., and Addison, Edward M.

Source: Journal of Wildlife Diseases, 30(1) : 123-125

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-30.1.123>

The BioOne Digital Library (<https://bioone.org/>) provides worldwide distribution for more than 580 journals and eBooks from BioOne's community of over 150 nonprofit societies, research institutions, and university presses in the biological, ecological, and environmental sciences. The BioOne Digital Library encompasses the flagship aggregation BioOne Complete (<https://bioone.org/subscribe>), the BioOne Complete Archive (<https://bioone.org/archive>), and the BioOne eBooks program offerings ESA eBook Collection (<https://bioone.org/esa-ebooks>) and CSIRO Publishing BioSelect Collection (<https://bioone.org/csiro-ebooks>).

Your use of this PDF, the BioOne Digital Library, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Digital Library content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne is an innovative nonprofit that sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Postpartum Immobilization of Adult Female Moose using Xylazine, Ketamine and Yohimbine Hydrochlorides

Dale L. Garner,¹ and Edward M. Addison,² ¹ Faculty of Environmental and Forest Biology, College of Environmental Science and Forestry, State University of New York, Syracuse, New York 13210, USA; ² Ontario Ministry of Natural Resources, Wildlife Research Section, P.O. Box 5000, Maple, Ontario L6A 1S9, Canada

ABSTRACT: Twenty-two free-ranging adult female moose (*Alces alces*) were immobilized with a 1:4 mixture of xylazine hydrochloride (XH) and ketamine hydrochloride (KH). Mean (SD) dosages/animal for XH and KH were 419 (148) and 1565 (433) mg, respectively. Mean (SD) induction time was 18.4 (9.7) minutes. Reversal with yohimbine hydrochloride using a mean dosage of 83 mg/animal resulted in a mean (SD) recovery time of 22.8 (28.5) minutes.

Key words: Immobilization, postpartum, moose, *Alces alces*, xylazine hydrochloride, ketamine hydrochloride, yohimbine hydrochloride.

A wide variety of wildlife species have been immobilized successfully using xylazine hydrochloride (XH) in combination with ketamine hydrochloride (KH), including members of the family Cervidae. Golightly and Hofstra (1989) reported success in immobilizing Roosevelt elk (*Cervus elaphus roosevelti*) using XH-KH reversed with yohimbine hydrochloride (YH). This combination also has been effective on mule deer (*Odocoileus hemionus*) (Jessup et al., 1983) and white-tailed deer (*O. virginianus*) (Mech et al., 1985).

Schwab et al. (1984) reported the use of XH-KH for immobilizing moose (*Alces alces*) during winter. However, they did not recommend its use for future studies because of the variability in animal response. We used XH-KH for immobilizing female moose following parturition and report the application of XH-KH for immobilizing postpartum female moose with YH used as an antagonist.

During each spring from 1990 to 1992, adult female moose were captured as part of an ongoing demographic study in Algonquin Provincial Park, Ontario, Canada (45°39'N, 78°39'W). The Park lies within south-central Ontario in which many lakes

with islands and peninsulas are present. Islands and peninsulas in the Park are known calving areas for female moose (Addison et al., 1990).

From mid-May to early June, crews searched islands and peninsulas for females with newborn calves by walking abreast, maintaining contact with one another visually, by voice, or through the use of portable radios. Once located, adult females were immobilized with a 1:4 (100 mg/ml) mixture of XH-(Rompun®, Bayvet Division Chemagro, Ltd., Etobicoke, Ontario) KH (Rogarsetic®, Rogar/STB Inc., London, Ontario) delivered intramuscularly (IM) from Cap-Chur equipment (Palmer Chemical and Equipment Company, Douglasville, Georgia, USA) or pole syringes. Due to the dilutions of the immobilizing drugs, all moose ($n = 22$) required >1 injection (range = 2 to 4; $\bar{x} = 2.5$).

Once immobilized, all moose received an eartag and a mortality sensing radio collar (Advanced Telemetry Systems Inc., Isanti, Minnesota, USA). Calves were captured by hand and received an eartag and a break-away mortality sensing radio collar (Lotek Engineering Inc., Newmarket, Ontario, Canada). Adult females in deep narcosis were given 5 mg/ml YH (Antagonil® Janssen Pharmaceutical, Mississauga, Ontario, Canada) intravenously and IM as an antagonist.

Drug doses are reported as mg administered/animal rather than mg/kg of body weight because of the difficulty of weighing moose in the field. We assumed live weights of adult females were similar to those reported by Quinn and Aho (1989). They reported a mean live weight of 435 kg for adult female moose in Algonquin

TABLE 1. Drug dosages and induction and recovery times of immobilized postpartum female moose in Algonquin Provincial Park, Ontario, Canada, 1990 to 1992.

	<i>n</i> ^a	Mean (SD)	Range
Xylazine hydrochloride (mg/animal)	22	418.6 (148.1)	280.0–820.0
Ketamine hydrochloride (mg/animal)	22	1,564.6 (432.9)	1,120.0–2,560.0
Induction time (min)	14	18.4 (9.7)	4.0–39.0
Yohimbine hydrochloride (mg/animal)	20	82.9 (11.0)	50.0–112.0
Recovery time ^b (min)	9	22.8 (28.5)	1.0–71.0

^a Number of moose tested.^b Time to standing.

Park. Induction time was recorded as the time (in min) from first injection until sternal or lateral recumbency was attained. Recovery time was the interval from injection of YH to standing.

Upon reversal, adult females were visually monitored until stable. All adult females and their calves were released at the capture site. After release, animals were monitored on a daily basis to the end of June. A weighted regression (SAS Institute Inc., 1990) was used to test for an effect of day of month captured on the amount of XH and KH used to achieve immobilization. Regression weights were computed as the inverse of the standard deviation (SD) of the mean dose of XH and KH per day. Days where only one moose was immobilized were not included in the analysis. We tested the regression slopes (β_1) against the null hypothesis that $\beta_1 = 0$ using *t*-tests (SAS Institute Inc., 1990).

Twenty-two adult female moose were immobilized. Mean (SD) dosage/animal for XH was 419 (148) mg and mean (SD) dosage/animal for KH was 1565 (433) mg. Mean (SD) induction time was 18.4 (9.7) minutes. Reversal with YH resulted in a mean recovery time of 22.8 (28.5) min (Table 1). The amount of XH used to achieve immobilization was significantly ($\beta_1 = 18.1$, $t = 2.2$, $P = 0.08$) affected by day of the month. Similarly, the amount of KH used also was affected ($\beta_1 = 81.0$, $t = 3.7$, $P = 0.01$). In total, an addition of approximately 100 mg/day of XH-KH was required to achieve effective immobilization as the calving season progressed.

The XH-KH combination produced satisfactory results for immobilizing postpartum adult female moose in this study. In contrast, Schwab et al. (1984) reported that immobilization of moose using XH-KH was not satisfactory during winter. We believe the disparate results may be related to the different physical and physiological status of the moose. Postpartum moose would be at a low annual physical and physiological state. Drawbacks of using XH-KH in this application included the large amounts (>1 injection at the concentration used) of drug required for immobilization, and the wide variation in induction times between animals.

The quantity of XH-KH could have been reduced significantly by concentrating the mixture through lyophilization. Variation in induction time between animals appeared to be a function of excitement level of the animal, which we believe was influenced by pursuit time (i.e., animals with longer pursuit times had longer induction times), and possibly by the strength of the cow-calf bond.

Larsen and Gauthier (1989), using a carfentanil, fentanil, XH, hyaluronidase (CFXH) mixture, attributed low survival of calves to both short-term and long-term effects of the drugs. They reported that using CFXH prepartum changed the female's normal defense behavior, which resulted in decreased postpartum calf survival. There was no evidence in our study that immobilization with XH-KH influenced either cow attentiveness to their young or postpartum calf survival. How-

ever, one calf died from injuries sustained when its partially immobilized mother fell on it. No adult mortality occurred as a result of XH-KH immobilization.

The increase of approximately 100 mg of drug/day needed to achieve effective anesthesia in this study raises the possibility of a metabolic effect on induction time through the calving season. Metabolic rate in moose is influenced by plant phenology (Van Ballenberghe and Miquelle, 1990). Leafout in our study area occurs coincident with calving in mid- to late May. Moose, like other northern cervids, have an increase in metabolic rate in a summer as compared to winter (Regelin et al., 1985). We suggest that increased metabolic rate affected the amount of drug needed for anesthesia. Future studies could address this hypothesis to identify time specific factors producing more effective immobilization of cervids.

Golightly and Hofstra (1989) reported that injection of YH \geq 35 minutes post-induction for elk always resulted in successful reversal. However, in our study recovery of moose following injection of YH was variable regardless of the time administered. Nevertheless, we recommend its use when using XH-KH for immobilizing moose.

While carfentanil currently is considered the drug of choice for moose (Franzmann et al., 1987), our primary reasons for recommending the use of XH-KH are adequate response by postpartum cow moose and operator safety. These responses included total survival of immobilized animals, wide dosage tolerance, adequate induction times, effective level of anesthesia, no adverse side effects, and potential reversibility. We contend that XH-KH is a desirable immobilizer for moose under the circumstances similar to those reported herein.

Sincerest thanks to all the volunteers (too numerous to name) who assisted us during the 1990, 1991 and 1992 spring field seasons. The contributions of L. Bruce, M. Donnelly, A. Felski, E. Klupfel, R. Masters,

J. Millsbaugh, L. O'Connor, B. Seybold, and A. Silver are greatly appreciated. Special thanks to H. B. Underwood for reviewing the manuscript. Funding was provided by the North American Wildlife Research Foundation, the New York Legislature under the auspices of the Adirondack Wildlife Program and the Ontario Ministry of Natural Resources.

LITERATURE CITED

- ADDISON, E. M., J. D. SMITH, R. F. McLAUGHLIN, D. J. H. FRASER, AND D. G. JOACHIM. 1990. Calving sites of moose in central Ontario. *Alces* 26: 142-153.
- FRANZMANN, A. W., C. C. SCHWARTZ, AND D. C. JOHNSON. 1987. Anesthesia of moose for vasectomy using carfentanil/xylazine and reversal with naloxone/yohimbine. *Alces* 23: 221-226.
- GOLIGHTLY, R. T., JR., AND T. D. HOFSTRA. 1989. Immobilization of elk with a ketamine-xylazine mix and rapid reversal with yohimbine hydrochloride. *Wildlife Society Bulletin* 17: 53-58.
- JESSUP, D. A., W. E. CLARK, P. A. GULLETT, AND K. R. JONES. 1983. Immobilization of mule deer with ketamine and xylazine, and reversal of immobilization with yohimbine. *Journal of the American Veterinary Medical Association* 183: 1339-1340.
- LARSEN, D. G., AND D. A. GAUTHIER. 1989. Effects of capturing pregnant moose and calves on calf survivorship. *The Journal of Wildlife Management* 53: 564-567.
- MECH, L. D., G. D. DEL GIUDICE, P. D. KARNS, AND U. S. SEAL. 1985. Yohimbine hydrochloride as an antagonist to xylazine hydrochloride-ketamine hydrochloride immobilization of white-tailed deer. *Journal of Wildlife Diseases* 21: 405-410.
- QUINN, N. W. S., AND R. W. AHO. 1989. Whole weights of moose from Algonquin Park. *Alces* 25: 48-51.
- REGELIN, W. L., C. C. SCHWARTZ, AND A. W. FRANZMANN. 1985. Seasonal energy metabolism of adult moose. *The Journal of Wildlife Management* 49: 388-393.
- SAS INSTITUTE INC. 1990. SAS procedures guide, version 6, 3rd ed. SAS Institute Inc., Cary, North Carolina, 705 pp.
- SCHWAB, F. E., S. W. SCHWAB, AND M. D. PITT. 1984. Moose immobilization program in north-central British Columbia. *Alces* 20: 209-221.
- VAN BALLEMBERGHE, V., AND D. G. MIQUELLE. 1990. Activity of moose during spring and summer in interior Alaska. *The Journal of Wildlife Management* 54: 391-396.

Received for publication 11 March 1993.